

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 April 2002 (18.04.2002)


PCT

(10) International Publication Number
WO 02/31117 A2

- (51) International Patent Classification⁷: C12N (74) Agents: SERAFINI, Andrew, T. et al.; Townsend and Townsend and Crew LLP, 2 Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).
- (21) International Application Number: PCT/US01/32202
- (22) International Filing Date: 15 October 2001 (15.10.2001) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/687,837 13 October 2000 (13.10.2000) US
- (71) Applicants (*for all designated States except US*): ARBOR VITA CORPORATION [US/US]; 772 Lucerne Drive, Sunnyvale, CA 94085 (US). GARMAN, Jonathan, David [US/US]; 5084 McCoy Avenue, San Jose, CA 95130 (US). CANDIA, III, Albert, Frederick [US/US]; 789 Marsh Road, Menlo Park, CA 94025 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): LU, Peter, S. [US/US]; 99 East Middlefield Road, No. 29, Mountain View, CA 94043 (US).
- Published:
— *without international search report and to be republished upon receipt of that report*

[Continued on next page]

(54) Title: CLASP-2 TRANSMEMBRANE PROTEINS



(57) Abstract: The present invention relates to a cell surface molecule, designated cadherin-like asymmetry protein-2 ("CLASP-2"). In particular, it relates to CLASP-2 polynucleotides, polypeptides, fusion proteins, and antibodies. The invention also relates to methods of modulating an immune response by interfering with CLASP-2 function.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

CLASP-2 TRANSMEMBRANE PROTEINS

0.0 CROSS-REFERENCES TO RELATED APPLICATIONS

This application is a continuation-in-part application of U.S. Patent Application No. 09/547,276 filed April 11, 2000, which claims the benefit of U.S. Provisional Application Nos. 60/182,296 filed February 14, 2000, 60/176,195 filed January 14, 2000, 60/170,453 filed December 13, 1999, 60/162,498 filed October 29, 1999, 60/160,860 filed October 21, 1999, 60/134,118 filed May 14, 1999, 60/134,117 filed May 14, 1999, 60/134,114 filed May 14, 1999, and 60/129,171 filed April 14, 1999, the disclosures of which are incorporated by reference.

1.0 FIELD OF THE INVENTION

The present invention relates to molecules expressed in cells of the immune system. In particular, the invention relates to a transmembrane protein that contains certain classical cadherin characteristics.

2.0 BACKGROUND OF THE INVENTION

The generation of an immune response against an antigen is carried out by a number of distinct immune cell types that work in concert within the context of a particular antigen. The helper T cell (T_H) plays a pivotal role to coordinate two types of antigen-specific immune response; *i.e.*, cellular and humoral immune response. Recognition of antigen by T cells requires the formation of a specialized junction between the T cell and the antigen-presenting cell (APC) called the "immunological synapse" (Dustin, *et al.*, 1998, Cell 94: 667-677). The immune synapse orchestrates recruitment and exclusion of specific proteins from the contact area by an unknown mechanism and is thought to be initiated by T-cell antigen receptor (TCR) recognition of peptides bound to MHC molecules (antigen) (Monk, *et al.* 1998, Nature 395: 82). However, the low affinity of the TCR for antigen as well as limited number of ligands makes it unlikely that TCR: antigen interaction alone is sufficient to drive the formation of the immunological synapse (Matsui *et al.*, 1994, Proc. Natl. Acad. Sci. U.S.A. 91: 12861-12866).

Costimulatory molecules such as CD4, ICAM-1, LFA-1, CD28, CD2 have been proposed to stabilize the cell-cell contact (Dustin, *et al.*, 1999, Science 283: 649). However, since these molecules are recruited to the synapse after activation they cannot

account for the high specificity and avidity during the early phases of T-cell antigen recognition. Recent work demonstrated that a portion of the T cell surface at the leading edge is specialized to mediate the early phases of synapse formation (Negulescu, *et al.*, 1996, Immunity 4: 421-430). Such a specialization must be a pre-formed structure containing cell surface adhesion proteins (ectodomains) to augment TCR engagement and corresponding cytoplasmic portions (endodomains) to transduce signals and bind cytoskeleton to maintain structural/functional polarity.

The ectodomain of the pre-formed synapse or "immune gateway" was recently discovered and is created in part by CLASP-1 (U.S.S.N. 09/411,328, filed October 1, 1999; PCT/US99/22996). In addition to cadherin motifs, CLASP-1 also contains a CRK-SH3 binding domain, tyrosine phosphorylation sites, and coiled/coil domains suggesting direct interaction with cytoskeleton and regulation by adaptor molecules such as CRK. The *CLASP-1* transcript is present in lymphoid organs and neural tissue, and the protein is expressed by T and B lymphocytes and macrophages in the MOMA-1 subregion of the marginal zone of the spleen, an area known to be important in T: B cell interaction. CLASP-1 staining of individual T and B cells exhibits a preactivation structural polarity, being organized as a "ball" or "cap" structure in B cells, and forming a "ring", "ball" or "cap" structure in T cells. The placement of these structures is adjacent to the microtubule-organizing center ("MTOC"). CLASP-1 antibody staining indicates that CLASP-1 is at the interface of T-B cell conjugates that are fully committed to differentiation. Antibodies to the extracellular domain of CLASP-1 also block T-B cell conjugate formation and T cell activation.

3.0 SUMMARY OF THE INVENTION

The present invention relates to a cell surface molecule, a member of a new multigene-family designated cadherin-like asymmetry protein(s) ("CLASP(s)"). In particular, it relates to a polynucleotide comprising a coding sequence for CLASP-2, a polynucleotide that selectively hybridizes to the complement of a CLASP-2 coding sequence, expression vectors containing such polynucleotides, genetically-engineered host cells containing such polynucleotides, CLASP-2 polypeptides, CLASP-2 fusion proteins, therapeutic compositions, CLASP-2 domain mutants, antibodies specific for CLASP-2 polypeptides, methods for detecting the expression of CLASP-2, and methods of inhibiting an immune response by interfering with CLASP-2 function. A wide variety of uses are encompassed by the invention, including but not limited to, treatment of autoimmune

diseases and hypersensitivities, prevention of transplantation rejection responses, and augmentation of immune responsiveness in immunodeficiency states.

In one aspect, the invention provides an isolated CLASP-2 polynucleotide that is: (a) a polynucleotide that has the sequence of SEQ ID NO: 1, 3, 5 or 9; (b) a polynucleotide that hybridizes under stringent hybridization conditions to (a) and encodes a polypeptide having the sequence of SEQ ID NO: 2, 4, 6 or 10 or an allelic variant or homologue of a polypeptide having the sequence of SEQ ID NO: 2, 4, 6 or 10; or (c) a polynucleotide that hybridizes under stringent hybridization conditions to (a) and encodes a polypeptide with at least 25 contiguous residues of the polypeptide of SEQ ID NO: 2, 4, 6 or 10; or (d) a polynucleotide that hybridizes under stringent hybridization conditions to (a) and has at least 12 contiguous bases identical to or exactly complementary to SEQ ID NO: 1, 3, 5, or 9. In a related aspect, the invention provides a CLASP-2 polynucleotide wherein the polynucleotide encodes a polypeptide that binds to the PDZ domain of PSD95, DLG1 or neDLG. 2. In another related aspect, the invention provides a CLASP-2 polynucleotide wherein the polynucleotide encodes a polypeptide that has a binding affinity of at least 10^4 M^{-1} for binding PSD95, DLG1 or neDLG.

In one aspect, the invention provides a CLASP-2 polynucleotide that encodes a polypeptide having the full-length sequence of SEQ ID NO: 2, 4, 6, or 10 or the cDNA sequence encoded by the inserts of ATCC Deposit Nos: PTA-1562, PTA-1563 and PTA-1573.

In another aspect, the invention provides a CLASP-2 polynucleotide that encodes a polypeptide having the full-length sequence of Isoform 1, Isoform 2, or Isoform 3 (SEQ ID NO: _____) or the cDNA sequence encoded by the inserts of AVC-PD14 (ATCC Deposit No. _____) and AVC-PD19 (ATCC Deposit No. _____).

In another aspect, the invention further provides an isolated CLASP-2 polynucleotide comprising a nucleotide sequence that has at least 90% percent identity to SEQ ID NO: 1, 3, 5 or 9 as calculated using FASTA wherein said sequences are aligned so that highest order match between said sequences is obtained.

The invention further provides an isolated polypeptide comprising a nucleotide sequence that has at least 90% sequence identity to SEQ ID NO: 2, 4, 6 or 10 and is immunologically crossreactive with SEQ ID NO: 2, 4, 6 or 10 or shares a biological function with native CLASP-2.

The invention also provides vectors, such as expression vectors, comprising a polynucleotide sequence of the invention. In other embodiments, the invention provides host

cells or progeny of the host cells comprising a vector of the invention. In certain embodiments, the host cell is a eukaryote. In other embodiments, the expression vector comprises a CLASP-2 polynucleotide in which the nucleotide sequence of the polynucleotide is operatively linked with a regulatory sequence that controls expression of the polynucleotide in a host cell. In certain embodiments, the invention provides a host cell comprising a CLASP-2 polynucleotide, wherein the nucleotide sequence of the polynucleotide is operatively linked with a regulatory sequence that controls expression of the polynucleotide in a host cell, or progeny of the cell.

In another aspect, the invention further provides a CLASP-2 polynucleotide that is an antisense polynucleotide. In a preferred embodiment, the antisense polynucleotide is less than about 200 bases in length. In other embodiments, the invention provides an antisense oligonucleotide complementary to a messenger RNA comprising SEQ ID NO: 1, 3, 5 or 9 and encoding CLASP-2, wherein the oligonucleotide inhibits the expression of CLASP-2.

In another aspect, the invention provides an isolated DNA that encodes a CLASP-2 protein as shown in SEQ ID NO: 2, 4, 6 or 10. In certain embodiments, the CLASP-2 polynucleotide is RNA.

The invention provides a method for producing a polypeptide comprising: (a) culturing the host cell containing a CLASP-2 polynucleotide under conditions such that the polypeptide is expressed; and (b) recovering the polypeptide from the cultured host cell or its cultured medium.

The invention further provides an isolated CLASP-2 polypeptide encoded by a CLASP-2 polynucleotide. In some embodiments, the CLASP-2 polypeptide has the amino acid sequence of SEQ ID NO: 2, 4, 6 or 10, or a fragment thereof. In some embodiments, the isolated CLASP-2 polypeptide is cell-membrane associated. In other embodiments, the isolated CLASP-2 polypeptide is soluble. In other embodiments, the soluble CLASP-2 polypeptide is fused with a heterologous polypeptide.

The invention further provides an isolated CLASP-2 protein having the sequence as shown in SEQ ID NO: 2, 4, 6 or 10. In some embodiments, the invention provides a CLASP-2 protein comprising the sequence as shown in SEQ. ID. NO: 1 and variants thereof that are at least 95% identical to SEQ ID. NO: 2 and specifically binds a cytoskeletal protein. In certain embodiments the cytoskeletal protein is spectrin.

The invention further provides an isolated antibody that specifically binds to a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 10, or a

binding fragment thereof. In some embodiments the antibody is monoclonal. In other embodiments, the invention provides a hybridoma capable of secreting the antibody.

The invention further provides a method of identifying a compound or agent that binds a CLASP-2 polypeptide comprising: i) contacting a CLASP-2 polypeptide with the compound or agent under conditions which allow binding of the compound to the CLASP-2 polypeptide to form a complex and ii) detecting the presence of the complex.

The invention further provides a method of detecting a CLASP-2 polypeptide in a sample, comprising: (a) contacting the sample with a CLASP-2 antibody or binding fragment and (b) determining whether a complex has been formed between the antibody and with CLASP-2 polypeptide.

The invention further provides a method of detecting a CLASP-2 polypeptide in a sample, comprising: (a) contacting the sample with a CLASP-2 polynucleotide or a polynucleotide that comprises a sequence of at least 12 nucleotides and is complementary to a contiguous sequence of the CLASP-2 polynucleotide and (b) determining whether a hybridization complex has been formed.

The invention further provides a method of detecting a CLASP-2 nucleotide in a sample, comprising: (a) using a polynucleotide that comprises a sequence of at least 12 nucleotides and is complementary to a contiguous sequence of a CLASP-2 polynucleotide in an amplification process; and (b) determining whether a specific amplification product has been formed.

The invention further provides pharmaceutical compositions comprising a CLASP-2 polynucleotide, a CLASP-2 polypeptide, or a CLASP-2 antibody and a pharmaceutically acceptable carrier.

In one aspect, the invention provides a method of inhibiting an immune response in a cell comprising: (a) interfering with the expression of a CLASP-2 gene in the cell; (b) interfering with the ability of a CLASP-2 protein to mediate cell-cell interaction (*e.g.*, interfering with a heterotypic and/or homotypic interaction) between CLASP-2 and an extracellular protein; (c) interfering with the ability of a CLASP-2 protein to bind to another protein. In some such methods, the cell is a T cell or a B cell. Some such methods comprise contacting the cell with an effective amount of a polypeptide which comprises the amino acid sequence of SEQ ID NO: 2, 4, 6 or 10 or a fragment thereof.

In another aspect, the invention provides a method of inhibiting an immune response in a subject, comprising administering to the subject a therapeutically effective

amount of an antibody which specifically binds a polypeptide having the sequence of SEQ ID NO: 2, 4, 6 or 10.

In another aspect, the invention provides a method of preventing or treating a CLASP-2-mediated disease comprising administering to a subject in need thereof a therapeutically effective amount of a CLASP-2 pharmaceutical composition. In some such methods, the CLASP-2-mediated disease is an autoimmune disease.

The invention further provides a method of treating an autoimmune disease in a subject caused or exacerbated by increased activity of T_H1 cells consisting of administering a therapeutically effective amount of a CLASP-2 pharmaceutical composition to the subject.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Nucleotide and predicted amino acid sequence of CLASP-2A cDNA. Notable protein motifs are indicated above the nucleotide sequence in bold. Potential initiator methionines are underscored. The notable, predicted protein motifs are: a cadherin cleavage site encoded by nucleotides 854-868, a cadherin ectodomain (EC) encoded by nucleotides 1253-1264, a transmembrane domain encoded by nucleotides 2861-2917, a coiled coil domain encoded by nucleotides 3579-3682, a second coiled coil domain encoded by nucleotides 3827-3937, and a PDZ binding motif (PBM) encoded by nucleotides 4046-4057.

Figure 2. A. Schematic of CLASP-2 splice variants. Splice variants are compared to Human (h) CLASP-2A. Numbers above hCLASP-2A line diagram indicate where splice variations comprising deletions and insertions relative to hCLASP-2A are found. Abbreviations: "KIAA" KIAA1058 sequence (Genbank Accession No. AB028981). **B.** Nucleotide and predicted amino acid sequence of CLASP-2A cDNA. Notable protein motifs are indicated above the nucleotide sequence in bold. Exact position of insertions and deletions are indicated above the CLASP-2A sequence with arrows and "x", respectively.

The nucleotide sequence of insertions schematized in FIG. 2A are indicated above the arrow. The insertions and deletions are as follows (numeration refers to Human CLASP-2A nucleotide sequence): Nucleotides 1966-2034 are deleted in CLASP-2D. Nucleotides 2219-2224 are deleted in CLASP-2B. There is an insertion of 69 amino acids at nucleotide 2927 found in CLASP-2D. The nucleotide sequence for this insertion is:

AAGCAGTCCAGTGGGAGCCGCCCTTCTCCCCACAGCCATAGCGCCTGCCTGAG GAGGAGCCGGGGAG and encodes amino acids AVQWEPPLPHSHSACLRRSRG (one letter amino acid abbreviation). This amino acid sequence encodes a putative SH3 binding

domain. There is another deletion at between nucleotides 3011-3079 found in CLASP-2E. CLASPs 2B, 2C, 2D and 2E contain an insertion at nucleotide 3153 with the nucleotide sequence of:

TGAGAGGCTGGCCCATCTGTATGACACGCTGCACCGGGCCTACAGCAAAGTGAC
 5 CGAGGTCATGCACTCGGGCCGCAGGCTTCTGGGGACCTACTTCCGGGTAGCCTTC
 TTCGGGCAGGCAGCGCAATACCAGTTTACAGACAGTGAAACAGATGTGGAGGGA

TT. The entire sequence is found in CLASP-2D and encodes amino acids

ERLAHL^YDTL^IHRAYSKVTEVMHSGRRL^LLGTYFRVAFFGQAAQYQFTDSETDVEG
 while the underline sequence is found in CLASPs 2B, 2C, and 2E and encodes amino acids

10 ERLAHL^YDTL^IHRAYSKVTEVMHSGRRL^LLGTYFRVAFFGQG. This amino acid
 sequence encodes a putative immunoreceptor tyrosine-based activation motif (ITAM). There
 is a two nucleotide deletion in Human CLASP-2C found at nucleotides 3586 and 3587.
 There is an insertion of 8 nucleotides found only in Human CLASP-2D with sequence:
 CTGGGATG at nucleotide 3937. This insertion puts a stop codon into the CLASP-2D
 15 nucleotide sequence.

Figure 3. A. Alignment of nucleotide sequences of the CLASP-2 isoforms.

Sequences were aligned using ClustalW B. Alignment of amino acid sequence of the
 CLASP-2 isoforms. Sequences were aligned using ClustalW. One letter amino acid
 abbreviation is used.

20 **Figure 4. Expression of CLASP-2 in human cell lines and human tissues as**
 determined by Northern hybridization. A CLASP-2-specific DNA fragment was generated
 by PCR from a CLASP-2 cDNA clone (HC2-5'), using primers HC2AS2 and HC2S1. The
 fragment was labeled by incorporation of radioactive ³²P dCTP. A. Expression in human
 tissues. The labeled DNA fragment was used as a probe on a human Multiple Tissue
 25 Northern (Clontech MTN Blot, #7780-1). A single band is clearly detect migrating at
 approximately 7.5 kb in placenta, heart kidney and lung in the Multiple Tissue Northern.
 Slight expression is detected in liver, skeletal muscle and brain. B. Expression in
 hematopoietic cell lines. A Northern with RNA from multiple cells lines was hybridized with
 the same hCLASP-2 probe. A similarly migrating band is detected in Jurkat (T-cell derived),
 30 9D10 (B-cell derived) and 293 (human kidney derived) cell lines. There are multiple weaker
 bands in the 9D10 lane indicating possible splice variants of hCLASP-2. Weak expression is
 also detected in the mouse cell lines CH27 (B cell lymphoma) and 3A9 (T-cell hybridoma).

Since hybridization and washing were carried out at high stringency, this indicates that the human CLASP-2 probe may cross-react with mouse CLASP mRNA.

Figure 5. A. Amino acid sequence of human and rat CLASP proteins.

Sequences were aligned using ClustalW. One letter amino acid abbreviation used. Protein motifs are found within the labeled boxes. A "-" indicates gaps that are placed to acquire a best overall alignment. Other abbreviations: "HC2A" Human CLASP-2 sequence, "KIAA" KIAA1058 sequence (Genbank Accession No. AB028981), "rat" TRG gene (Genbank Accession No. X68101), "HC4" Human CLASP-4 sequence, "HC1" Human CLASP-1 sequence, "HC3" Human CLASP-3 sequence, "HC5" Human CLASP-5 sequence. **B.**

Alignment of DOCK motifs found within the human CLASPs and compared to canonical DOCK motifs. Consensus amino acids found within all DOCK motifs are also indicated.

Figure 6. A. Nucleotide and predicted amino acid sequence of CLASP-2A

cDNA. Notable protein motifs are indicated (see FIG. 1 legend for details). Additionally, boundaries between exons and introns are indicated by arrows. These boundaries were defined by sequencing Bacterial Artificial Chromosomes (BACs) containing genomic DNA corresponding to CLASP-2. BACs were sequenced using primers derived from exon sequences corresponding to the CLASP-2 cDNA. Each exon/intron boundary is noted (as "Ref" with an appropriate reference number) above the cDNA sequence. The references contain exact nucleotide location of introns. The names and nucleotide numbers of the primers that were used in sequence reactions are also indicated. All nucleotide numbers refer to CLASP-2A cDNA sequence. As shown in the reference, not all of the sequence from sequencing reactions produced sequence matching the cDNA. These nucleotide sequences that did not match the exon sequence for CLASP-2 were considered to be intron sequences.

B. Alignment of human and rat CLASP amino acid sequences by ClustalW. Notable protein motifs are indicated (see FIG. 1 Legend for additional details). Additionally, the exon/intron borders described in part A are indicated with vertical lines between appropriate amino acids. Reference numbers are indicated in the right margin and correspond to references in Fig 6A and B.

Figure 7. Southern hybridization analysis of CLASP-2. Genomic DNA from

HeLa cells or a BAC DNA clone was digested with EcoRI or HindIII (genomic DNA) or Pst I (BAC DNA) and electrophoresed and transferred to nylon membrane by standard methods. For a probe, a CLASP-2-specific DNA fragment was generated by PCR from a CLASP-2

cDNA clone (HC2-5'), using primers HC2AS2 and HC2S1. The fragment was labeled by incorporation of radioactive ^{32}P dCTP. Probe HC2.1 is 800 bp long and it recognizes two fragments (~4.5 kb and 1.85 kb) on Eco RI digested genomic DNA. Three fragments are revealed by this probe when hybridized to digested DNA of BACs 4 and 6, with the two major ones identical in size to those detected on genomic DNA.

Figure 8. Expression of human CLASP-1 (hCLASP-1) CLASP-1 and CLASP-2 Glutathion-S-Transferase (GST) fusion proteins. Nucleotides encoding a portion of the hCLASP-2A intracellular domain (nucleotides 3230-4065) were subcloned into pGEX vectors (Pharmacia). Recombinant plasmids were transformed into *E. coli* (strain DH5 α), and transformed strains were grown by standard conditions. While in log phase cells were either induced (I) with IPTG (0.1 mM final concentration) or left uninduced (U). After several additional hours of growth cells were harvested and soluble protein lysates generated by standard methods. Aliquots of the protein lysates were electrophoresed on SDS-PAGE along with molecular mass standards. The gel was stained with Coomassie Blue and shows that fusion proteins migrated with their predicted molecular masses of 59 and 57 kD for hCLASP-1 and hCLASP-2, respectively.

Figure 9. A. Binding of CLASP-2 C-terminal 20 amino acids to PDZ domains. 20 μM biotinylated synthetic peptide corresponding to the C-terminal 20 amino acids of CLASP-2 was reacted with the indicated plate bound GST fusion proteins (none = no GST fusion protein coated onto plate). Error bars indicate standard deviation of duplicate measurements. **B.** Affinity of CLASP 2 – PDZ interactions. Varying concentrations of biotinylated CLASP-2 peptide were reacted with plate bound GST alone, GST-DLG1, GST-NeDLG, and GST-PSD95 fusion proteins. The binding to GST alone (< 0.1 OD units) was subtracted from the binding to the fusion proteins and the remaining signal was divided by the signal observed upon addition of 30 μM CLASP-2 peptide to each PDZ domain-containing protein (0.4 – 1.0 OD units) and plotted. The plotted data was fit to a saturation binding curve, yielding an apparent affinity of 7.5 μM for NeDLG- CLASP-2 interaction, 21 μM for DLG1- CLASP-2 interaction, and 45 μM for PSD95-CLASP-2 interaction. Data are means of duplicate data points, with standard errors between duplicate data points < 10%. **C.** Inhibition of CLASP-2 – PDZ binding. 5 μM biotinylated synthetic peptide corresponding to the C-terminal 20 amino acids of CLASP-2 was reacted with the indicated, plate-bound PDZ domain-containing GST fusion proteins in the presence or absence of 100 μM competitor

peptide. CLASP-2 Inhibitor refers to a synthetic peptide composed of the eight C-terminal amino acids of CLASP-2. KV1.3 Inhibitor refers to a synthetic peptide composed of the 19 C-terminal amino acids of KV1.3, a lymphocyte potassium channel. The amino acid sequence of the KV1.3 inhibitor is TTNNNPNSAVNIKKIFTDV. **D. Inhibition of KV1.3 – PDZ binding.** 5 μ M biotinylated synthetic peptide corresponding to the C-terminal 19 amino acids of KV1.3 was reacted with the indicated, plate-bound PDZ-domain containing GST fusion proteins in the presence or absence of 100 μ M CLASP-2 Inhibitor (see FIG. 9C legend).

Figure 10. Preliminary nucleotide sequences of CLASP-2 cDNAs.

Figure 11. A) Full length cDNA sequence and predicted amino acid translation of the human CLASP-2 gene. Predicted initiator methionine starts at nucleotide +1. Three independent 1st exons (indicated as 11A, 11B and 11C) splice into the second exon starting at nucleotide -101. The sequence appearing in FIG. 1 corresponds to nucleotides 1884 through 6690 of FIG 11A. B) Differences between the human CLASP-2 cDNA isoforms. In addition to the differential first exon usage indicated in A, sequencing multiple, independent cDNA products revealed nucleotide polymorphisms (allelic variations) between CLASP-2 cDNA isoforms. Additionally, differential exon usage through alternative splicing events was discovered. The use of the exon in B leads to a premature stop codon that can generate a soluble form of CLASP-2. C. Schematic of human CLASP-2 cDNAs. The top line represents nucleotide numbering found in FIG. 11A. Line (i) represents CLASP-2 cDNA shown in FIG. 1 above; line (ii) represents the full length CLASP-2 isoforms, where there are three CLASP-2 full length cDNA isoforms (A + Z, B + Z, and C + Z). Each of the isoforms uses a unique first exon (A, B, and C) (see FIG. 11A) that splices into the rest of the cDNA from exon 2 onwards represented by Z. The portion of the cDNA represented by Z itself has alternative splice and nucleotide polymorphisms that are shown in FIG. 2 above. Line (iii) represents the additional 5' sequence with a small region of overlap between nucleotides 1884 to 2109 in FIG. 11A and nucleotides 1-225 of FIG. 1.

Figure 12. Sequence of human CLASP-2 exons and intron boundaries. A Sequence of human CLASP-2 exons and intron borders. Stretches of noncontiguous genomic sequence from the Human Genome Project (GENBANK entry gi9988160) were aligned using the human CLASP-2 cDNA as a template and Sequencher sequence analysis software (Gene Codes Corp). 22 exons representing approximately the 5' 20% of the human CLASP-

2 cDNA sequence are presented in predicted 5' to 3' order. Exon sequences are underlined and are flanked by intron sequence. Nucleotide numbers in parentheses refer to the exon sequence within the uniquely-generated, contiguous gi9988160 sequence, which is located in B. B. Ordered stretch of human genomic DNA at the CLASP-2 locus aligned from noncontiguous, shotgun sequencing from the Human Genome Project using the human CLASP-2 sequence from FIG. 5A to determine genomic DNA fragment order and orientation.

Figure 13. Amino acid alignment and comparison between the human (h) CLASP family members. Amino acid sequences were aligned using ClustalW. The alignment is presented in order of their greatest pairwise similarity scores. Single letter amino acid abbreviations are used. Astericks indicate complete identity, while colons and periods indicate sequence similarity among CLASP family members. Dashes indicate gaps inserted in the amino acid sequence to facilitate alignment. Labelled boxes are domains with similarity to known protein motifs; unlabelled boxes represent regions of similarity between all CLASPs and may represent CLASP-specific domains.

Figure 14. Expression of CLASP-2 upon T-cell activation as assayed by Northern analysis. Jurkat cells were activated using PMA, Ionomycin, and α CD28. RNA was prepared from cell culture aliquots at 0, 1, 2, 4, 8, 14 hours post activation and Northern analysis was performed (A). Hybridization signals obtained with a CLASP-2-specific probe were quantified using a phosphor imager system. Relative signal intensities (refers to total signal of the specific probe used) are shown in the bar diagram (B). The ethidium staining of the Northern gel (A) demonstrates even RNA loading.

DETAILED DESCRIPTION

5.0 Definitions

Except when noted, the terms "patient" or "subject" are used interchangeably and refer to mammals such as human patients and non-human primates, as well as experimental animals such as rabbits, rats, and mice, and other animals.

The term "biological sample" as used herein is a sample of biological tissue, fluid, or cells that contains hCLASP-2 or nucleic acid encoding hCLASP-2 protein. Such samples include, but are not limited to, tissue isolated from humans. Biological samples may also include sections of tissues such as frozen sections taken for histologic purposes. A biological sample is typically obtained from a eukaryotic organism, preferably eukaryotes

such as fungi, plants, insects, protozoa, birds, fish, reptiles, and preferably a mammal such as rat, mice, cow, dog, guinea pig, or rabbit, and most preferably a primate such as chimpanzees or humans.

5 The term "treating" includes the administration of the compounds or agents of the present invention to prevent or delay the onset of the symptoms, complications, or biochemical indicia of a disease, alleviating the symptoms or arresting or inhibiting further development of the disease, condition, or disorder (*e.g.*, autoimmune disease). Treatment may be prophylactic (to prevent or delay the onset of the disease, or to prevent the manifestation of clinical or subclinical symptoms thereof) or therapeutic suppression or
10 alleviation of symptoms after the manifestation of the disease.

The term "lymphocyte" as used herein has the normal meaning in the art, and refers to any of the mononuclear, nonphagocytic leukocytes, found in the blood, lymph, and lymphoid tissues, *i.e.*, B and T lymphocytes.

The terms "isolated," or "purified," refer to material that is substantially free
15 from components that normally accompany it as found in its native state (*e.g.*, recombinantly produced or purified away from other cell components with which it is naturally associated). Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. The term "purified" denotes that a nucleic acid or protein gives rise to essentially one band in an
20 electrophoretic gel. Particularly, it means that the nucleic acid or protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure.

The terms "nucleic acid" and "polynucleotide" are used interchangeably" and refer to refers to DNA, RNA and nucleic acid polymers containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-
25 naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

30 The terms "polypeptide," "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The amino acids may be natural amino acids, or include an artificial chemical mimetic of a corresponding naturally occurring amino acid.

As used herein a “nucleic acid probe” is defined as a nucleic acid capable of specifically binding to a target nucleic acid of complementary sequence (*e.g.*, through complementary base pairing). As used herein, a probe may include natural (*i.e.*, A, G, C, or T) or modified bases (7-deazaguanosine, inosine, and the like). In addition, the bases in a probe may be joined by a linkage other than a phosphodiester bond, so long as it does not interfere with hybridization (*e.g.*, probes may be peptide nucleic acids). The probes can be directly labeled as with isotopes, chromophores, lumiphores, chromogens, or indirectly labeled such as with biotin to which a streptavidin complex may later bind.

The term “recombinant” when used with reference, *e.g.*, to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or, in the case of cells, to progeny of a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

The term “heterologous” when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, *e.g.*, a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (*e.g.*, a fusion protein).

The term “sequence identity” refers to a measure of similarity between amino acid or nucleotide sequences, and can be measured using methods known in the art, such as those described below:

The terms “identical” or percent “identity,” in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (*i.e.*, 60% identity, preferably 65%, 70%, 75%, 80%, 85%, 90%, or 95% identity over a specified region (see, *e.g.*, SEQ ID NO: 1), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection.

The phrase "substantially identical," in the context of two nucleic acids or polypeptides, refers to two or more sequences or subsequences that have at least of at least 60%, often at least 70%, preferably at least 80%, most preferably at least 90% or at least 95% nucleotide or amino acid residue identity, when compared and aligned for maximum
5 correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. Preferably, the substantial identity exists over a region of the sequences that is at least about 50 bases or residues in length, more preferably over a region of at least about 100 bases or residues, and most preferably the sequences are substantially identical over at least about 150 bases or residues. In a most preferred embodiment, the sequences are
10 substantially identical over the entire length of the coding regions.

The phrase "sequence similarity" in the context of two nucleic acids or polypeptides, refers to two or more sequences that are identical or in the case of amino acids, have homologous amino acid substitutions at either 50%, often at least 60%, often at least 70%, preferably at least 80%, most preferably at least 90% or at least 95% of the
15 indicated positions.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program
20 parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters. For sequence comparison of nucleic acids and proteins to CLASP-2 nucleic acids and proteins, the BLAST and BLAST 2.0 algorithms and the default parameters discussed below are used.

25 A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences
30 for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, 1981, Adv. Appl. Math. 2: 482), by the homology alignment algorithm of Needleman & Wunsch, 1970, J. Mol. Biol. 48: 443, by the search for similarity method of Pearson & Lipman, 1988, Proc. Natl. Acad. Sci. U.S.A. 85: 2444, by computerized implementations of these algorithms

(FASTDB (Intelligenetics), BLAST (National Center for Biomedical Information), GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, *e.g.*, Ausubel *et al.*, 1987 (1999 Suppl.), Current Protocols in Molecular Biology, Greene Publishing Associates and Wiley Interscience, N.Y.)

A preferred example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the FASTA algorithm, which is described in Pearson, W.R. & Lipman, D.J., 1988, Proc. Natl. Acad. Sci. U.S.A. 85: 2444. See also W. R. Pearson, 1996, Methods Enzymol. 266: 227-258. Preferred parameters used in a FASTA alignment of DNA sequences to calculate percent identity are optimized, BL50 Matrix 15: -5, k-tuple= 2; joining penalty= 40, optimization= 28; gap penalty -12, gap length penalty =2; and width= 16.

Another preferred example of algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, 1977, Nuc. Acids Res. 25: 3389-3402 and Altschul *et al.*, 1990, J. Mol. Biol. 215: 403-410, respectively. BLAST and BLAST 2.0 are used, with the parameters described herein, to determine percent sequence identity for the nucleic acids and proteins of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences)

uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, 1989, Proc. Natl. Acad. Sci. U.S.A. 89: 10915) alignments (B) of
5 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, 1993, Proc. Natl. Acad. Sci. U.S.A. 90: 5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between
10 two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

Another example of a useful algorithm is PILEUP. PILEUP creates a multiple
15 sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, 1987, J. Mol. Evol. 35: 351-360. The method used is similar to the method described by Higgins & Sharp, 1989, CABIOS 5: 151-
20 153. The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two
25 individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default
30 gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, *e.g.*, version 7.0 (Devereaux *et al.*, 1984, Nuc. Acids Res. 12: 387-395).

Another preferred example of an algorithm that is suitable for multiple DNA and amino acid sequence alignments is the CLUSTALW program (Thompson, J. D. *et al.*,

1994, Nucl. Acids. Res. 22: 4673-4680). ClustalW performs multiple pairwise comparisons between groups of sequences and assembles them into a multiple alignment based on homology. Gap open and Gap extension penalties were 10 and 0.05 respectively. For amino acid alignments, the BLOSUM algorithm can be used as a protein weight matrix (Henikoff and Henikoff, 1992, Proc. Natl. Acad. Sci. U.S.A. 89: 10915-10919).

A "label" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (*e.g.*, as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins for which antisera or monoclonal antibodies are available (*e.g.*, the polypeptide of SEQ ID NO: 1 can be made detectable, *e.g.*, by incorporating a radiolabel into the peptide, and used to detect antibodies specifically reactive with the peptide).

The term "sorting" in the context of cells as used herein to refers to both physical sorting of the cells, as can be accomplished using, *e.g.*, a fluorescence activated cell sorter, as well as to analysis of cells based on expression of cell surface markers, *e.g.*, FACS analysis in the absence of sorting.

The phrase "selectively (or specifically) hybridizes to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (*e.g.*, total cellular or library DNA or RNA).

The phrase "specifically (or selectively) binds" to an antibody refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample.

The phrase "specifically bind(s)" or "bind(s) specifically" when referring to a peptide refers to a peptide molecule which has intermediate or high binding affinity, exclusively or predominately, to a target molecule. The phrases "specifically binds to" refers to a binding reaction which is determinative of the presence of a target protein in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated assay conditions, the specified binding moieties bind preferentially to a particular target protein and do not bind in a significant amount to other components present in a test sample. Specific binding to a target protein under such conditions may require a binding moiety that is selected for its specificity for a particular target antigen. A variety of assay formats may be

used to select ligands that are specifically reactive with a particular protein. For example, solid-phase ELISA immunoassays, immunoprecipitation, Biacore and Western blot are used to identify peptides that specifically react with PDZ domain-containing proteins. Typically a specific or selective reaction will be at least twice background signal or noise and more
5 typically more than 10 times background. Specific binding between a monovalent peptide and a PDZ-containing protein means a binding affinity of at least 10^4 M^{-1} , and preferably 10^5 or 10^6 M^{-1} .

The phrase "homotypic interaction" refers to the binding of a given protein to another molecule of the same protein (*e.g.*, the binding of hCLASP-2 to hCLASP-2). The
10 phrase "heterotypic interaction" refers to the binding of a given protein to a different protein or other molecule (*e.g.*, the binding of hCLASP-2 to a PDZ domain-containing protein or the binding of a transcription factor to DNA).

The phrase "immune cell response" refers to the response of immune system cells to external or internal stimuli (*e.g.*, antigen, cytokines, chemokines, and other cells)
15 producing biochemical changes in the immune cells that result in immune cell migration, killing of target cells, phagocytosis, production of antibodies, other soluble effectors of the immune response, and the like.

The terms "B lymphocyte response" and "B lymphocyte activity" are used interchangeably to refer to the component of immune response carried out by B lymphocytes
20 (*i.e.* the proliferation and maturation of B lymphocytes, the binding of antigen to cell surface immunoglobulin, the internalization of antigen and presentation of that antigen via MHC molecules to T lymphocytes, and the synthesis and secretion of antibodies).

The terms "T lymphocyte response" and "T lymphocyte activity" are used here interchangeably to refer to the component of immune response dependent on T
25 lymphocytes (*i.e.*, the proliferation and/or differentiation of T lymphocytes into helper, cytotoxic killer, or suppressor T lymphocytes, the provision of signals by helper T lymphocytes to B lymphocytes that cause or prevent antibody production, the killing of specific target cells by cytotoxic T lymphocytes, and the release of soluble factors such as cytokines that modulate the function of other immune cells).

The term "immune response" refers to the concerted action of lymphocytes,
30 antigen presenting cells, phagocytic cells, granulocytes, and soluble macromolecules produced by the above cells or the liver (including antibodies, cytokines, and complement) that results in selective damage to, destruction of, or elimination from the human body of

invading pathogens, cells or tissues infected with pathogens, cancerous cells, or, in cases of autoimmunity or pathological inflammation, normal human cells or tissues.

Components of an immune response may be detected in vitro by various methods that are well known to those of ordinary skill in the art. For example, (1) cytotoxic
5 T lymphocytes can be incubated with radioactively labeled target cells and the lysis of these target cells detected by the release of radioactivity, (2) helper T lymphocytes can be incubated with antigens and antigen presenting cells and the synthesis and secretion of cytokines measured by standard methods (Windhagen A; *et al.*, 1995, Immunity 2(4): 373-80), (3) antigen presenting cells can be incubated with whole protein antigen and the
10 presentation of that antigen on MHC detected by either T lymphocyte activation assays or biophysical methods (Harding *et al.*, 1989, Proc. Natl. Acad. Sci., 86: 4230-4), (4) mast cells can be incubated with reagents that cross-link their Fc-epsilon receptors and histamine release measured by enzyme immunoassay (Siraganian, *et al.*, 1983, TIPS 4: 432-437).

Similarly, products of an immune response in either a model organism (*e.g.*,
15 mouse) or a human patient can also be detected by various methods that are well known to those of ordinary skill in the art. For example, (1) the production of antibodies in response to vaccination can be readily detected by standard methods currently used in clinical laboratories, *e.g.*, an ELISA; (2) the migration of immune cells to sites of inflammation can be detected by scratching the surface of skin and placing a sterile container to capture the
20 migrating cells over scratch site (Peters *et al.*, 1988, Blood 72: 1310-5); (3) the proliferation of peripheral blood mononuclear cells in response to mitogens or mixed lymphocyte reaction can be measured using ³H-thymidine; (4) the phagocytic capacity of granulocytes, macrophages, and other phagocytes in PBMCs can be measured by placing PMBCs in wells together with labeled particles (Peters *et al.*, 1988); and (5) the differentiation of immune
25 system cells can be measured by labeling PBMCs with antibodies to CD molecules such as CD4 and CD8 and measuring the fraction of the PBMCs expressing these markers.

As used herein, the phrase "signal transduction pathway" or "signal transduction event" refers to at least one biochemical reaction, but more commonly a series of biochemical reactions, which result from interaction of a cell with a stimulatory compound
30 or agent. Thus, the interaction of a stimulatory compound with a cell generates a "signal" that is transmitted through the signal transduction pathway, ultimately resulting in a cellular response, *e.g.*, an immune response described above.

A signal transduction pathway refers to the biochemical relationship between a variety of signal transduction molecules that play a role in the transmission of a signal from

one portion of a cell to another portion of a cell. Signal transduction molecules of the present invention include, for example, extracellular and intracellular domains of CLASP-2. As used herein, the phrase "cell surface receptor" includes molecules and complexes of molecules capable of receiving a signal and the transmission of such a signal across the plasma membrane of a cell. An example of a "cell surface receptor" of the present invention is the T cell receptor (TCR). As used herein, the phrase "intracellular signal transduction molecule" includes those molecules or complexes of molecules involved in transmitting a signal from the plasma membrane of a cell through the cytoplasm of the cell, and in some instances, into the cell's nucleus. In the present invention, CLASP-2 can be referred to as an "intracellular signal transduction molecule", but can also be referred to as a "signal transduction molecule".

A signal transduction pathway in a cell can be initiated by interaction of a cell with a stimulator that is inside or outside of the cell. If an exterior (*i.e.*, outside of the cell) stimulator (*e.g.*, an MHC-antigen complex on an antigen presenting cell) interacts with a cell surface receptor (*e.g.*, a T cell receptor), a signal transduction pathway can transmit a signal across the cell's membrane, through the cytoplasm of the cell, and in some instances into the nucleus. If an interior (*e.g.*, inside the cell) stimulator interacts with an intracellular signal transduction molecule, a signal transduction pathway can result in transmission of a signal through the cell's cytoplasm, and in some instances into the cell's nucleus.

Signal transduction can occur through, *e.g.*, the phosphorylation of a molecule; non-covalent allosteric interactions; complexing of molecules; the conformational change of a molecule; calcium release; inositol phosphate production; proteolytic cleavage; cyclic nucleotide production and diacylglyceride production. Typically, signal transduction occurs through phosphorylating a signal transduction molecule. According to the present invention, a CLASP-2 signal transduction pathway refers generally to a pathway in which CLASP-2 protein regulates a pathway that includes engaged-receptors, PKC-substrates, G proteins, and other molecules.

5.1. Introduction

The present invention relates to a novel transmembrane protein, CLASP-2, a new member of the CLASP family that contains an endodomain that displays the appropriate properties to organize the cytoskeleton and signal transduction apparatus of the immune gateway.

CLASP-2 functions in cells of the immune system, *e.g.*, T cells and B cells, as well as non-immune cells. The CLASP-2 protein functions in a variety of cellular processes,

particularly related to immune function, regulation of T cell and B cell interactions, T cell activation, and in the organization, establishment and maintenance of the “immunological synapse” (see Dustin *et al.*, 1999, Science 283: 680-682; Paul *et al.*, 1994, Cell 76: 241-251; Dustin *et al.*, 1996, J. Immunol. 157: 2014; Dustin *et al.*, 1998, Cell 94: 667), including
5 signal transduction, cytoskeletal interactions, and membrane organization.

Without intending to be bound by a particular mechanism or limited in any way, the CLASP-2 protein is believed to be a component of the lymphocyte organelle called the “immune gateway” that creates a docking site or portal for cell-cell contact during antigen-presentation. It is believed the cytoplasmic domains of CLASP-2 proteins organize it
10 into a patch at the leading edge of T cells. The carboxy-terminus encoded sequences mediate interaction with PDZ domain proteins and with cytoskeletal proteins (*e.g.*, spectrin or ankyrin) to connect CLASP-2 to the microtubule network and hold the receptors at a polarized configuration just above the microtubule-organizing center (“MTOC”). Thus, when T cells engages a B cell acting as an APC, the CLASP-2 molecules engage one another
15 to dock the two cells and organize the immune synapse.

Modulating the expression of the CLASP-2 protein, and interference with, or enhancement of, CLASP-2 protein interactions with other proteins has a number of beneficial physiological effects, *e.g.*, altered signaling in response to antigen, altered T and B cell response to antigen, and modulation of T cell activation. In one aspect, the CLASP-2
20 extracellular domain is targeted (*e.g.*, using anti-CLASP-2 antibody, soluble CLASP-2 fragments, and the like) to regulate T cell activation (and thus regulate immune responses). Disorders that can be treated by disrupting CLASP-2 function, include without limitation, multiple sclerosis, juvenile diabetes, rheumatoid arthritis, pemphigus, pemphigoid, epidermolysis bullosa acquista, lupus, endometriosis, toxemia or pregnancy induced
25 hypertension, pruritic urticarial papules and plaques of pregnancy (PUPPP), herpes gestationis, impetigo herpetiformis, pruritus gravidarum, placenta-related disorders, and Rh incompatibility.

In another aspect, the present invention provides methods and reagents for detection of CLASP-2 expression and CLASP-2-expressing cells. Abnormal expression
30 patterns or expression levels are diagnostic for immune and other disorders. For example, diseases characterized by overproduction or depletion of lymphocytes in blood or other organs may be detected or monitored by monitoring the level of CLASP-2 polypeptide or mRNA in a biological sample (*e.g.*, peripheral blood), *e.g.*, the number or percentage of CLASP-2 expressing cells. Diseases characterized by overproduction of T cells include, *e.g.*,

leukemia (both ALL and CLL), lymphoma (including non-Hodgkins lymphoma, Burkitt's lymphoma, mycosis fungoides, and sezary syndrome), EBV, CMV, toxoplasmosis, syphilis, typhoid, brucellosis, tuberculosis, influenza, hepatitis, serum sickness, and thyrotoxicosis. Diseases associated with the depletion of T cells include, *e.g.*, HIV and myelodysplasia.

- 5 Diseases associated with the overproduction of B cells include, *e.g.*, leukemia (both ALL and CLL), non-Hodgkins lymphoma, Burkitt's lymphoma, myeloma, EBV, CMV, toxoplasmosis, syphilis, typhoid, brucellosis, tuberculosis, influenza, hepatitis, serum sickness, and thyrotoxicosis. Diseases associated with the depletion of B cells include, *e.g.*, myelodysplasia.

10 5.2. CLASP-2 cDNA and Polypeptide Structure

The CLASP-2 protein is type I transmembrane glycoprotein, characterized by multiple forms produced by alternative exon usage (*i.e.*, production of splice variants). In one naturally occurring form, CLASP-2 has the structure shown in FIG. 1. However, as discussed in detail *infra*, the CLASP-2 gene encodes a variety of gene product due to
 15 alternative splicing of mRNA. FIG. 2 shows the nucleotide sequence and conceptual translation of human CLASP-2 polypeptides:

- hCLASP-2A cDNA (SEQ ID NO: 1) and hCLASP-2A polypeptide (SEQ ID NO: 2).
 hCLASP-2B cDNA (SEQ ID NO: 3) and hCLASP-2B polypeptide (SEQ ID
 20 NO: 4).
 hCLASP-2C cDNA (SEQ ID NO: 5) and hCLASP-2C polypeptide (SEQ ID NO: 6).
 hCLASP-2D cDNA (SEQ ID NO: 7) and hCLASP-2D polypeptide (SEQ ID NO: 8).
 25 hCLASP-2E cDNA (SEQ ID NO: 9) and hCLASP-2E polypeptide (SEQ ID NO: 10).

Unless specifically referred to, the phrase "human CLASP-2 (hCLASP-2)" is used herein refers to hCLASP-2A, hCLASP-2B, hCLASP-2C and hCLASP-2E. "hCLASP-2D" cDNA is also known as KIAA1058, which was described by Kikuno *et al.*, 1999, *DNA*
 30 *Res.* 6, 197-205 as a cDNA from brain encoding a protein of unknown function.

CLASP-2 polypeptides typically include an approximately 120 residue leader sequence, followed by a cadherin proteolytic cleavage signal RXXR, an extracellular domain, a transmembrane domain, and an intracellular domain. The present invention provides a polynucleotide having the sequence of SEQ. ID. NO: 1, or a fragment thereof, and a
 35 polypeptide having the sequence of SEQ. ID NO: 2, or a fragment thereof. In addition, the

invention provides polynucleotides comprising hCLASP-2 genomic sequences, CLASP-2 homologs from other species, naturally occurring alleles of hCLASP-2, and hCLASP-2 variants as described herein, and methods for using CLASP-2 polynucleotide, polypeptides, antibodies and other reagents.

5 5.2.1. CLASP-2 Polypeptide Domains

As is shown in FIG. 1, one naturally occurring CLASP-2 cDNA encodes a polypeptide characterized by several structural and functional domains and defined sequence motifs. To provide guidance to the practitioner, the structural features are described *infra*. However, it will be understood that the present invention is not limited to polypeptides that include all, or any particular one of these domains or motifs. For example, a CLASP-2 fusion protein of the invention contains only the extracellular domain of CLASP-2. Similarly, the CLASP-2A polypeptide of SEQ ID NO: 2 does not have the ITAM motifs (discussed *infra*) found in the CLASP-2B and 2C polypeptides.

It will be appreciated that the structurally (and functionally) different domains of CLASP-2 polypeptides (and the corresponding region of the mRNA) are of interest, in part, because they may be separately targeted or modified (*e.g.*, deleted or mutated) to affect the activity or expression of a CLASP-2 gene product (in order to, for example, modulate an immune response). For example, the extracellular domain of a CLASP-2 protein can be targeted (*e.g.*, using an anti-CLASP monoclonal antibody to (a) block the interaction of a CLASP-2-expressing cell (*e.g.*, a T cell) and a second cell (*e.g.*, a B cell) displaying a protein that is bound by CLASP-2 (*i.e.*, a CLASP-2 ligand). Similarly, an intracellular domain (*e.g.*, ITAM or DOCK, see *infra*) can be targeted to interfere with signal transduction without interfering with extracellular ligand binding.

Generally, inhibiting CLASP-2 expression or CLASP-2 polypeptide function will result in modulation of immune function including, for example, changing the threshold for T cell activation by affecting formation of the immune synapse. Modulation of immune function can be screened and quantitated by a number of assays known in the art and described herein (see also §5.14).

5.2.1.1. Signal Peptide

The human CLASP-2 sequence presented in FIG. 1 encodes two potential start sites for translation. The first predicted methionine appears at nucleotide 278 (ATG). The second methionine appears at nucleotide 476. Both have an acceptable consensus sequence

for a translational start (A/GxxATGG; Kozak, M., 1996, Mamm. Genome 7(8): 563-74). A polypeptide beginning at the second methionine is also predicted to encode a signal peptide capable of localizing the protein to the secretory pathway by SignalP, a signal sequence prediction program (Nielsen, H. *et al.*, 1997, Protein Eng. 10(1): 1-6). Polypeptides beginning at the first methionine are not predicted to contain a signal sequence; however, the consensus for signal prediction is only 80-90% accurate for known signal sequences. A third possibility for a translational start is that the cDNA listed in FIG. 1 is incomplete and another methionine is encoded in frame and upstream of the sequence shown in FIG.1 .

5.2.1.2. Extracellular Domain

The CLASP-2 extracellular domain is characterized by one cadherin EC-like motif (Pigott, R. and Power, C., 1993, The Adhesion Molecule Factbook. Academic Press, pg. 6; Jackson, R. M. and Russell, R. B., 2000, J. Mol. Biol. 296: 325-34). Several highly conserved cysteines are found in the extracellular domain, as well as various glycosylation signals. Through its extracellular domains, CLASP-2 may interact with ligands in a homotypic and/or heterotypic manner to establish the immunological synapse in conjunction with molecules such as TCR, MHC class I, MHC class II, CD3 complex and accessory molecules such as CD4, CD3, ICAM-1, LFA-1, and others. Many cadherins contain a pro-domain of approximately 50 to 150 amino acids that is removed before localization to the plasma membrane. This cleavage is presumed to be carried out by Furin (Posthaus, H. *et al.*, 1998, FEBS Let 438: 306-10) at a consensus sequence of RKQR. Furin is a protease that is at least partially responsible for the maturation of certain cadherins. CLASP-2 has the sequence RNQR at nucleotides 945 through 957. By homology, this region is around 120 amino acids into the predicted protein start site for hCLASP-2A. This region may be a pro-domain and cleavage may be required for CLASP-2 function, or aspects of CLASP-2 function.

Antibodies raised against the extracellular domain can be added to cells expressing CLASP-2. These antibodies can either block the interaction of CLASP-2 with potential ligands or stabilize these interactions. Any immunoassay known in the art, *e.g.*, listed and described herein, may be used to assess the modulation of immune function brought about by this approach.

Similarly, portions of the extracellular domain of CLASP-2 can be expressed as soluble protein. This soluble protein can then be added to cells expressing CLASP-2. These proteins may interact with potential ligands to competitively inhibit their binding to endogenous CLASP-2. This could modulate CLASP-2 function via the immunoassays

described herein. Recombinant proteins could interfere in a positive or negative fashion with CLASP-2 interactions.

5.2.1.3. Transmembrane Domain

CLASP-2 predicted amino acid sequence was analyzed using the PHDhtm analysis software for prediction of transmembrane helices (Rost, B., *et al.*, 1996, *Prot. Science* 7: 1704-1718). Using the PPHDhtm analysis software, it was determined that the transmembrane domain is located from nucleotides 2861-2917 (see FIG. 1), as well as three other potential transmembrane domains located near the amino terminal end.

5.2.1.4. Intracellular Domains

The CLASP-2 intracellular domains contain motifs corresponding to several types of protein domains. Depending on the specific CLASP-2 (*i.e.*, specific family member or splice variant) all or only some of the domains can be present. Listed from amino terminus to carboxy terminus, the domains include: (1) ITAM (Chan *et al.* 1994, *Annual Review of Immunology* 12: 555-592), (2) a newly discovered DOCK/CLASP-2 motif, (3) a coiled-coil motif, and (4) a C-terminal PDZ binding motif (PBM) (also referred to as PDZ ligand or "PL").

5.2.1.5. ITAM

Immunoreceptor Tyrosine-based Activation Motifs (ITAM motifs; also known as ARAM, or antigen recognition activation motifs) are motifs contained within antigen receptors for T and B cells, and Fc receptors on other leukocytes, and are necessary for proper activation and signal transduction in these cells. They are characterized by the consensus sequence YXXL/I - X7/8- YXXL/I (Grucza *et al.*, 1999, *Biochemistry* 38: 5024-5033), usually separated by 6-8 amino acids (Watson *et al.*, 1998, *Immunol. Today* 19: 260-264; Isakov, J. *Leukoc. Biol.* 61: 6-16). ITAM is used as an intracellular regulatory motif through its ability to be tyrosine phosphorylated by src-family tyrosine kinases such as Lyn that are involved in leukocyte signal transduction. Once phosphorylated, the ITAM acts as a high affinity binding site for SH2 containing proteins. Signal transduction components including ZAP-70, Syk, Lyn, Shc, PI3 kinase, and Grb2 contain SH2 domains and have been shown to bind ITAMs (Clements *et al.*, 1999, *Annu. Rev. Immunol.* 17: 89-108). This places ITAM-containing molecules in a central role of intracellular signal regulation in leukocytes. ITAM motifs in leukocyte signaling can facilitate signal transduction (*e.g.*,

tyrosine kinase signaling) by acting as temporal scaffolds where other transduction components could bind and be properly positioned to mediate transduction. ITAM motifs often appear in multiples in a protein, however, it is known that one set of YXXL/I alone can transduce signals of the PTK pathway, though weakly.

5 CLASP-2 proteins typically have ITAM YXXL/I motifs (where X is any amino acid) separated by 3 or 13 amino acids. In various embodiments the CLASP-2 polypeptide of the invention is characterized by one or more of the motifs shown in Table 1.

Table 1

CLASP-2 ITAM Motifs

Motif No.	Sequence Motif
1	YXX(I/L)-X ₃ -YXX(I/L)
2	YXX(I/L)-X ₁₃ -YXX(I/L)
3	YXX(I/L)-X ₃ -YXX(I/L)-X ₁₃ -YXX(I/L)

The presence of multiple ITAM motifs in CLASPs proteins indicates that they may be engaged by multiple signal transduction components (*e.g.*, ZAP-70/Syk, Shc, PI3 kinase, and Grb2). In general, the ITAM motif in CLASP proteins match identically to the canonical ITAM motif with some motifs containing a conservative amino acid change (*i.e.* valine instead of isoleucine or leucine). As previously described for other ITAMs, the ITAMs within CLASPs can bind SH2-containing proteins including ZAP-70, Syk, Shc, PI3 kinase, and Grb2. Since CLASPs have an extracellular domain, CLASPs protein can independently initiate a signal transduction cascade through engagement of its extracellular domain. Otherwise CLASPs may cooperate with an antigen receptor signaling complex (*e.g.*, with CD3/TCR, BCR, FcR), to facilitate tyrosine kinase signal transduction

20 The ITAMs have demonstrated different binding specificity and affinities for SH2 domains (Clements, *et al.*, 1999, Ann. Rev. Immunol. 17: 89-108). For example, Shc, PI3 kinase, and Grb2 bind to dual and mono phosphorylated ITAMs with different affinities. Thus the ITAMs in CLASPs are believed to provide quantitative as well as qualitative differences in signal transduction depending up their phosphorylation state, as well as to inhibit or augment specific protein interactions and hence specific tyrosine kinase-mediated signaling pathways in leukocytes.

Antagonizing the PTK-CLASP-2 interaction (*e.g.*, phosphorylation of CLASP-2) will thus inhibit immune function. In one embodiment, interactions between ITAM-bearing human CLASPs and their binding partners are believed to be antagonized by the alpha subtype (SIRPalpha) of signal regulatory proteins that has been shown to negatively

regulate ITAM-dependent lymphocyte activation (Lienard H; 1999, J Biol Chem-274: 32493-9). Also, a recently recognized family of immunoreceptor tyrosine-based inhibition motif (ITIM) receptors are thought to inhibit the ITAM-induced activation of immune competent cells (Gergely, *et al.*, 1999, J. Immunol Lett 68: 3-15) and therefore may block CLASP-partner interaction.

5.2.1.6. DOCK

CLASP-2 polypeptides contain a new "DOCK" motif, not previously described in the scientific literature. The CLASP DOCK motif includes a series of five tyrosines surrounded by conserved sequences in regions A, B, C, D, and G (see FIG. 5B).

10 There are also two highly conserved non-tyrosine containing regions (E and G) separated by nine amino acids (P+EXAI+XM) and (LXMXL+GXVXXXVNXG) (where X is any amino acid).

The cytoplasmic region of CLASP-2 immediately following the ITAM domains exhibits sequence similarity to the C-terminal third of the so-called "DOCK" proteins. The DOCK gene family includes three molecules that are the human homologues of the *C. elegans* CED proteins known to be involved in apoptosis. CED-5 (DOCK180), a major CRK-binding protein, alters cell morphology upon translocation to the membrane (mediates the membrane motion that scavenger cells exhibit as they surround and engulf dying cells; its function can be partially rescued by the human DOCK180 (Wu *et al.*, 1998, *Nature* 20 392: 501-504). Myoblast City in *Drosophila* (MBC) is another member of the DOCK protein family and has been found to be involved in myoblast fusion (Erickson, *et al.*, 1997, *J. Cell Biol.* 138: 589). Since CLASP-2 expression is found in syncytial tissues such as placenta, muscle, and heart, it is believed that CLASP-2 is involved in mediating or inhibiting cell fusion.

25 The DOCK family has been implicated in the control of cell shape. DOCK1, when transfected into spindle cells, can make them flattened and polygonal (Takai, *et al.*, 1996, *Genomics* 35: 403-303). DOCK1 expression is ubiquitous except in hematopoietic cells. DOCK2 is expressed in hematopoietic cells and when transfected into spindle cells can make them round up (Nishihara, H., 1999, *Hokkaido Igaku Zasshi* 74: 157-66). DOCK2 is 30 expressed in peripheral blood lymphocytes, thymus, spleen, and liver.

5.2.1.7. COILED-COIL

CLASP-2s have the two coiled-coil domains (Lupas *et al.*, 1991, *Science* 252: 1162-64; Lupas, A., 1996, *Meth. Enzymology* 266: 513-525). Coiled-coil domains are known to interact directly with cytoskeleton, indicating that that CLASP-2 proteins interact directly with the cytoskeleton. Thus, it is believed that CLASP-2 binds cytoskeletal proteins, *e.g.*, spectrin, ankyrin, hsp70, talin, ezrin, tropomyosin, myosin, plectin, syndecans, paralemmin, Band 3 protein, Cytoskeletal protein 4.1, Tyrosine phosphatase PTP36 and other molecules.

5.2.1.8. PDZ Ligand

CLASP-2 proteins contain a PDZ-ligand motif ("PBM" or "PL") at the C-terminus of the protein. This short (3 – 8 amino acid) motif mediates the binding of proteins terminating at their carboxyl terminus in the motif (most commonly S/T – X – V – free carboxyl-terminus) to other proteins containing one or more specific PDZ domains (See Songyang *et al.*, 1997, *Science* 275: 72 and Doyle *et al.*, 1996, *Cell* 85: 1067 for a discussion of PDZ-ligand structures).

PDZ domain-containing proteins are involved in the organization of ion channels and receptors at the neurological synapse and in establishing and maintaining polarity in epithelial cells via their binding to the C-termini of transmembrane receptors. It has been shown that PDZ-domain containing proteins can mediate protein-protein interactions in immune system cells (*e.g.*, DLG1 binds to the lymphocyte potassium channel KV1.3 in human T lymphocytes, (Hanada *et al.*, 1997, *J. Biol. Chem.* 272: 26899).

Biochemical evidence that CLASP-2 interacts with the PDZ domains of three closely related proteins is shown in FIG 9A-D. FIG. 9A demonstrates the specificity of the interaction, as the C-terminal 20 amino acids of CLASP-2 bind PSD-95, NeDLG, and DLG1, but not to the PDZ domains of the TIAM-1 protein. FIG. 9B demonstrates the affinity of the interaction. Notably, the highest affinity interaction occurs between CLASP-2 and NeDLG, with a specific binding affinity of at least 10^4 M^{-1} . Affinities in the micromolar range have been found for other biologically important PDZ-ligand interactions. FIG. 9C demonstrates the ability to inhibit CLASP-2 PDZ interactions using either a short fragment of CLASP-2 (the eight C-terminal amino acids) or the C-terminus of KV1.3. As noted above, KV1.3 is known to bind to DLG1 in live lymphocytes. FIG. 9D demonstrates that CLASP-2 and KV1.3 compete for PDZ binding; *i.e.*, not only does KV1.3 block CLASP-2 binding but

CLASP-2 also blocks KV1.3 binding. The ability of the eight C-terminal residues of CLASP-2 to inhibit the interaction of both CLASP-2 and KV1.3 with selected PDZ domains suggests that compounds related to the C-terminal eight-amino acids of CLASP-2, when introduced into cells, will mediate changes in multiple protein-protein interactions involved in the function of lymphoid tissues and other tissues that express these proteins (including heart, lung, and kidney).

Evidence that the C-terminal 8 amino acids of CLASP-2, when introduced into cells, can effect cellular function comes from the experiments in which these amino acids were introduced into cells as a fusion, *e.g.*, with the HIV-derived TAT transporter peptide sequence. Addition of the TAT-CLASP-2 fusion peptide to Jurkat T lymphocytes (compared to controls using the TAT peptide alone) results in subtle, time-dependent alterations in intracellular calcium concentrations as measured using the calcium indicator dye Fluo-4. While these results are consistent with the hypothesis that the TAT-CLASP-2 fusion changes T cell ion fluxes. In particular, the results indicate that the CLASP-2 C-terminal sequence can slightly increase basal intracellular calcium concentrations and can slightly decrease the proportional increase in calcium upon activation of the cells with anti-CD3 antibody. Such changes would be expected for a compound that disrupts localization of the T cell activation-associated CLASP-2 protein and the KV1.3 potassium channel. Small changes in T cell calcium flux can result in large changes in the functional activity of the cells (Wulfig *et al.*, 1997, J. Exp. Med. 185: 1815).

5.2.1.9. Modulation of Immune Responses

CLASP-2 proteins, as described above, modulate immune function in a variety of ways and through a variety of mechanisms (*i.e.*, changing the threshold for T cell activation) by affecting formation of the immunological synapse. Establishment and maintenance of the immunological synapse can involve: (A) signal transduction, (B) cell-cell interactions, and (C) membrane organization.

(A) Signal transduction

Human CLASP proteins, as discussed above, contain SH3 domains and tyrosine phosphorylation sites. These regions have been shown to be involved in signal transduction in a variety of cells including lymphocytes. Thus, human CLASP proteins are believed to interact with these regions during signal transduction events which lead to modulation of immune responses.

CLASP proteins can interact with Tec sub-family of nonreceptor tyrosine kinases. The Tec sub-family of nonreceptor tyrosine kinases consists of Tec, Btk, Tsk/Itk/Emt Itk, and Bmx, and is defined by the presence of SH3 and SH2 domains adjacent to the catalytic domain and an amino-terminal region containing a pleckstrin homology (PH) domain, a Tec homology (TH) domain, and a proline-rich region (Mano, H.; 1999, Cytokine Growth Factor Rev 10: 267-80). The T cell specific Tsk/Itk/Emt, and Btk expressed in most hematopoietic cells other than T cells are important components of antigen receptor signaling pathways in hematopoietic cells.

Btk has been identified as the gene defective in murine X-linked immunodeficiency (*xid*) and human X-linked agammaglobulinemia (XLA) (Nisitani, S., 2000, Proc Natl Acad Sci U.S.A. 97: 2737-42). In *xid* mice, B cell numbers are reduced to one-half of normal and the titers of specific immunoglobulin isotypes are significantly reduced; in addition, *xid* B cells are insensitive to a number of mitogenic stimuli. The human disorder is much more severe, resulting in nearly complete elimination of the B cell compartment and dramatically reduced immunoglobulin levels. Biochemical studies have supported multiple roles for Btk in B cell activation. Btk kinase activity and tyrosine phosphorylation are increased after cross-linking either the B cell receptor on B cells or the high affinity IgE receptor, FcRI, on mast cells. Interleukin-5 and interleukin-6 treatment have also been shown to lead to the activation of Btk.

Itk, like Btk, is tyrosine-phosphorylated upon antigen receptor cross-linking (Mano, H., 1999, Cytokine Growth Factor Rev, 10: 267-80). In addition, peripheral T cells from mice lacking functional Itk are refractory to stimulation by antibodies to CD3 plus antigen presenting cells. These Itk-deficient T cells can be stimulated by phorbol ester and calcium ionophore, demonstrating that Itk acts in signaling pathways proximal to the TCR.

Unlike the related Src family tyrosine kinases including Lyn, Lck, Fyn, ZAP-70, SyK, and CSK, the Tec family kinases lack the amino-terminal myristylation site crucial for the membrane localization of Src family kinases, suggesting that some adaptor proteins are required for the their membrane localization (Mano, H., 1999, Cytokine Growth Factor Rev 10: 267-80). Since all the Tec family kinases contain a proline-rich region which could be bound by a SH3 domain, and since all the human CLASPs contain a SH3 domain, it is believed that human CLASPs could serve as adaptors for the members in the Tec family in different hematopoietic cells.

GTP-binding proteins play an important role in immune response (Mach, B., 1999, Science 285: 1367). A number of biochemical events triggered by TCR/CD3-induced

T cell activation are ablated by agents that modulate the action of G proteins. Pertinent to this is the ability of cholera toxin to inhibit the cellular proliferation and intracellular Ca^{2+} mobilization that is mediated by anti-CD3 antibody treatment of T cells. The G protein competitive inhibitor GDPS, can impede the extent of inositol phosphates generated upon stimulation in peripheral T lymphocytes. Nonhydrolyzable analogs of GTP, such as GTPS, or other agents such as ALF that activate G proteins by circumventing the need for receptor engagement, can result in T cell activation.

The Guq/11 subfamily (Stanners, J., 1995, J Biol Chem 270: 30635-42) and Rap1 (Lafont, V., 1998, Biochem Pharmacol 55: 319-24) of GTP-binding proteins have been shown to be involved in human T cell receptor/CD3-mediated signal transduction pathway. Also, Cdc42, a Rho family small GTPase, is known to play a critical role in the formation of actin microspikes in response to external stimuli (Miki, H.; 1998, Nature, 391: 93-6). Interestingly, a Cdc42 binding protein, WASP, has a proline-rich domain which could interact with the SH3 domain present in all the human CLASPs. Human CLASPs may interact with these GTP-binding proteins.

Several adaptor proteins including NCK, CBL (Bachmaier, K., 2000 Nature 403: 211-6), SHC, LNK, SLP-76, HS1, SIT, VAV, GrB2, and BRDG1, and two tyrosine phosphatases, EZRIN, SHP-1 and SHP-2 have been shown to interact with ITAM or SH3 domains. These proteins may also interact with CLASP-2. Several proteins have been shown to interact with ITAM or SH3 domains and may also interact with CLASP-2. These include adaptor proteins such as NCK, CBL (Bachmaier, K., 2000, Nature 403: 211-6), SHC, LAT, LNK, SLP-76 (Krause M *et al.*, 2000, J Cell Biol 149: 181-94), HS1, SIT, VAV, GrB2 (Zhang W. and Samelson, L.E., 2000, Semin Immunol 12: 35-41), and BRDG1, kinases such as SYK and LCK, and tyrosine phosphatases such as SHP-1 and SHP-2. These interactions can be defined by a number of different biochemical or cell biological methods including in vitro binding assays, co-immunoprecipitation assays, co-immunostaining (Harlow, E. and Lane, D., 1999, Using Antibodies: A laboratory Manual. Cold Spring Harbor Press) or genetic assays such as yeast the yeast two hybrid system, in which a CLASP-2 protein or fragment can be used as "bait" (Zervos *et al.*, 1993, Cell 72: 223-232; Madura *et al.*, 1993, J. Biol. Chem 268: 12046-12054).

Other assays include in vitro binding assays, co-immunoprecipitation assays, co-immunostaining assays, and yeast two hybrid system screening assays in which a CLASP-2 domain or fragment can be used as "bait" or "trap" protein (Zervos *et al.* (1993), Cell 72: 223-232; Madura *et al.* (1993) J. Biol. Chem. 268: 12046-12054).

In other embodiments, CLASP polypeptides are transfected into lymphocytes. After transfection, a variety of standard assays can be used to evaluate, for example, CLASP modulation of T cell activation. These assays include calcium influx assays, NF-AT nuclear translocation assays (*e.g.*, Cell, 1998, 93: 851-61), NF-AT/luciferase reporter assays (*e.g.*, MCB 1996 16: 7151-7160), tyrosine phosphorylation of early response proteins such as Hs1, PLC- γ , ZAP-70, and Vav (*e.g.*, J. Biol. Chem. 1997, 272: 14562-14570).

(B) Cell-Cell Interaction

As discussed above, human CLASP proteins are homologues of E-cadherin. As shown in FIG. 1, CLASP-2 contains both a cadherin cleavage domain and a cadherin ectodomain. Therefore CLASP-2 proteins may interact with cadherins through these domains. The cadherins constitute a family of cell surface adhesion molecules that are involved in calcium-dependent cell to cell adhesion. Human cadherins, E-, P- N- and VE-cadherin, have a restricted tissue distribution: E- and P-cadherin are expressed in epithelial tissues, N-cadherin is found mainly on neural cells, and VE-cadherin is found on vascular endothelium. Homophilic binding between cadherins on adjacent cells is vital for the maintenance of strong cell to cell adhesion in these tissues. For example E-cadherin is required for the formation of adherens junctions between mature epithelial cells and is involved in Langerhans cell adhesion to keratinocytes, and VE-cadherin is needed for the maintenance of lateral association between endothelial cells. The extracellular regions of mature mammalian cadherins are comprised of five "CAD" modules of approximately 1110 amino acids. Crystallographic and biochemical studies indicate that cadherins can form dimers on the cell surface, and that interaction with dimeric cadherins on opposing cell surfaces can lead to the formation of "zipper-like" cell junctions.

The integrins are a second family of transmembrane adhesion molecules that are involved in both cell to cell and cell to matrix interactions. At least 15 chains associate with 8 chains to form a large number of heterodimeric integrins that can be classified into several major subfamilies based on their shared use of a particular chain. Members of three subfamilies, the 1, 2, and 7 integrins, are commonly found on leukocytes. The expression of 1 integrins is widespread (for example, 51, CD49e/CD29, is found on T cells, granulocytes, platelets, fibroblasts, endothelium, and epithelium), whereas the 2 and 7 integrins have a restricted pattern of expression.

Interestingly, E-cadherin on human epithelial cells has been found to be a ligand for the mucosal lymphocyte integrin, E7, and a similar interaction has been indicated

in the mouse. Monoclonal antibodies to E-cadherin or to E7 block IEL adherence to epithelial cells, and transfection of cells with E7 confers upon them the ability to adhere to cells transfected with E-cadherin.

L929 cells can be transfected with CLASP-2 and Neomycin. G418-resistant clones can be screened for CLASP-expression with anti-CLASP peptide-specific antibodies. CLASP-expressing clones can be used to test for homotypic and/or heterotypic calcium dependent cell adhesion using the "cell aggregation assay" described for cadherin molecules (Murphy-Erdosh, C. *et al.*, 1995, J. Cell Biol. 129: 1379-1390).

Several approaches can be used to identify the amino acids involved in the binding domains. Soluble fusion molecules (*e.g.*, EC12-IgG, ECC-IgG, ECM-IgG, and GST-EC12), peptides, and peptide-specific anti-CLASP antibodies are available for blocking experiments in the above-described assay. Transfectants generated by site-directed mutagenesis can also be used.

(C) Membrane Anchoring/Cytoskeletal Interactions

Interestingly, tyrosine-phosphorylated ITAMs interact with actin cytoskeleton upon activation of mature T lymphocytes (Rozdzial, M. M., 1995, Immunity 3: 623-633). Since human CLASPs contain both ITAMs and coiled-coil domains which have been shown to interact with cytoskeletal proteins, CLASPs are believed to play an important role in modulating cell surface molecule expression by re-organizing cytoskeletal structure.

F-actin microfilament cytoskeletal organization has been known to be involved in the modulation of cell surface molecule expression. WASP, a GTPase-binding protein, plays a critical role in the formation of actin microspikes in response to external stimuli and ectopic expression of WASP induces the formation of F-actin filament clusters that overlap with the expressed WASP itself. Another WASP family protein, N-WASP, has also been shown to play important roles in filopodium formation. Both of these proteins cause actin polymerization, but with different features when they are expressed in cells; WASP mainly localizes at perinuclear areas and causes actin clustering, but most N-WASP is present at plasma membranes and induces filopodium formation (Miki, H.; 1998, Nature 391: 93-6). Both WASP and N-WASP, contain a proline-rich domain which could interact with the SH3 domain present in all the human CLASPs. CLASP-2 may interact with F-actin filament through CLASP-2 binding to WASP or WASP-like proteins.

Standard assays can be used for detecting CLASP protein interaction with cytoskeletal proteins. These assays include co-sedimentation assays, far western blot analysis (Ohba, T., 1998, Anal. Biochem. 262: 185-192), surface plasmon resonance, F-actin staining

with phalloidin in CLASP-transfected lymphocytes (*e.g.*, Small, J. *et al.* 1999, Microsc. Res. Tech. 4: 3-17), and immunocytochemical analysis of subcellular distribution of focal adhesion proteins (such as paxillin, tensin, vinculin, talin, and FAK in CLASP-transfected lymphocytes; see, *e.g.*, Ridyard, M.S., 1998, Biochem. Cell Biol. 76: 45-58).

5 5.2.2. CLASP-2 Exon Structure and Genomic Domains

Alternative splicing is likely to represent a regulatory switch that governs different functions of CLASP-2 in immune responses. Additionally, alternative splice variants affecting the untranslated regions of an RNA can be a way of regulating RNA stability.

10 As noted *supra*, CLASP-2 gene expression is characterized by alternative exon usage. Intron/exon structure can be predicted by computer analysis of genomic DNA, however, splice junctions and alternative splicing can only be elucidated by comparison of genomic clones to cDNA clones. Alternative splicing and RNA editing are mechanisms generate a variety of proteins from the same gene. An example for how alternative splicing is used to generate thousands of different proteins from only a few genes is represented by the 15 Neurexin gene family (for review of Neurexins, see Missler M. and Suedhof, T., 1998, Trends in Genetics, 14: 20-25). Comparative analysis of CLASP-2 genomic clones and cDNA clones revealed that CLASP-2 is composed of numerous exons and that distinct CLASP-2 transcripts are generated by alternative splicing. The protein encoding portion of 20 CLASP-2 is covered by at least 14 exons (FIG. 6A).

Numerous diseases are caused or are thought to be caused by splice site mutations that can cause exon skipping or otherwise result in a truncated protein product. Some of these diseases include, *e.g.*, Marfan Syndrome (Liu W, *et al.*, 1997, Nat. Genet. 16: 328-9), Hunter disease (Bonucelli G, *et al.*, 2000, Hum. Mutat. (Online) 2000 15(4): 389, 25 Duchenne muscular dystrophy (Wibawa T, *et al.*, 2000, Brain Dev. 22(2): 107-112), Myelomonocytic leukemia (Wutz D, *et al.*, 1999, Leuk. Lymphoma 35: 491-9.), and Isovaleric acidemia (Vockley J, *et al.*, 2000, Am. J. Hum. Genet. 66: 356-67). This is especially true for genes composed of many exons (such as CLASP-2). The genomic sequence around CLASP-2 exon/intron boundaries is useful for diagnostic approaches 30 towards the identification of diseases caused by splice site mutations. The abundance or presence of CLASP-2 isoforms in cell populations (*e.g.*, hematopoietic cells, lymphocytes) is correlated with a disease state by comparing the abundance of CLASP-2 in cells from

subjects suffering from the disease with the level of CLASP-2 in cells from healthy subjects. This can be accomplished by utilizing any number of assays (e.g., PCR).

Alignment of the CLASP-2 intron/exon splice sites with the CLASP-2 protein sequence and the finding of conserved exon/intron boundaries within the CLASP gene family (FIG. 6) suggest that specific CLASP-2 exons encode functionally distinct protein domains (see FIG. 6 and Example 4). ITAM and DOCK motifs 1 and 2 are encompassed by splice sites (amino acid residues 946 and 1063); DOCK motif 3 and COILED-COIL motif 1 and 2 are also encompassed by splice sites (amino acid residues 1102, 1170 and 1246, respectively).

CLASP-2 alternative transcripts are summarized in FIG. 3 and FIG. 11B. Briefly, one alternative exon missing in CLASP-2A is present in CLASP-2B and CLASP-2D. This exon contains the DNA portion encoding the ITAM motif and DOCK motif 1. The CLASP-2D protein product does not contain the C-terminal 38 amino acids of CLASP-2A and CLASP-2B: Thus, a PDZ binding motif (SSVV; amino acid residue 1286 through 1289) that is only present in the CLASP-2A/B-specific C-terminal end is missing in the CLASP-2D gene product. The presence or absence of this PDZ binding motif can be attributed to alternative RNA processing. Additionally, a CLASP-2 alternative transcript has been found that deletes nucleotides 209-291 that results in a premature stop codon. The protein encoded by this transcript appears to be a soluble form of CLASP-2 that may regulate (e.g., is an antagonist or an agonist) the function other CLASP family members and isoforms.

5.2.3. CLASP Superfamily Members

As is illustrated in FIG. 5, CLASP-2 is a member of a superfamily of immune-cell associated proteins with similar motifs. CLASP-1 was described in U.S.S.N. 09/411,328, filed October 1, 1999. CLASP-1 uniquely among the known CLASPs contains SH3 binding domain motifs. CLASP-2A, -B, -C, and -E polypeptides have no adaptor binding sites or SH3 binding domains found in CLASP-1. CLASP-3, CLASP-4, CLASP-5 and CLASP-7 are described in copending U.S.S.N. 60/182,296, filed February 14, 2000, and which is incorporated by reference herein in its entirety for all purposes.

5.3. CLASP-2 mRNA Expression

As described in Example 2, CLASP-2 mRNA expression was assayed in tissues and cell lines by Northern analysis. The results are shown in FIG. 4A and B. The

results of Northern Analysis of CLASP-2 expression and expression of other members of the CLASP family are summarized in Table 2.

Table 2

Tissue/Cell Line ¹	CLASP					
	1	2 ^{3,4}	3	4	5	7
PBL	+ ²	-	-	+++	++	-
Lung	-	+	-	-	-/+	+++
Placenta	-/+	+++	+	-/+	+	+
Sm Intestine	-/+	-	-	-	-/+	+
Liver	-/+	-/+	-/+	-	-/+	+
Kidney	-/+	+	+++	-/+	+	++
Spleen	++	-	-	-/+	+	-/+
Thymus	++	-	-	-/+	+	-
Colon	-	-	-	-	-	-
Skel Muscle	-	-/+	++	-	-	-/+
Heart	-/+	++	+++	-/+	-	+++
Brain	+++	-/+	-/+	-	-	-
Jurkat	++	++	++	+	-	-
MV411	++	-	++	+	+	+
THP1	++	-	-	-	-	-/+
HL60	-	-	-	-	-/+	-
9D10	++	++ ⁵	+	+	+	+
3A9	+	-/+	-	-	-	-
CH27	+	-/+	-	-	-	-
293	-	++	+++	+	-	+

1. Jurkat = human T cell line; MV4-11 = B myelomonocyte; 9D10 = B cell line; THP-1 = monocyte; 3A9 = mouse T cell; CH27 = mouse B cell line; HL60 = human promyelocyte; 293 = embryonic kidney epithelial cells (293)

2. Table Legend (based on Northern blot results): - = no expression; -/+ = low expression; + = medium expression; ++ medium high expression; +++ high expression.

3. A CLASP-2 EST (EST 815795) was identified from a bone marrow cDNA library.

4. The probe used (HC2.2) did not distinguish between CLASP-2A, -2B, -2C and 2D.. This probe encompasses nucleotides 3920 to 4650 (731 bp long) from CLASP-2A cDNA.

5. In RNA from 9D10, the major transcript runs substantially shorter than the major transcripts seen in Jurkat and 293 cells; however, the longer transcript is also present in 9D10. Hybridization of probe HC2.2 with 9D10 total RNA reveals at least 3 different transcripts. See FIG. 4B

As indicated in Table 2 and shown in FIG. 4, CLASP-2 is expressed most strongly in placenta followed by lung, kidney and heart; CLASP-3 is expressed strongly in kidney and heart, and less strongly in placenta and skeletal muscle ; CLASP-4 is expressed exclusively in peripheral blood lymphocytes; CLASP-5 is expressed strongly in peripheral blood leukocytes, present in placenta, kidney; spleen and thymus, and weakly in lung, small

intestine and liver. It is not expressed in brain, heart, skeletal muscle and large intestine; CLASP-7 is expressed strongly in lung, heart, liver and kidney, but not in PBL, brain or thymus.

Differences in tissue expression patterns for different CLASP proteins indicate
5 different CLASPs have differential roles in immune function and, accordingly, can be separately targeted to achieve different functions. For example, since CLASP proteins are necessary for proper function or signaling by the T cell receptor (TCR), the tissue specific distribution of different CLASPs permits differential modulation of the immune response in different tissues. Since CLASP-2 is present in heart, blocking CLASP-2 function or
10 expression is useful to selectively block immune response in the heart (for example, to selectively stop immune response in the heart compartment, *e.g.*, following cardiac transplant rejection or post-MI inflammation, without compromising immunity elsewhere. Similarly, blocking CLASP-3 can block rejection of the kidney following kidney transplant. Furthermore, by adjusting the level of inhibition, the degree of immune blockage versus
15 response can be modulated in the compartments represented by each CLASP.

5.4. CLASP-2 Polynucleotides And Methods Of Use

The present invention provides a variety of CLASP-2 polynucleotides and methods for using them. In one aspect, the polynucleotide of the invention encodes a polypeptide comprising at least a fragment (*e.g.*, an immunogenic fragment) of a CLASP-2
20 protein (*e.g.*, at least a fragment of SEQ. ID. NO: 2, 4, 6 or 10) or variant thereof. In another aspect, the molecules that comprise a CLASP-2 polynucleotide that, while not necessarily encoding a CLASP-2 protein or fragment, is useful as a probe or primer for detecting CLASP-2 expression, for inhibition of CLASP-2 expression (*e.g.*, antisense or ribozyme-mediated inhibition), for gene knockout, and the like.

5.4.1. CLASP-2 Polynucleotides

The invention also provides isolated or purified nucleic acids having at least 8 nucleotides (*i.e.*, a hybridizable portion) of a CLASP-2 sequence or its complement; in other embodiments, the nucleic acids consist of at least about 25 (continuous) nucleotides, about 50 nucleotides, about 100 nucleotides, about 150 nucleotides, about 200 nucleotides, about 250
30 nucleotides, about 500 nucleotides, about 550 nucleotides, about 600 nucleotides, or about 650 nucleotides or more of a CLASP-2 sequence, or a full-length CLASP-2 coding sequence. In another embodiment, the nucleic acids are smaller than about 35, about 200 or about 500

nucleotides in length. Polynucleotides can be single or double stranded, and may be DNA, RNA, PNA or a hybrid molecule.

In specific aspects, nucleic acids are provided which comprise a sequence complementary to at least about 10, 25, 50, 100, 150, 200, 250, 500, 550, 600, or 650 nucleotides or the entire coding region of a CLASP-2 coding sequence. Usually, the isolated polynucleotide is less than about 100 kbp, generally less than about 50 kbp, and often less than about 20 kbp, less than about 10 kbp, less than about 5 kbp, or less than about 1000 nucleotides in length.

In a specific embodiment, a nucleic acid that is hybridizable to a CLASP-2 nucleic acid or its complement, or to a nucleic acid encoding a CLASP-2 derivative, under conditions of low stringency is provided. Derivatives of CLASP-2 contemplated include, but are not limited to, splice variants of a gene encoding a CLASP-2, other members of a CLASP-2 gene family which differ from one of the CLASP-2 nucleotide or amino acid sequences disclosed herein by the insertion or deletion of one or several domains, and the like.

In one embodiment, the CLASP-2 polynucleotide is identical or exactly complementary to SEQ. ID NO: 1, 3, 5 or 9 or selectively hybridizes to an aforementioned sequence. In various embodiments, the polynucleotide is identical or exactly complementary to, or selectively hybridizes to, the nucleotide sequence encoding a particular protein domain or region, or a particular gene exon of the CLASP-2 mRNA or genomic sequence. Such polynucleotides are particularly useful as probes, because they can be selected to identify a defined species of CLASP-2.

In addition to the polypeptide and polynucleotide sequences specifically exemplified herein, the invention contemplates CLASP-2 homologues from other species, allelic and splice variants, and other variants disclosed herein. The CLASP-2 gene exhibits evidence of alternative splicing of transcripts.

For example, CLASP-2A and CLASP-2C are related to each other as apparent splice variants, with CLASP-2C containing an exon not found in CLASP-2A. The exon sequence is 5'-AGG GAT TTT GAG AGG CTG GCC CAT CTG TAT GAC ACG CTG CAC CGG GCC TAC AGC AAA GTG ACC GAG GTC ATG CAC TCG GGC CGC AGT TNC TGG GGA CCT ACT TCC GGG TAG CCT TCT TCG GGC AG-3' (encoding the peptide sequence: RDFERLAHLYDTLHRAYSKVTEVMHSGRLLGTYFRVAFFGQGF). It will be apparent to one of skill that, by using polynucleotide probes or primers corresponding to the nucleic acid sequence above, or by using antibodies that specifically

recognize the peptide above, or those polynucleotide probes or primers shown in Table 3 below, it is possible to distinguish between different CLASP isoforms (e.g., to detect differential expression).

Table 3

	Found in/will detect	Exemplary Probe/Primer (5' - 3')	Notes/Comments
1	full length hC2A	F1: CCCAGATTTTATGATGAG R1: GATAATGACAAAGTTCTGAC	
2	full length hC2D	F2: CTGGAAATCTTGACAAAAATGC R2: GTCTTTTAAATACAGATGTGG	
3	hC2B, hC2C, hC2E	F3: GAGAGGCTGGCCCATCTGTATG R3: ATCTTCAAAGAATCCCTGCC	Distinction based upon product size differences following PCR
4	hC2D	F4: GAAGCAGTCCAGTGGGAGCCG R4: GCCTCCCCGGCTCCTCCTCAGG	Recognizes hC2D-specific insertion
5	hC2D	F3: GAGAGGCTGGCCCATCTGTATG R5: CCTCCACATCTGTTCACTGTC	
6	hC2E	F5: CTCCATGATGGAAGACGTGGG R6: GATGAGCTCGTAGCGTCCGGC	Spans deletion unique to hC2E. Distinction based upon product size differences following PCR
7	hC2B	F6: CATTGGCGTTTAAAGCTCCTG R3: ATCTTCAAAGAATCCCTGCC	F6 primer spans deletion unique to hC2E
8	hC2A	F7: GGACCCATAGTTTCATGATCG R4: CTCATCTTCAAGAAATCCCTC	R4 primer spans the region where other CLASPs have an insert

5

5.4.1.1. Substantial Identity

In some embodiments, the CLASP-2 polynucleotides of the invention are substantially identical to SEQ ID NOs: 1, 3, 5, or 9, or to a fragment thereof.

10 An indication that two nucleic acid sequences are substantially identical is that the two polynucleotides have a specified percentage sequence identity *e.g.*, usually at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98 identity over a specified region when optimally aligned.

15 Another indication that two nucleic acid sequences are substantially identical is that a polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication

that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below.

Yet another indication that two nucleic acid sequences are substantially identical (*e.g.*, a naturally occurring allele of the CLASP-2 sequence of SEQ ID NO: 1) is that the same primers can be used to amplify the sequence. For example, CLASP-2 polynucleotides can be PCR amplified from cDNA derived from human lymphocytes using the primer pairs shown in Table 3.

The primers of Table 3 are also useful for amplification of CLASP-2 splice variants. Another indication that two nucleic acid sequences are substantially identical is that they selective hybridize under stringent conditions (*i.e.*, one sequence hybridizes to the complement of the second sequence), as described *infra*.

5.4.1.2. Selective Hybridization

The invention also relates to nucleic acids that selectively hybridize to exemplified CLASP-2 sequences (including hybridizing to the exact complements of these sequences). Selective hybridization can occur under conditions of high stringency (also called "stringent hybridization conditions"), moderate stringency, or low stringency.

5.4.1.2.1. High Stringency

"Stringent hybridization conditions" are conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but not to other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Probes, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m , 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (*e.g.*, 10 to 50 nucleotides)

and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For high stringency hybridization, a positive signal is at least two times background, preferably 10 times background hybridization. Exemplary high stringency or stringent hybridization conditions include: 50% formamide, 5x SSC and 1% SDS incubated at 42° C or 5x SSC and 1% SDS incubated at 65° C, with a wash in 0.2x SSC and 0.1% SDS at 65° C. In a specific embodiment, a nucleic acid which is hybridizable to a CLASP-2 nucleic acid under the following conditions of high stringency is provided: Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 8-16 h at 65°C in prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20 X 10⁶ cpm of ³²P-labeled probe. Washing of filters is done at 65°C for 15-30 h in a solution containing 2X SSC, 0.1% SDS. This is followed by a wash in 0.2X SSC and 0.1% at 50°C for 15-30 min before autoradiography.

5.4.1.2.2. Moderate Stringency

In another specific embodiment, a nucleic acid, which is hybridizable to a CLASP-2 nucleic acid under conditions of moderate stringency is provided. Examples of procedures using such conditions of moderate stringency are as follows: Filters containing DNA are pretreated for 6 h at 55°C in a solution containing 6X SSC, 5X Denhart's solution, 0.5% SDS and 100 µg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution and 5-20 X 10⁶ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 12-16 h at 55°C, and then washed twice for 30 minutes at 50°C in a solution containing 1X SSC and 0.1% SDS. Filters are blotted dry and exposed for autoradiography. Other conditions of moderate stringency which can be used are well-known in the art. Washing of filters is done at 45°C for 1 h in a solution containing 0.2X SSC and 0.1% SDS.

5.4.1.2.3. Low Stringency

By way of example and not limitation, procedures using such conditions of low stringency are as follows (see also Shilo and Weinberg, 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 6789-6792): Filters containing DNA are pretreated for 6 h at 40 C in a solution

containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 g/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20 X 10⁶ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 h at 40 C, and then washed for 1.5 h at 55 C in a solution containing 2X SSC and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 30 minutes at 50-55°C. Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 60-65°C and reexposed to film. Other conditions of low stringency that can be used are well known in the art (e.g., as employed for cross-species hybridizations).

5.4.1.3. CLASP-2 Variants and Fragments

The CLASP-2 variants of the invention can contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. CLASP-2 polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as E. coli).

Exemplary CLASP-2 polynucleotide fragments are preferably at least about 15 nucleotides, and more preferably at least about 20 nucleotides, still more preferably at least about 30 nucleotides, and even more preferably, at least about 40 nucleotides in length, or larger 50, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650 nucleotides. In one embodiment, exemplary fragments include fragments having at least a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600 to the end of SEQ ID NO: 1 or SEQ ID NO: ___ or comprising the cDNA coding sequence in the deposited clones. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein.

In other embodiments, CLASP-2 polynucleotides of the invention are other than SEQ ID NO:1 or fragments of SEQ ID NO:1.

As shown in FIG 11 above, there are at least three CLASP-2 full length cDNA isoforms (A + Z, B + Z, and C + Z). Each of the isoforms uses a unique first exon (labelled exon 1A, 1B, and 1C) (see FIG. 11 and Table 4 below).

Table 4: CLASP-2 Isoforms

CLASP-2 Isoform	FIG 11C Schematic	Nucleotides
Isoform 1	A + Z	-182 to 6690
Isoform 2	B + Z	-219 to 6690
Isoform 3	C + Z	-143 to 6690

5

In one embodiment, the CLASP-2 polynucleotide has the sequence shown in FIG. 11 (Isoform 1, Isoform 2, or Isoform 3 as indicated in Table 4 above) or a fragment of the sequence shown in FIG. 11 comprising at least about 1, 5, 10, 25 or 50 or more contiguous nucleotides from nucleotides -182 to 1883 of Isoform 1, nucleotides -219 to 1883 of Isoform 2, or nucleotides -143 to 1883 of Isoform 3.

10

In another embodiment, CLASP-2 primers or probes comprise at least about 5, 10, 25 or 50 or more contiguous nucleotides from nucleotides -182 to 1883 of Isoform 1, nucleotides -219 to 1883 of Isoform 2, or nucleotides -143 to 1883 of Isoform 3 as shown in FIG. 11 and Table 4 above alone or in combination with SEQ ID NO:1 or a fragment of SEQ ID NO:1.

15

In an aspect, the invention provides antibodies or binding fragments that bind the CLASP-2 isoforms 1-3. In another embodiment, the invention provides antibodies that specifically bind to the CLASP-2 isoforms shown in FIG. 11 but not to the polypeptide encoded by SEQ ID NO:1

20

In one embodiment, the CLASP-2 variants differ from those shown in FIG. 1 or FIG. 11 (SEQ ID NOs 1, 3, 5, 7 9, _____) by virtue of incorporating a different combination of exons than found in the exemplified sequences. For example, 81g01 (Genbank Accession Number AF85864; Locus.HUMYN81g01; 526 bp; EST sequence submitted August 29, 1998 by Washington University at St. Louis; see FIG. 3A and FIG. 3B) is a variant of hCLASP-2 on the basis of CLASP-2 identity along certain stretches of the sequence. From 5' to 3', it begins with a 315 nucleotide stretch which is identical to CLASP-2A. It then has a gap relative to CLASP-2A that is identical to the GAP in another CLASP-2 isoform, hCLASP-2D (KIAA1058). In place of that gap, a 16 amino acid insert (48 nucleotides) is present which is not found in other isoforms, followed by another

25

approximately 150 bp stretch of nucleotides once again identical to CLASP-2A. This is characteristic of an alternate splice due to the specific sequence identity on both sides of a differential stretch of nucleotides.

Using known methods of protein engineering and recombinant DNA technology, variants can be generated to improve or alter the characteristics of the CLASP-2 polypeptides. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the CLASP-2 protein without substantial loss of biological function.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities can still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes CLASP-2 polypeptide variants which show biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, Science 247: 1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicate that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at 30 specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells,

1989, Science 244: 1081-1085) The resulting mutant molecules can then be tested for biological activity.

In various embodiments, CLASP-2 polynucleotide fragments include coding regions for, or regions hybridizable to, the CLASP-2 structural or functional domains described *supra*. As set out in the Figures, such preferred regions include the following domains/motifs: ITAM, DOCK, COILED/COILED, and PBM. Thus, for example, polypeptide fragments of CLASP-2 as shown in FIG. 1 and FIG. 11-(SEQ ID NO: 2, 4, 6 10, _____) falling within conserved domains are specifically contemplated by the present invention (see FIG. 3). Moreover, polynucleotide fragments encoding these domains are also contemplated. Such polypeptide fragments find use, for example, as inhibitors of CLASP-2 function in CLASP-2-expressing cells.

5.4.2. Uses of CLASP-2 Polynucleotides

The CLASP-2 polynucleotides of the invention are useful in a variety of applications. In one aspect of the invention, the polypeptide-encoding CLASP-2 polynucleotides of the invention are used to express CLASP-2 polypeptides (*e.g.*, as described herein) for example to produce anti-CLASP-antibodies or for use as therapeutic polypeptides. In another aspect, the CLASP-2 polynucleotide or fragments thereof can be used for diagnostic purposes (*e.g.*, as probes for CLASP-2 expression). In particular, since CLASP-2s can be expressed in lymphocytes, a CLASP-2 polynucleotide can be used to detect the expression of CLASP-2 as a lymphocyte marker. For diagnostic purposes, a CLASP-2 polynucleotide can be used to detect CLASP-2 gene expression or aberrant CLASP-2 gene expression in disease states. In another aspect, the CLASP-2 polynucleotide or fragments are used for therapeutic purposes. For example, included in the scope of the invention are methods for inhibiting CLASP-2 expression, *e.g.*, using oligonucleotide sequences, such as antisense RNA and DNA molecules and ribozymes, that function to inhibit expression of CLASP-2. In another aspect, CLASP-2 polynucleotides can be used to construct transgenic and knockout animals, *e.g.*, for screening of CLASP-2 agonists and antagonists. In another aspect, CLASP-2 polynucleotides can be used for screening of CLASP-2 agonists and antagonists.

5.4.2.1. Use of CLASP-2 Polynucleotides for Detection, Diagnosis, and Treatment

The CLASP-2 polynucleotides of the invention are useful for detection of CLASP-2 expression in cells and in the diagnosis of diseases or disorders (*e.g.*,

immunodeficient states) resulting from aberrant expression of CLASP-2. Aberrant expression of CLASP-2 mRNA or protein means expression in lymphocytes (*e.g.*, T lymphocytes or B lymphocytes) or other CLASP-2 expressing cells of at least 2-fold, preferably at least 5-fold greater or less than expression in control lymphocytes obtained from a healthy subject. CLASP-2 polypeptide expression is easily measured by ELISA using anti-CLASP-2 antibodies of the invention. CLASP-2 mRNA expression (including expression of specific species or splice variants of CLASP-2) can be measured by quantitative Northern analysis or quantitative PCR, LCR, or other methods, using the probes and primers of the invention.

In one embodiment, the assays of the present invention are amplification-based assays for detection of an CLASP-2 gene product. In an amplification based assay, all or part of a CLASP-2 mRNA or cDNA (hereinafter also referred to as "target") is amplified, and the amplification product is then detected directly or indirectly. When there is no underlying gene product to act as a template, no amplification product is produced (*e.g.*, of the expected size), or amplification is non-specific and typically there is no single amplification product. In contrast, when the underlying gene or gene product is present, the target sequence is amplified, providing an indication of the presence and/or quantity of the underlying gene or mRNA. Target amplification-based assays are well known to those of skill in the art.

The present invention provides a wide variety of primers and probes for detecting CLASP-2 genes and gene products. Such primers and probes are sufficiently complementary to the CLASP-2 gene or gene product to hybridize to the target nucleic acid. Primers are typically at least 6 bases in length, usually between about 10 and about 100 bases, typically between about 12 and about 50 bases, and often between about 14 and about 25 bases in length, often PCR primers of 15-30 (*e.g.*, 18-22 nucleotides) are used. However, the length of primers can be adjusted by one skilled in the art. One of skill, having reviewed the present disclosure, will be able, using routine methods, to select primers to amplify all, or any portion, of the CLASP-2 gene or gene product, or to distinguish between variant gene products, CLASP-2 alleles, and the like. Single oligomers (*e.g.*, U.S. Pat. No. 5,545,522), nested sets of oligomers, or even a degenerate pool of oligomers can be employed for amplification.

It will be appreciated that probes and primers can be selected to distinguish between species and splice variants based on the guidance of this disclosure, by targeting

primers or probes to differentially used exons (or exon-exon junctions that differ between variants).

Methods can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (*e.g.*, genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to an CLASP-2 gene under conditions such that hybridization and amplification of the CLASP-2-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. See U.S. Pat. Nos. 4,683,195 and 4,683,202, Landegran *et al.*, 1988, Science 241: 1077-1080; Nakazawa *et al.*, 1994, Proc. Natl. Acad. Sci. U.S.A. 91: 360-364, Abravaya *et al.*, 1995, Nucleic Acids Res. 23: 675-682).

Because CLASP-2 gene products are expressed in the immune system (*e.g.*, T lymphocytes, B lymphocytes and macrophages), expression will be typically assayed in these cells. Methods which are well known to those skilled in the art can be used to isolate lymphocytes, macrophages, and alike (*See, e.g.*, Coligan, J. E., *et al.* (eds.), 1991, Current Protocols in Immunology, John Wiley & Sons, NY; this reference is incorporated by reference for all purposes). In one embodiment, assays are carried out on biopsy or autopsy-derived tissue.

In various embodiments, CLASP-2 gene expression is detected by hybridization of a detectable probe to mRNA or cDNA obtained from cells (*e.g.*, lymphocytes). A variety of methods for specific DNA and RNA measurement using nucleic acid hybridization techniques are known to those of skill in the art (see Sambrook *et al.*, *supra*). Hybridization based assays refer to assays in which a probe nucleic acid is hybridized to a target nucleic acid, forming a hybridization complex. Usually the nucleic acid hybridization probes of the invention are entirely or substantially identical to a contiguous sequence of the CLASP-2 gene or RNA sequence. Preferably, nucleic acid probes are at least about 50 bases, often at least about 20 bases, and sometimes at least about 200 bases, at least about 300-500 nucleotides or more in length. Various hybridization techniques are well known in the art, and are in fact the basis of many commercially available diagnostic kits.

Methods of selecting nucleic acid probe sequences for use in nucleic acid hybridization are discussed in Sambrook *et al.*, *supra*. In some formats, at least one of the target and probe is immobilized. The immobilized nucleic acid can be DNA, RNA, or another oligo- or poly-nucleotide, and can comprise natural or non-naturally occurring

nucleotides, nucleotide analogs, or backbones. Such assays can be in any of several formats including: Southern, Northern, dot and slot blots, high-density polynucleotide or oligonucleotide arrays (e.g., GeneChips™ Affymetrix), dip sticks, pins, chips, or beads. All of these techniques are well known in the art and are the basis of many commercially available diagnostic kits. Hybridization techniques are generally described in Hames *et al.*, ed., 1985, *Nucleic Acid Hybridization, A Practical Approach* IRL Press; Gall and Pardue, 1969, *Proc. Natl. Acad. Sci. U.S.A.*, 63: 378-383; and John *et al.*, 1969, *Nature*, 223: 582-587.

A variety of nucleic acid hybridization formats are known to those skilled in the art. For example, one common format is direct hybridization, in which a target nucleic acid is hybridized to a labeled, complementary probe. Typically, labeled nucleic acids are used for hybridization, with the label providing the detectable signal. One method for evaluating the presence, absence, or quantity of CLASP-2 mRNA is carrying out a Northern transfer of RNA from a sample and hybridization of a labeled CLASP-2 specific nucleic acid probe. A useful method for evaluating the presence, absence, or quantity of DNA encoding CLASP-2 proteins in a sample involves a Southern transfer of DNA from a sample and hybridization of a labeled CLASP-2 specific nucleic acid probe.

Other common hybridization formats include sandwich assays and competition or displacement assays. Sandwich assays are commercially useful hybridization assays for detecting or isolating nucleic acid sequences. Such assays utilize a "capture" nucleic acid covalently immobilized to a solid support and a labeled "signal" nucleic acid in solution. The biological or clinical sample will provide the target nucleic acid. The "capture" nucleic acid and "signal" nucleic acid probe hybridize with the target nucleic acid to form a "sandwich" hybridization complex. To be effective, the signal nucleic acid cannot hybridize with the capture nucleic acid.

In one embodiment, CLASP-2 polypeptides or polynucleotides are useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the activation, differentiation of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders can be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious.

In another embodiment, CLASP-2 polynucleotides or polypeptides are useful in treating or detecting deficiencies or disorders of hematopoietic cells. CLASP-2

polypeptides or polynucleotides could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g., agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

In one embodiment, CLASP-2 polynucleotides or polypeptides are useful in treating or detecting autoimmune diseases. The term "autoimmune disease" as used herein has the normal meaning in the art and refers to a spontaneous or induced malfunction of the immune system of mammals in which the immune system fails to distinguish between foreign immunogenic substances within the mammal and/or autologous ("self") substances and, as a result, treats autologous ("self") tissues and substances as if they were foreign and mounts an immune response against them. Autoimmune disease is characterized by production of either antibodies that react with self tissue, and/or the activation of immune effector T cells that are autoreactive to endogenous self antigens. Three main immunopathologic mechanisms act to mediate autoimmune diseases: 1) autoantibodies are directed against functional cellular receptors or other cell surface molecules, and either stimulate or inhibit specialized cellular function with or without destruction of cells or tissues; 2) autoantigen--autoantibody immune complexes form in intercellular fluids or in the general circulation and ultimately mediate tissue damage; and 3) lymphocytes produce tissue lesions by release of cytokines or by attracting other destructive inflammatory cell types to the lesions. These inflammatory cells in turn lead to production of lipid mediators and cytokines with associated inflammatory disease.

Since many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of CLASP-2 polypeptides or polynucleotides that can inhibit an immune response, particularly the proliferation, or differentiation of T-cells, can be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected by CLASP-2 include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid

syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitis, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, can also be treated by CLASP-2 polypeptides or polynucleotides. Moreover, CLASP-2 can be used to treat anaphylaxis or hypersensitivity to an antigenic molecules.

In one embodiment CLASP-2 polynucleotides or polypeptides are used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of CLASP-2 polypeptides or polynucleotides that inhibits an immune response, particularly the proliferation, differentiation of T-cells, can be an effective therapy in preventing organ rejection or GVHD.

Similarly, in another embodiment, CLASP-2 polypeptides or polynucleotides are used to modulate inflammation. The term "inflammation" refers to both acute responses (*i.e.*, responses in which the inflammatory processes are active) and chronic responses (*i.e.*, responses marked by slow progression and formation of new connective tissue). Acute and chronic inflammation can be distinguished by the cell types involved. Acute inflammation often involves polymorphonuclear neutrophils; whereas chronic inflammation is normally characterized by a lymphohistiocytic and/or granulomatous response. Inflammation includes reactions of both the specific and non-specific defense systems. A specific defense system reaction is a specific immune system reaction response to an antigen (possibly including an autoantigen). A non-specific defense system reaction is an inflammatory response mediated by leukocytes incapable of immunological memory. Such cells include granulocytes, macrophages, neutrophils and eosinophils.

For example, CLASP-2 polypeptides or polynucleotides can inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (*e.g.*, septic shock, sepsis, or systemic

inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1). Examples of specific types of inflammation are
5 diffuse inflammation, focal inflammation, croupous inflammation, interstitial inflammation, obliterative inflammation, parenchymatous inflammation, reactive inflammation, specific inflammation, toxic inflammation and traumatic inflammation.

In another embodiment CLASP-2 polypeptides or polynucleotides are used to treat or detect infectious agents. For example, by increasing the immune response,
10 particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases can be treated. The immune response can be increased by either enhancing an existing immune response, or by initiating a new immune response. CLASP-2 polypeptides or polynucleotides can be used to treat or detect any of these symptoms or diseases.

5.4.2.2. Use of CLASP-2 Polynucleotides in Screening

15 The presence or absence of hCLASP-2 nucleotide and amino acid sequences in a biological sample can be used in screening assays as medical diagnostics to aid in clinical decision-making. In one embodiment, hCLASP-2-based diagnostics involves screening assays for vaginal bleeding of unknown cause. In several examples discussed below, the cause of the bleeding can be in part differentiated by knowledge of whether the vaginal
20 bleeding contains placental components (Hart FD, Ed., 1985, French's Index of Differential Diagnosis, 12th Ed. John Wright & Sons, pp. 561-63). In these cases, the high expression of hCLASP-2 nucleotide sequences in placenta relative to its low expression in blood (FIG. 4A) will allow the detection of the presence of placenta based on the presence of the hCLASP-2 nucleotide or protein. Such detection can be achieved by quantitative RT-PCR, Northern
25 analysis, Western analysis, ELISAs, and fluorescence activated cell sorting (FACS) by using labeled anti-hCLASP-2 antibodies (Sambrook *et al.*, 1989, Molecular Cloning, 2nd Ed., Cold Spring Harbor Lab. Press; Harlow *et. al.*, 1988, Antibodies, a laboratory manual, Cold Spring Harbor Lab. Press).

For example, hCLASP-2 can be used in the following screening assays:

30 (1) A woman gives birth and presents with post-partum bleeding. In this case the presence of placental tissue indicates a condition called "retained products of

conception" that requires surgical evacuation of the uterus (Decherney and Pernol, Eds., 1996, Current Obstetric & Gynecologic Diagnosis & Treatment, 8th Ed. McGraw Hill).

(2) A pregnant woman suffers from vaginal bleeding of unknown origin. In this case the presence of placental tissue indicates a condition called "threatened abortion" that implies a poor prognosis for carrying the fetus to term (Decherney and Pernol, Eds., 1996, Current Obstetric & Gynecologic Diagnosis & Treatment, 8th Ed. McGraw Hill).

(3) A woman of child bearing age presents with vaginal bleeding and is found to have a positive pregnancy test without evidence of an intra-uterine pregnancy. In this case, the most serious of the differential diagnoses is ectopic pregnancy, a medical emergency. However, another common diagnosis is a completed abortion or miscarriage. The presence of products of conception (*i.e.* placenta) in the vaginal bleeding strongly favors the diagnosis of completed abortion over that of ectopic pregnancy (Decherney and Pernol, Eds., 1996, Current Obstetric & Gynecologic Diagnosis & Treatment, 8th Ed. McGraw Hill).

In another embodiment, hCLASP-2-based diagnostics involve screening assays to determine injury to vital tissues that express hCLASP-2 at high levels. Such tissues include kidney, heart, and lung (Fig 4A). Injury to these tissues can result in leakage of cells and cellular constituents including hCLASP-2 into surrounding fluids (specified below). Detection of abnormally high levels of hCLASP-2 protein in these surrounding fluids by Western analysis or ELISA, or detection of abnormally high levels of hCLASP-2 RNA in these fluids by RT-PCR or Northern analysis is expected to aid in the diagnosis of tissue injury.

In the case of renal injury, the hCLASP-2 nucleotide or amino acid sequences or fragments thereof would be expected to appear in the urine. Detection of abnormally high levels of hCLASP-2 can aid in the diagnosis of both nephritis and tubular necrosis, and differentiate from non-renal causes of proteinuria. Early diagnosis of nephritis is of particular value in patients with clinical signs and symptoms suggestive of systemic lupus erythematosus in whom early diagnosis and treatment of lupus nephritis can prevent irreversible kidney damage (Cameron J.S., 1999, J Nephrol 12 Suppl 2: S29-41). While tubular necrosis currently cannot be reversed by pharmacotherapy, differentiation of tubular necrosis from pre-renal failure is critical in formulating a treatment plan for oligouric hospitalized patients (Bidani A. and Churchill P.C., 1989, Dis Mon 35: 57-132).

In the case of myocardial injury, the hCLASP-2 nucleic or amino acid sequence or fragments thereof are expected to appear in the blood. This is analogous to current standard practice of monitoring for other elevated levels myocardial proteins (*e.g.*,

creatine kinase, troponin) in the blood following myocardial infarction and ischemia by standard ELISA or electrophoretic methodologies (Fauci *et al.*, (eds.), 1998, Harrison's Principles of Internal Medicine, 14th Ed., McGraw Hill, pp. 1352-1375). The presence of hCLASP-2 in cardiac muscle and its absence in skeletal muscle and blood makes hCLASP-2 an ideal marker to diagnose and monitor myocardial injury.

Unlike myocardial injury, pulmonary injury is not routinely diagnosed by assaying serum for lung-specific proteins. By analogy to myocardial infarction, pulmonary infarction also releases lung-specific proteins and cells into systemic circulation. Pulmonary infarction due to pulmonary embolism (PE) or pneumonia is expected to release hCLASP-2-bearing cells or protein/peptides into systemic circulation. Detection of hCLASP-2 protein in serum or RNA in blood can aid in the diagnosis of pulmonary infarction in the appropriate clinical setting. Current methods to diagnose PE are not only expensive but lack specificity and sensitivity, and the misdiagnosis of this condition is a leading cause of preventable death in hospitalized patients (Raskob G.E. and Hull R.D., 1999, *Curr Opin Hematol.* 6(5): 280-4).

In another embodiment, hCLASP-2-based diagnostics involve screening assays for identifying disorders of cells of hematopoietic lineage. hCLASP-2 is expressed in human T cells, B cells but not cells from the myeloid lineage. Different hCLASP-2 isoforms in T and B cells permit further discrimination between malignancies of T and B lineage (FIG. 4B). Precise identification of hematopoietic cell types is vital to guide chemotherapy and radiation therapy of patients with leukemia and lymphoma (Fauci *et al* Eds., 1998, Harrison's Principles of Internal Medicine, 14th Ed. McGraw Hill, pp. 695-712). hCLASP-2 expression differences can be detected by using FACS, immunofluorescence, immunoperoxidase staining, RT-PCR, in situ hybridization or RNA blot analysis (Sambrook, Fritsch and Maniatis, *Molecular Cloning*, 2nd Ed. Cold Spring Harbor Lab. Press, 1989; Ward MS, *Pathology* 1999 Nov; 31(4): 382-92).

In another embodiment, hCLASP-2-based diagnostics involve screening assays for identifying activated immune system cells. Although hCLASP-2 is generally expressed at quite low levels in PBMCs (which is critical for some of the above applications), it is known that the surface expression of the closely related mouse CLASP-1 protein is altered during the process of lymphocyte activation. An analogous change in expression is expected for the hCLASP-2 protein. Subtyping lymphocytes specific for a particular antigen, for example, using MHC-based multimeric staining reagents (Altman *et. al.*, 1996, *Science* 274: 94-6), for separating cell populations into hCLASP-2 high and hCLASP-2 low populations, can aid in determining the nature of the immune response against that antigen.

Such understanding is critical, for example, in predicting the course of chronic viral infections such as hepatitis B, hepatitis C, and HIV, and to designing appropriate treatment regimens for patients suffering from these infections.

hCLASP-2 can also serve as a potential therapeutic agent for Wilms' tumor.

5 Wilms tumor is the most common primary renal tumor of childhood (Cotran, Kumar, and Collins, 1999, Robbins Pathologic Basis of Disease, 6th Ed. W.B. Saunders, pp. 487-89). As discussed herein, hCLASP-2 is highly expressed in 293 cells, embryonic kidney epithelial cells. Therefore, hCLASP-2 nucleic or amino acid sequence or fragments can serve as tumor markers for Wilms' tumor. Antibodies directed against a hCLASP-2 variant that is expressed
10 only in Wilms' tumor can serve as novel therapeutic agents for Wilms' tumor, and can also function as delivery vehicles for other targeted therapeutics that may be attached to the anti-hCLASP-2 antibody (*e.g.*, chemotherapeutics or radiolabeling).

5.4.2.2.1. CLASP-2 Antisense, Ribozyme and Triplex Polynucleotides and Methods of Use

15 Oligonucleotide sequences, that include anti-sense RNA and DNA molecules and ribozymes that function to inhibit the translation of a CLASP-2 mRNA are within the scope of the invention. Such molecules are useful in cases where downregulation of CLASP-2 expression is desired. Anti-sense RNA and DNA molecules act to directly block the translation of mRNA by binding to targeted mRNA and preventing protein translation. The
20 invention provides methods and antisense oligonucleotide or polynucleotide reagents which can be used to reduce expression of CLASP-2 gene products in vitro or in vivo. Administration of the antisense reagents of the invention to a target cell results in reduced CLASP activity. As will be apparent to one of skill and as discussed *supra* (Table 3), specific CLASP-2 splice variants can be specifically targeted for inhibition. Alternatively, by
25 designing an, *e.g.*, antisense molecule that recognizes a sequence found in several or all CLASP-2 species, a general inhibition can be achieved.

A. Antisense

Without intending to be limited to any particular mechanism, it is believed that antisense oligonucleotides bind to, and interfere with the translation of, the sense CLASP-2
30 mRNA. Alternatively, the antisense molecule can render the CLASP-2 mRNA susceptible to nuclease digestion, interfere with transcription, interfere with processing, localization or otherwise with RNA precursors ("pre-mRNA"), repress transcription of mRNA from the

CLASP-2 gene, or act through some other mechanism. However, the particular mechanism by which the antisense molecule reduces CLASP-2 expression is not critical.

The antisense polynucleotides of the invention comprise an antisense sequence of at least 7 to 10 to typically 20 or more nucleotides that specifically hybridize to a sequence from mRNA encoding CLASP-2 or mRNA transcribed from the CLASP-2 gene. More often, the antisense polynucleotide of the invention is from about 10 to about 50 nucleotides in length or from about 14 to about 35 nucleotides in length. In other embodiments, antisense polynucleotides are polynucleotides of less than about 100 nucleotides or less than about 200 nucleotides. In general, the antisense polynucleotide should be long enough to form a stable duplex but short enough, depending on the mode of delivery, to administer in vivo, if desired. The minimum length of a polynucleotide required for specific hybridization to a target sequence depends on several factors, such as G/C content, positioning of mismatched bases (if any), degree of uniqueness of the sequence as compared to the population of target polynucleotides, and chemical nature of the polynucleotide (*e.g.*, methylphosphonate backbone, peptide nucleic acid, phosphorothioate), among other factors. Generally, to assure specific hybridization, the antisense sequence is substantially complementary to the target CLASP-2 mRNA sequence. In certain embodiments, the antisense sequence is exactly complementary to the target sequence. The antisense polynucleotides can also include, however, nucleotide substitutions, additions, deletions, transitions, transpositions, or modifications, or other nucleic acid sequences or non-nucleic acid moieties so long as specific binding to the relevant target sequence corresponding to CLASP-2 RNA or its gene is retained as a functional property of the polynucleotide.

It will be appreciated that the CLASP-2 polynucleotides and oligonucleotides of the invention can be made using nonstandard bases (*e.g.*, other than adenine, cytidine, guanine, thymine, and uridine) or nonstandard backbone structures to provides desirable properties (*e.g.*, increased nuclease-resistance, tighter-binding, stability or a desired TM). Techniques for rendering oligonucleotides nuclease-resistant include those described in PCT publication WO 94/12633. A wide variety of useful modified oligonucleotides may be produced, including oligonucleotides having a peptide-nucleic acid (PNA) backbone (Nielsen *et al.*, 1991, Science 254: 1497) or incorporating 2'-O-methyl ribonucleotides, phosphorothioate nucleotides, methyl phosphonate nucleotides, phosphotriester nucleotides, phosphorothioate nucleotides, phosphoramidates. Still other useful oligonucleotides may contain alkyl and halogen-substituted sugar moieties comprising one of the following at the 2' position: OH, SH, SCH₃, F, OCN, OCH₃OCH₃, OCH₃O(CH₂)_nCH₃, O(CH₂)_nNH₂ or

O(CH₂)_nCH₃, where n is from 1 to about 10; C1 to C10 lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl; Br; CN; CF₃; OCF₃; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃; SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a cholesteryl group; a folate group; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. Folate, cholesterol or other groups that facilitate oligonucleotide uptake, such as lipid analogs, may be conjugated directly or via a linker at the 2' position of any nucleoside or at the 3' or 5' position of the 3'-terminal or 5'-terminal nucleoside, respectively. One or more such conjugates may be used. Oligonucleotides may also have sugar mimetics such as cyclobutyls in place of the pentofuranosyl group. Other embodiments may include at least one modified base form or "universal base" such as inosine, or inclusion of other nonstandard bases such as queosine and wybutosine as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases. The antisense oligonucleotide can comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)_w, and 2,6-diaminopurine.

The invention further provides oligonucleotides having backbone analogues such as phosphodiester, phosphorothioate, phosphorodithioate, methylphosphonate, phosphoramidate, alkyl phosphotriester, sulfamate, 3'-thioacetal, methylene(methylimino), 3'-N-carbamate, morpholino carbamate, chiral-methyl phosphonates, nucleotides with short chain alkyl or cycloalkyl intersugar linkages, short chain heteroatomic or heterocyclic

intersugar ("backbone") linkages, or CH₂-NH-O-CH₂, CH₂-N(CH₃)-OCH₂, CH₂-O-N(CH₃)-CH₂, CH₂-N(CH₃)-N(CH₃)-CH₂ and O-N(CH₃)-CH₂-CH₂ backbones (where phosphodiester is O-P-O-CH₂), or mixtures of the same. Also useful are oligonucleotides having morpholino backbone structures (U.S. Patent No. 5,034,506).

- 5 Useful references include Oligonucleotides and Analogues, A Practical Approach, edited by F. Eckstein, IRL Press at Oxford University Press (1991); Antisense Strategies, Annals of the New York Academy of Sciences, Volume 600, Eds. Baserga and Denhardt (NYAS 1992); Milligan *et al.*, 9 July 1993, J. Med. Chem. 36(14): 1923-1937; Antisense Research and Applications (1993, CRC Press), in its entirety and specifically
- 10 Chapter 15, by Sanghvi, entitled "Heterocyclic base modifications in nucleic acids and their applications in antisense oligonucleotides;" and Antisense Therapeutics, ed. Sudhir Agrawal (Humana Press, Totowa, New Jersey, 1996).

- In one embodiment, the antisense sequence is complementary to relatively accessible sequences of the CLASP-2 mRNA (*e.g.*, relatively devoid of secondary structure).
- 15 This can be determined by analyzing predicted RNA secondary structures using, for example, the MFOLD program (Genetics Computer Group, Madison WI) and testing in vitro or in vivo as is known in the art. Another useful method for identifying effective antisense compositions uses combinatorial arrays of oligonucleotides (see, *e.g.*, Milner *et al.*, 1997, Nature Biotechnology 15: 537). Examples of oligonucleotides that can be tested in cells for antisense
- 20 suppression of CLASP-2 function are those capable of hybridizing to (*i.e.*, substantially complementary to) the following positions from SEQUENCE ID NO: 1:

- | | | |
|----|----|---------------------------------|
| | 1) | GAAGGCGATCATCACGTGGCCTTCCATCGC |
| | 2) | GCTTCAAGTAATGACTGGTGCAGAACATCTG |
| | 3) | GCTCCTCCTCAGGCAGGCGCTATGGCTGTGG |
| 25 | 4) | GTAGGCCCGGTGCAGCGTGTCATACAGATGG |

(See also Example 8)

- In some embodiments, administration of antisense oligonucleotides will result in reduction of hCLASP-mRNA expression by at least about 50%, as assessed by Northern analysis after administration of an antisense phosphorothioate oligonucleotide at a
- 30 concentration of 1 μ M, 5 μ M, 10 μ M or 20 μ M.

The invention also provides an antisense polynucleotide that has sequences in addition to the antisense sequence (*i.e.*, in addition to anti-CLASP-2-sense sequence). In this

case, the antisense sequence is contained within a polynucleotide of longer sequence. In another embodiment, the sequence of the polynucleotide consists essentially of, or is, the antisense sequence.

The antisense nucleic acids (DNA, RNA, modified, analogues, and the like) can be made using any suitable method for producing a nucleic acid, such as the chemical synthesis and recombinant methods disclosed herein. In one embodiment, for example, antisense RNA molecules of the invention can be prepared by *de novo* chemical synthesis or by cloning. For example, an antisense RNA that hybridizes to CLASP-2 mRNA can be made by inserting (ligating) an CLASP-2 DNA sequence (e.g., SEQUENCE ID No: 1, or fragment thereof) in reverse orientation operably linked to a promoter in a vector (e.g., plasmid). Provided that the promoter and, preferably termination and polyadenylation signals, are properly positioned, the strand of the inserted sequence corresponding to the noncoding strand will be transcribed and act as an antisense oligonucleotide of the invention. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter or enhancer) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

In one embodiment, antisense DNA oligodeoxyribonucleotides derived from the translation initiation site, e.g., between -10 and +10 regions of a CLASP-2 nucleotide sequence, are used. For general methods relating to antisense polynucleotides, see ANTISENSE RNA AND DNA, 1988, D.A. Melton, Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY). See also, Dagle *et al.*, 1991, Nucleic Acids Research, 19: 1805. For a review of antisense therapy, see, e.g., Uhlmann *et al.*, 1990, Chem. Reviews, 90: 543-584.

B. Ribozyme

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyze endonucleolytic cleavage of CLASP-2 RNA sequences.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences, GUA, GUU and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing

the cleavage site can be evaluated for predicted structural features such as secondary structure that can render the oligonucleotide sequence unsuitable. The suitability of candidate targets can also be evaluated by testing their accessibility to hybridization with complementary oligonucleotides, using ribonuclease protection assays.

5 C. Triplex

Alternatively, endogenous target gene expression can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the target gene (i.e., the target gene promoter and/or enhancers) to form triple helical structures that prevent transcription of the target gene in target cells in the body. (See generally, Helene, 1991, 10 *Anticancer Drug Des.*, 6(6): 569-584; Helene *et al.*, 1992, *Ann. N.Y. Acad. Sci.*, 660: 27-36; and Maher, 1992, *Bioassays* 14(12): 807-815).

Nucleic acid molecules to be used in triplex helix formation for the inhibition of transcription should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via 15 Hoogsteen base pairing rules, which generally require sizable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences can be pyrimidine-based, which will result in TAT and CGC⁺ triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel 20 orientation to that strand. In addition, nucleic acid molecules can be chosen that are purine-rich, for example, contain a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

25 Alternatively, the potential sequences that can be targeted for triple helix formation can be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizable stretch of either purines or pyrimidines to be present on one strand of a duplex.

30 D. General

The anti-sense RNA and DNA molecules, ribozymes and triple helix molecules of the invention can be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides well known in the art such as for example solid phase

phosphoramidite chemical synthesis. Alternatively, RNA molecules can be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences can be incorporated into a wide variety of vectors which contain suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters.

- 5 Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

Various modifications to the DNA molecules can be introduced as a means of increasing intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences of ribo- or deoxy- nucleotides to the 5' and/or
10 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

Methods for introducing polynucleotides into such cells or tissue include methods for *in vitro* introduction of polynucleotides such as the insertion of naked polynucleotide, *i.e.*, by injection into tissue, the introduction of a CLASP-2 polynucleotide in
15 a cell *ex vivo*, the use of a vector such as a virus, (*e.g.*, a retrovirus, adenovirus, adeno-associated virus, and the like), phage or plasmid, and the like or techniques such as electroporation or calcium phosphate precipitation.

5.4.2.2.2. Gene Therapy

By introducing gene sequences into cells, gene therapy can be used to treat
20 conditions in which the cells do not express normal CLASP-2 or express abnormal/inactive CLASP-2. In some instances, the polynucleotide encoding a CLASP-2 is intended to replace or act in the place of a functionally deficient endogenous gene. Alternatively, abnormal conditions characterized by overexpression can be treated using the gene therapy techniques described below.

25 In a specific embodiment, nucleic acids comprising a sequence encoding a CLASP-2 protein or functional derivative thereof, are administered to promote CLASP-2 function, by way of gene therapy. Gene therapy refers to therapy performed by the administration of a nucleic acid to a subject. In this embodiment of the invention, the nucleic acid produces its encoded protein that mediates a therapeutic effect by promoting CLASP-2
30 function.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, *see*, Goldspiel *et al.*, 1993, Clinical Pharmacy 12: 488-505; Wu and Wu, 1991, Biotherapy 3: 87-95; Tolstoshev, 1993, Ann. Rev. Pharmacol. Toxicol. 32: 573-596; Mulligan, 1993, Science 260: 926-932; and Morgan and Anderson, 1993, Ann. Rev. Biochem. 62: 191-217; Can, 1993, TIBTECH 11(5): 155-215). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel *et al.*, *supra*; and Kriegler, 1990, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY.

In one aspect, the therapeutic composition comprises a CLASP-2 nucleic acid that is part of an expression vector that encodes a CLASP-2 protein or fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the CLASP-2 coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the CLASP-2 coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the CLASP-2 nucleic acid (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. U.S.A. 86: 8932-8935; Zijlstra *et al.*, 1989, Nature 342: 435-438).

Delivery of the nucleic acid into a patient can be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

In a specific embodiment, the nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, *e.g.*, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by infection using a defective or attenuated retroviral or other viral vector (*see*, U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (*see, e.g.*, Wu and Wu, 1987, J. Biol. Chem. 262: 4429-4432) (which can be used to target cell types specifically expressing the receptors), and the like. In another embodiment, a nucleic acid-ligand

complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (*see, e.g.*, PCT Publications WO 92/06180 dated April 16, 5 1992; WO 92/22635 dated December 23, 1992; WO 92/20316 dated November 26, 1992; WO 93/14188 dated July 22, 1993; WO 93/20221 dated October 14, 1993). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. U.S.A. 86: 8932-8935; Zijlstra *et al.*, 1989, Nature 342: 435-438).

10 In a specific embodiment, a viral vector that contains the CLASP-2 nucleic acid is used. For example, a retroviral vector can be used (*see*, Miller *et al.*, 1993, Meth. Enzymol. 217: 581-599). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The CLASP-2 nucleic acid to be used in gene therapy is cloned into the vector, 15 which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen *et al.*, 1994, Biotherapy 6: 291-302, which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes *et al.*, 1994, J. Clin. Invest. 93: 644-651; Kiem *et al.*, 20 1994, Blood 83: 1467-1473; Salmons and Gunzberg, 1993, Human Gene Therapy 4: 129-141; and Grossman and Wilson, 1993, Curr. Opin. in Genetics and Devel. 3: 110-114.

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other 25 targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson 1993, Current Opinion in Genetics and Development 3: 499-503) present a review of adenovirus-based gene therapy. Bout *et al.*, 1994, Human Gene Therapy 5: 3-10, demonstrated the use of adenovirus vectors to transfer 30 genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld *et al.*, 1991, Science 252: 431-434; Rosenfeld *et al.*, 1992, Cell 68: 143-155; and Mastrangeli *et al.*, 1993, J. Clin. Invest. 91: 225-234. Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh *et al.*, 1993, Proc. Soc. Exp. Biol. Med. 204: 289-300).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, and the like. Numerous techniques are known in the art for the introduction of foreign genes into cells (*see, e.g.,* Loeffler and Behr, 1993, Meth. Enzymol. 217: 599-618; Cohen *et al.*, 1993, Meth. Enzymol. 217: 618-644; Cline, 1985, Pharmac. Ther. 29: 69-92) and can be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. In a preferred embodiment, epithelial cells are injected, *e.g.,* subcutaneously. In another embodiment, recombinant skin cells can be applied as a skin graft onto the patient. Recombinant blood cells (*e.g.,* hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, and the like., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.,* as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, and the like. In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

5 5.4.2.3. Knockout Cells

In one aspect of the invention, endogenous target gene expression can also be reduced by inactivating or "knocking out" the target gene or its promoter using targeted homologous recombination (*see, e.g.,* Smithies *et al.*, 1985, *Nature* 317: 230-234; Thomas and Capecchi, 1987, *Cell* 51: 503-512; Thompson *et al.*, 1989, *Cell* 5: 313-321; each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional target gene (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous target gene (either the coding regions or regulatory regions of the target gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express the target gene *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches are particularly suited in the agricultural field where modifications to ES (embryonic stem) cells can be used to generate animal offspring with an inactive target gene (*see, e.g.,* Thomas and Capecchi, 1987 and Thompson, 1989, *supra*). However, this approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors.

5.4.2.4. Transgenic and Knockout Animals

The CLASP-2 gene product can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, sheep, and non-human primates, *e.g.,* baboons, monkeys, and chimpanzees can be used to generate CLASP-2 transgenic animals. The term "transgenic," as used herein, refers to animals expressing CLASP-2 gene sequences from a different species (*e.g.,* mice expressing human CLASP-2 gene sequences), as well as animals that have been genetically engineered to overexpress endogenous (*i.e.,* same species) CLASP-2 sequences or animals that have been genetically engineered to no longer express endogenous CLASP-2 gene sequences (*i.e.,* "knock-out" animals), and their progeny.

Any technique known in the art can be used to introduce a CLASP-2 transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but

are not limited to pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Pat.-No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten *et al.*, 1985, Proc. Natl. Acad. Sci., U.S.A. 82: 6148-6152); gene targeting in embryonic stem cells (Thompson *et al.*, 1989, Cell 56: 313-321); electroporation of embryos (Lo, 1983, Mol. Cell. Biol. 3: 1803-1814); and sperm-mediated gene transfer (Lavitrano *et al.*, 1989, Cell 57: 717-723) (For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115, 171-229)

Any technique known in the art can be used to produce transgenic animal clones containing a CLASP-2 transgene, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal or adult cells induced to quiescence (Campbell *et al.*, 1996, Nature 380: 64-66; Wilmut *et al.*, Nature 385: 810-813).

The present invention provides for transgenic animals that carry a CLASP-2 transgene in all their cells, as well as animals that carry the transgene in some, but not all their cells, *i.e.*, mosaic animals. The transgene can be integrated as a single transgene or in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene can also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko *et al.* (1992, Proc. Natl. Acad. Sci. U.S.A. 89: 6232-6236). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the CLASP-2 transgene be integrated into the chromosomal site of the endogenous CLASP-2 gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous CLASP-2 gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous CLASP-2 gene. The transgene can also be selectively introduced into a particular cell type, thus inactivating the endogenous CLASP-2 gene in only that cell type, by following, for example, the teaching of Gu *et al.* (1994, Science 265: 103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant CLASP-2 gene can be assayed utilizing standard techniques. Initial screening can be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals can also be assessed using techniques

that include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR (reverse transcriptase PCR). Samples of CLASP-2 gene-expressing tissue, can also be evaluated immunocytochemically using antibodies specific for the CLASP-2 transgene product.

5 **5.4.2.5. Other Uses of CLASP-2 Polynucleotides**

There exists an ongoing need to identify new chromosome marking reagents. Sequences can be mapped to chromosomes by preparing PCR primers from SEQ ID NO: 1, 3, 5, or 9. These primers can be less than 50 nucleotides in length, generally less than 46 nucleotides, more generally less than 41 nucleotides, most generally less than 36
10 nucleotides, preferably less than 31 nucleotides, more preferably less than 26 nucleotides, and most preferably less than 21 nucleotides in length. The probes can also be less than 16 nucleotides, less than 13 nucleotides in length, less than 9 nucleotides in length and less than 7 nucleotides in length. Primers can be selected so that the primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of
15 somatic cell hybrids containing individual human chromosomes (*i.e.*, chromosome 13). Only those hybrids containing the human CLASP-2 gene corresponding to SEQ ID NO: 1, 3, 5, or 9 will yield an amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Precise chromosomal location of the CLASP-2
20 polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. See Verma, et al, Human Chromosomes: A Manual of Basic Techniques, Pergamon Press. NY, 1988. Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location
25 and presentation of a particular disease. See McKusick, V., 1998, Mendelian Inheritance in Man : A Catalog of Human Genes and Genetic Disorders, 12th Ed, Johns Hopkins University Press.

The CLASP-2 polynucleotides can be used for identifying individuals from minute biological samples as DNA markers for restriction fragment length polymorphism
30 (RFLP). An individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot with CLASP-2 DNA markers to yield unique bands for identifying the individual.

As described above, upon sequencing of numerous independent cDNA products, single nucleotide polymorphisms (SNPs) have been discovered within CLASP-2. These alterations and differences are presented in FIG 11B. They represent mis-sense alterations.

5 If it is determined that certain SNPs are deleterious or advantageous, SNPs can be used as a diagnostic tool through SNP mapping or direct sequencing of the SNP region to determine which isoform is expressed. Additionally, the SNPs can be used as a general SNP marker for chromosomal defects such as rearrangement and translocations.

CLASP-2 polynucleotides can be also be used as polymorphic markers for
10 forensic analysis. See generally National Research Council, *The Evaluation of Forensic DNA Evidence* (Eds. 1996, Pollard *et al.*, National Academy Press, Washington D.C.). The capacity to identify a distinguishing or unique set of forensic markers in an individual is useful for forensic analysis. For example, one can determine whether a blood sample from a suspect matches a blood or other tissue sample from a crime scene by determining whether
15 the set of polymorphic forms occupying selected polymorphic sites is the same in the suspect and the sample. If the set of polymorphic markers does not match between a suspect and a sample, it can be concluded (barring experimental error) that the suspect was not the source of the sample. If the set of markers does match, one can conclude that the DNA from the suspect is consistent with that found at the crime scene. If frequencies of the polymorphic
20 forms at the loci tested have been determined (*e.g.*, by analysis of a suitable population of individuals), one can perform a statistical analysis to determine the probability that a match of suspect and crime scene sample would occur by chance.

To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, *e.g.*, hair or skin, or body
25 fluids, *e.g.*, blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample. The CLASP-2 polynucleotide sequences of the present invention can be used to provide polynucleotide reagents, *e.g.*, PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for
30 example, providing another "identification marker" (*i.e.* another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to noncoding regions of SEQ ID NO: 1, 3, 5 or 9 are particularly appropriate for this use as greater numbers of polymorphisms occur in the

noncoding regions, making it easier to differentiate individuals using this technique.

Examples of polynucleotide reagents include the CLASP-2 nucleotide sequences or portions thereof, *e.g.*, fragments derived from the noncoding regions of SEQ ID NO: 1, 3, 5, or 9.

having a length of at least 20 bases, preferably at least 25 bases, and more preferably at least

5 30 bases.

CLASP-2 polynucleotides can also be used as reagents for paternity testing.

The object of paternity testing is usually to determine whether a male is the father of a child.

In most cases, the mother of the child is known and thus, the mother's contribution to the child's genotype can be traced. Paternity testing investigates whether the part of the child's

10 genotype not attributable to the mother is consistent with that of the putative father. Paternity

testing can be performed by analyzing sets of polymorphisms in the putative father and the

child. Of course, the present invention can be expanded to the use of this procedure to

determine if one individual is related to another. Even more broadly, the present invention

can be employed to determine how related one individual is to another, for example, between

15 races or species.

Bacterial infections are a major cause of health-related problems. However, the emergence of drug resistant bacteria is compromising the therapeutic value of the present spectrum of antibiotics. All the currently used antibiotics are small organic molecules, with certain level of structural similarity. This provides an advantage for bacteria to develop drug

20 resistance, since they need to modify a limited number of genes in order to become resistant to a wide variety of antibiotics. The development of antibiotics with different chemical

structure and targets can overcome antibiotic resistance, and provide therapeutic superiority in preventing infection by bacterial pathogens. Additionally, most antibiotics are not naturally

occurring compounds and cause minor or sometimes serious side effects. For example,

25 antibiotics used to treat TB can cause hearing loss.

The present invention provides new antibacterial agents. Certain CLASP-2 DNA sequences were difficult to clone and subclone (*see* Example 1). Bacteria harboring certain pieces of CLASP cDNA products were unable to be isolated, indicating that introduction of CLASP sequences compromised bacterial viability. There can be at least two

30 possible reasons why the CLASP cDNA were unable to be cloned, which can reflect a

variation of the well-established Modification and Restriction systems found in bacteria

(reviewed in Wilson and Murray. (1991) *Annu. Rev. Genet.* 25:585-627; Bickle and Kruger

(1993) *Microbiol. Rev.* 57:29-67). This well-described system is used by bacteria to prevent

deleterious effects caused by the introduction of foreign DNA. Bacteria can recognize

foreign DNA since it does not have the same modifications (e.g. methylation) as the native DNA. After recognition, the bacteria then digest and eliminate the foreign DNA (restriction). In the first scenario, the CLASP cDNA can be recognized as foreign DNA, and digested and eliminated as in the Modification and Restriction system. However, this would be unique for CLASP cDNA since the bacteria used for cloning cDNA are compromised in the Modification and Restriction system, which makes cloning of cDNA into bacteria a practice common in the art. If this is the case, the bacterial apparatus that specifically recognizes or eliminates CLASP cDNA can provide a novel target to develop antimicrobial agents. The CLASP DNA sequence would be useful in targeting the apparatus as well as an entry point for designing screens to identify potential targets. The second possibility is that CLASP cDNA behaves as an antimicrobial agent (i.e., antibiotic), and prevents bacterial growth. This, in effect, would create a new type of antibiotic mediated by the presence of foreign DNA (i.e. CLASP cDNA). In the case for the CLASP cDNA, the bacteria can recognize the DNA but instead of digesting and eliminating the DNA, the CLASP cDNA can cause a variation of the restriction and prevent the bacteria from growing, imposing a bacteriicidal effect upon the bacteria.

DNA as an antimicrobial agent has significant advantages over currently available agents. First, it is structurally unrelated to any existing antibiotics, and can overcome the present growing drug-resistance problem to structurally common agents. Second, since DNA antimicrobials composed of naturally-occurring human DNA, are expected to have minimal side effects and immune rejection. Third, DNA sequences can be tailored with sequence variation and numerous chemical modifications to circumvent the problem of resistance. Fourth, the antimicrobial DNA can be delivered specifically to bacterial cells through the use of bacteriophages (i.e., bacterial virus) which specifically infect bacteria and do not infect human cells. Further specificity can be generated to infect certain bacteria and bacterial subpopulations. Finally, this system can be economically robust since the generation of DNA and delivery vehicles are inexpensive.

5.5. Polypeptides Encoded by the CLASP-2 Gene Coding Sequence

In accordance with the invention, a CLASP-2 polynucleotide which encodes the CLASP-2 polypeptides, mutant polypeptides, peptide fragments, CLASP-2 fusion proteins or functional equivalents thereof, can be used to express CLASP-2 proteins in appropriate host cells. In various embodiments, the CLASP-2 polypeptides expressed will be identical or substantially similar to SEQ ID NOs: 2, 4, 6 or 10 or a fragment thereof.

In some embodiments, altered DNA sequences which can be used in accordance with the invention include deletions, additions or substitutions of different nucleotide residues resulting in a sequence that encodes the same or a functionally equivalent gene product. For example, due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence, can be used in the practice of the invention for the expression of the CLASP-2 protein. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. One of skill will recognize that each codon in a nucleic acid sequence such as SEQ ID NO: 1 (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence. Thus, for example, due to the degeneracy of the genetic code, a polypeptide having the sequence of SEQ ID NO: 2 or a fragment thereof, can be encoded by numerous polynucleotides other than SEQ ID NO: 1. Typically, the degenerate sequence will hybridize with SEQ ID NO: 1 under high or moderate stringency conditions, but this is not strictly required (e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions.)

The gene product itself can contain deletions, additions or substitutions of amino acid residues within a CLASP-2 sequence, which result in a silent change thus producing a functionally equivalent CLASP-2 protein. Such conservative amino acid substitutions can be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine, histidine and arginine; amino acids with uncharged polar head groups having similar hydrophilicity values include the following: glycine, asparagine, glutamine, serine, threonine, tyrosine; and amino acids with nonpolar head groups include alanine, valine, isoleucine, leucine, phenylalanine, proline, methionine, tryptophan. Creighton, 1984, PROTEINS, has grouped amino acids that are conservative substitutions for

one another as follows: (1) Alanine (A), Glycine (G); (2) Aspartic acid (D), Glutamic acid (E); (3) Asparagine (N), Glutamine (Q); (4) Arginine (R), Lysine (K); (5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); (6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); (7) Serine (S), Threonine (T); and (8) Cysteine (C), Methionine (M).

5 The DNA sequences of the invention can be engineered in order to alter a CLASP-2 coding sequence for a variety of ends, including but not limited to, alterations which modify processing and expression of the gene product. For example, mutations can be introduced using techniques which are well known in the art, *e.g.*, site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, phosphorylation, and the like.

10 Based on the domain organization of the CLASP-2 proteins, a large number of CLASP-2 mutant polypeptides can be constructed by modifying or rearranging the nucleotide sequences that encode the CLASP-2 extracellular, transmembrane and cytoplasmic domains.

 In various embodiments, the present invention provides homologues of the CLASP-2 polypeptides which function as either an CLASP-2 agonists or an CLASP-2
15 antagonist. In a preferred embodiment, the CLASP-2 agonists and antagonists stimulate or inhibit, respectively, a subset of the biological activities of the naturally occurring form of the CLASP-2 polypeptide. Thus, specific biological effects can be elicited by treatment with a homologue of limited function. In one embodiment, treatment of a subject with a homologue having a subset of the biological activities of the naturally occurring form of the polypeptide
20 has fewer side effects in a subject relative to treatment with the naturally occurring form of the CLASP-2 polypeptide.

 The invention contemplates both full-length CLASP-2 polypeptides and fragments, *e.g.*, fragments having a length of at least about 10, often 20, frequently 50 or 100 residues substantially identical to the exemplified CLASP-2 polypeptide sequences of the
25 invention. Protein fragments can be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, 161-180, 181-200, or 201 to the end of the coding
30 region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 200 amino acids in length. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes.

Preferred polypeptide fragments include the CLASP-2 protein. Further preferred polypeptide fragments include the CLASP-2 protein having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-X, can be deleted from the amino terminus of either the CLASP-2 polypeptide. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotide fragments encoding these CLASP-2 polypeptide fragments are also preferred.

Even if deletion of one or more amino acids from the N-terminus of a protein results in modification or loss of one or more biological functions of the protein, other biological activities can still be retained. Thus, the ability of shortened CLASP-2 muteins to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptides generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a CLASP-2 mutein with a large number of deleted N-terminal amino acid residues can retain some biological or immunogenic activities. In fact, peptides composed of as few as four CLASP-2 amino acid residues can often evoke an immune response.

Homologues of the CLASP-2 polypeptide can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of the CLASP-2 polypeptide. As used herein, the term "homologue" refers to a variant form of the CLASP-2 polypeptide which acts as an agonist or antagonist of the activity of the CLASP-2 polypeptide. An agonist of the CLASP-2 polypeptide can retain substantially the same, or a subset, of the biological activities of the CLASP-2 polypeptide. An antagonist of the CLASP-2 polypeptide can inhibit one or more of the activities of the naturally occurring form of the CLASP-2 polypeptide, by, for example, competitively binding to a downstream or upstream member of the CLASP-2 molecular pathway which includes the CLASP-2 polypeptide.

Modulation can be assayed by determining any parameter that is indirectly or directly affected by the expression of the target gene. Such parameters include, *e.g.*, changes in RNA or protein levels, changes in protein activity, changes in product levels, changes in downstream gene expression, changes in reporter gene transcription (luciferase, CAT, β -galactosidase, β -glucuronidase, GFP (*see, e.g.*, Mistili & Spector, 1997, *Nature Biotechnology* 15: 961-964); changes in signal transduction, phosphorylation and

dephosphorylation, receptor-ligand interactions, second messenger concentrations (*e.g.*, cGMP, cAMP, IP₃, and Ca²⁺), and cell growth. These assays can be *in vitro*, *in vivo*, and *ex vivo*. Such functional effects can be measured by any means known to those skilled in the art, *e.g.*, measurement of RNA or protein levels, measurement of RNA stability, identification of downstream or reporter gene expression, *e.g.*, via chemiluminescence, fluorescence, colorimetric reactions, antibody binding, inducible markers, ligand binding assays; changes in intracellular second messengers such as cGMP and inositol triphosphate (IP₃); changes in intracellular calcium levels; cytokine release, and the like.

5.5.1. Synthesis or Expression of CLASP-2 Polypeptide Expression Systems

In order to express a biologically active CLASP-2, the nucleotide sequence coding for CLASP-2, or a functional equivalent, is inserted into an appropriate expression vector. The CLASP-2 gene product as well as host cells or cell lines transfected or transformed with recombinant CLASP-2 expression vectors can be used for a variety of purposes. These include, but are not limited to, generating antibodies (*i.e.*, monoclonal or polyclonal) that competitively inhibit activity of CLASP-2 protein and neutralize its activity; antibodies that activate CLASP-2 function and antibodies that detect its presence on the cell surface or in solution. Anti-CLASP-2 antibodies can be used in detecting and quantifying expression of CLASP-2 levels in cells and tissues such as lymphocytes and macrophages, as well as isolating CLASP-2-positive cells from a cell mixture.

Methods which are well known to those skilled in the art can be used to construct recombinant expression vectors containing the CLASP-2 coding sequence and appropriate transcriptional/translational control signals. These methods include *in vitro* recombinant DNA techniques, synthetic techniques and *in vivo* recombination/genetic recombination. (*See, e.g.*, the techniques described in Sambrook *et al.*, 1989, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y. and Ausubel *et al.*, *supra*). The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of polypeptide desired, and the like. The expression vectors of the invention can be introduced into host cells to thereby produce polypeptides or

peptides, including fusion polypeptides or peptides, encoded by nucleic acids as described herein (e.g., CLASP-2 polypeptides, mutant forms of CLASP-2, fusion polypeptides, and the like).

A variety of host-expression vector systems can be utilized to express a CLASP-2 coding sequence. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA, or cosmid DNA expression vectors containing the CLASP-2 coding sequence; yeast transformed with recombinant yeast expression vectors containing the CLASP-2 coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the CLASP-2 coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing the CLASP-2 coding sequence; or animal cell systems. The expression elements of these systems vary in their strength and specificities. Depending on the host/vector system utilized, any of a number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used in the expression vector. For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage λ , plac, ptrp, ptac (ptrp-lac hybrid promoter; cytomegalovirus promoter) and the like can be used; when cloning in insect cell systems, promoters such as the baculovirus polyhedron promoter can be used; when cloning in plant cell systems, promoters derived from the genome of plant cells (e.g., heat shock promoters; the promoter for the small subunit of RUBISCO; the promoter for the chlorophyll α/β binding protein) or from plant viruses (e.g., the ³⁵S RNA promoter of CaMV; the coat protein promoter of TMV) can be used; when cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter) can be used; when generating cell lines that contain multiple copies of the CLASP-2 DNA, SV40-, BPV- and EBV-based vectors can be used with an appropriate selectable marker.

In bacterial systems a number of expression vectors can be advantageously selected depending upon the use intended for the expressed CLASP-2 product. For example, when large quantities of CLASP-2 protein are to be produced for the generation of antibodies or to screen peptide libraries, vectors which direct the expression of high levels of fusion protein products that are readily purified can be desirable. Such vectors include, but are not

limited to, the *E. coli* expression vector pUR278 (Ruther *et al.*, 1983, EMBO J. 2: 1791), in which the CLASP-2 coding sequence can be ligated into the vector in frame with the lacZ coding region so that a hybrid protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic acids Res. 13: 3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264: 5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In yeast, a number of vectors containing constitutive or inducible promoters can be used. (Current Protocols in Molecular Biology, Vol. 2, 1988 (Suppl. 1999), Ed. Ausubel *et al.*, Greene Publish. Assoc. & Wiley Interscience, Ch. 13; Grant *et al.*, 1987, Expression and Secretion Vectors for Yeast, in Methods in Enzymology, Eds. Wu & Grossman, 1987, Acad. Press, N.Y., Vol. 153, pp. 516-544; Glover, 1986, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3; and Bitter, 1987, Heterologous Gene Expression in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684; and The Molecular Biology of the Yeast *Saccharomyces*, 1982, Eds. Strathern *et al.*, Cold Spring Harbor Press, Vols. I and II.)

In cases where plant expression vectors are used, the expression of the CLASP-2 coding sequence can be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson *et al.*, 1984, Nature 310: 511-514), or the coat protein promoter of TMV (Takamatsu *et al.*, 1987, EMBO J. 6: 307-311) can be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi *et al.*, 1984, EMBO J. 3: 1671-1680; Broglie *et al.*, 1984, Science 224: 838-843); or heat shock promoters, *e.g.*, soybean hsp17.5-E or hsp17.3-B (Gurley *et al.*, 1986, Mol. Cell. Biol. 6: 559-565) can be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, and the like. (Weissbach & Weissbach, 1988, Methods for Plant Molecular Biology, Academic Press, NY, Section VIII, pp. 421-463; and Grierson & Corey, 1988, Plant Molecular Biology, 2d Ed., Blackie, London, Ch. 7-9.)

An alternative expression system which could be used to express CLASP-2 is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera*

frugiperda cells. The CLASP-2 coding sequence can be cloned into non-essential regions (e.g., the polyhedron gene) of the virus and placed under control of an AcNPV promoter (e.g., the polyhedron promoter). Successful insertion of the *CLASP-2* coding sequence will result in inactivation of the polyhedron gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedron gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (see, e.g., Smith *et al.*, 1983, J. Viol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral based expression systems can be utilized. In cases where an adenovirus is used as an expression vector, the CLASP-2 coding sequence can be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene can then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing CLASP-2 in infected hosts. (See, e.g., Logan & Shenk, 1984, Proc. Natl. Acad. Sci. U.S.A. 81: 3655-3659). Alternatively, the vaccinia 7.5K promoter can be used. (See, e.g., Mackett *et al.*, 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 7415-7419; Mackett *et al.*, 1984, J. Virol. 49: 857-864; Panicali *et al.*, 1982, Proc. Natl. Acad. Sci. U.S.A. 79: 4927-4931). Regulatable expression vectors such as the tetracycline repressible vectors can also be used to express a coding sequence in a controlled fashion.

Specific initiation signals can also be required for efficient translation of inserted CLASP-2 coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where the entire CLASP-2 gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals can be needed. However, in cases where only a portion of the CLASP-2 coding sequence is inserted, exogenous translational control signals, including the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the CLASP-2 coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, and the like. (see Bittner *et al.*, 1987, Methods in Enzymol. 153: 516-544).

In addition, a host cell strain can be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in a specific fashion desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products can be important for the function of the protein. The presence of several consensus N-glycosylation sites in CLASP-2 extracellular domains support the possibility that proper modification can play a role in CLASP-2 function. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product can be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, WI38, and the like.

Host cells transformed with nucleotide sequences encoding CLASP-2 may be cultured under conditions suitable for the expression and recovery of the soluble protein from cell culture. The protein produced by a transformed cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode CLASP-2 may be designed to contain signal sequences which direct secretion of CLASP-2 through a prokaryotic or eukaryotic cell membrane. Other constructions may be used to join sequences encoding CLASP-2 to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin,

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express CLASP-2 proteins can be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with the CLASP-2 DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, and the like.), and a selectable marker. Following the introduction of foreign DNA, engineered cells can be allowed to grow for 1-2 days in an enriched medium, and then switched to a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

This method can advantageously be used to engineer cell lines which express the CLASP-2 protein(s) on the cell surface. Such engineered cell lines are particularly useful in screening for molecules or drugs that affect CLASP-2 function.

A number of selection systems can be used, including but not limited to, the
5 herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, Cell 11: 223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. U.S.A. 48: 2026), and adenine phosphoribosyltransferase (Lowy *et al.*, 1980, Cell 22: 817) genes which can be employed in tk⁻, hgp⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for *dhfr*, which confers resistance to
10 methotrexate (Wigler *et al.*, 1980, Natl. Acad. Sci. U.S.A. 77: 3567; O'Hare *et al.*, 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 1527); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. U.S.A. 78: 2072); *neo*, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin *et al.*, 1981, J. Mol. Biol. 150: 1); and *hygro*, which confers resistance to hygromycin (Santerre *et al.*, 1984, Gene 30: 147).
15 Additional selectable genes have been described, namely *trpB*, which allows cells to utilize indole in place of tryptophan; *hisD*, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, 1988, Proc. Natl. Acad. Sci. U.S.A. 85: 8047); *ODC* (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., 1987, In: Current Communications
20 in Molecular Biology, Cold Spring Harbor Laboratory ed.) and glutamine synthetase (Bebbington *et al.*, 1992, Biotech 10: 169).

In an alternate embodiment of the invention, the coding sequence of CLASP-2 could be synthesized in whole or in part, using chemical methods well known in the art. (See, *e.g.*, Caruthers *et al.*, 1980, Nuc. Acids Res. Symp. Ser. 7: 215-233; Crea and Horn, 180,
25 Nuc. Acids Res. 9(10): 2331; Matteucci and Caruthers, 1980, Tetrahedron Letter 21: 719; and Chow and Kempe, 1981, Nuc. Acids Res. 9(12): 2807-2817.) Alternatively, the protein itself could be produced using chemical methods to synthesize a CLASP-2 amino acid sequence in whole or in part. For example, peptides can be synthesized by solid phase techniques, cleaved from the resin, and purified by preparative high performance liquid chromatography.
30 (See Creighton, 1983, Proteins Structures And Molecular Principles, W.H. Freeman and Co., N.Y. pp. 50-60). The composition of the synthetic polypeptides can be confirmed by amino acid analysis or sequencing (*e.g.*, the Edman degradation procedure; see Creighton, 1983, Proteins, Structures and Molecular Principles, W.H. Freeman and Co., N.Y., pp. 34-49).

In some embodiments, the CLASP-2 polypeptide contains non-naturally occurring amino acids or amino acid analogs (*i.e.*, compounds that have the same basic chemical structure as a naturally occurring amino acid, *i.e.*, an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium).

5.5.2. Identification of Cells That Express CLASP-2

The recombinant host cells which contain the coding sequence and which express a CLASP-2 gene product or fragments thereof can be identified by at least four general approaches; (a) DNA-DNA or DNA-RNA hybridization; (b) the presence or absence of "marker" gene functions; (c) assessing the level of transcription as measured by the expression of CLASP-2 mRNA transcripts in the host cell; and (d) detection of the gene product as measured by immunoassay or by its biological activity. Prior to the identification of gene expression, the host cells can be first mutagenized in an effort to increase the level of expression of CLASP-2, especially in cell lines that produce low amounts of CLASP-2.

In the first approach, the presence of the CLASP-2 coding sequence inserted in the expression vector can be detected by DNA-DNA or DNA-RNA hybridization using probes comprising nucleotide sequences that are homologous to the CLASP-2 coding sequence, respectively, or portions or derivatives thereof.

In the second approach, the recombinant expression vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (*e.g.*, thymidine kinase activity, resistance to antibiotics, resistance to methotrexate, transformation phenotype, occlusion body formation in baculovirus, and the like). For example, if the CLASP-2 coding sequence is inserted within a marker gene sequence of the vector, recombinants containing the CLASP-2 coding sequence can be identified by the absence of the marker gene function. Alternatively, a marker gene can be placed in tandem with the CLASP-2 sequence under the control of the same or different promoter used to control the expression of the CLASP-2 coding sequence. Expression of the marker in response to induction or selection indicates expression of the CLASP-2 coding sequence.

In the third approach, transcriptional activity for the CLASP-2 coding region can be assessed by hybridization assays. For example, RNA can be isolated and analyzed by Northern blot using a probe homologous to the CLASP-2 coding sequence or particular portions thereof. Alternatively, total nucleic acids of the host cell can be extracted and

assayed for hybridization to such probes. Additionally, reverse transcription-polymerase chain reactions can be used to detect low levels of gene expression.

In the fourth approach, the expression of the CLASP-2 protein product can be assessed immunologically, for example by Western blots, immunoassays such as radioimmuno-precipitation, enzyme-linked immunoassays, fluorescent activated cell sorting ("FACS"), and the like. This can be achieved by using an anti-CLASP-2 antibody. Alternatively, CLASP-2 protein can be expressed as a fusion protein with green-fluorescent protein to facilitate its detection in cells (United States Patent Nos. 5,491,084; 5,804,387; 5,777,079).

Identification of cells or tissues expressing CLASP protein or mRNA, especially CLASP-2 isoforms, can be useful for determining normal and abnormal CLASP expression in a given cell or tissue. As discussed above, a number of CLASP-2 isoforms have been identified, *e.g.*, in Jurkat cells, peripheral blood, and brain. The identification of mRNA or protein expression in various cell types and tissues can allow for identification of isoforms improperly expressed in either a spatial or temporal manner. Expression of hCLASP-2D isoform in hematopoietic cells may cause problems due to the presence of the SH3 domain not seen in the Jurkat and peripheral blood isoforms.

Other molecules in the immune system may also interact with portions of hCLASP2D. However, the absence of the PBM domain in the hCLASP-2D isoform may be necessary for function in certain cell types or tissues. Similarly, expression of CLASP isoforms 2A, 2B, and 2C in brain may cause problems for different reasons: the PBM present in these isoforms may interfere with a particular function by binding any of the known PDZ domain protein involved in formation of the neurological synapse. Similarly, the lack of an SH3 domain may cause an inappropriate response due to interactions with only a subset of molecules required for CLASP-2 function in the brain.

5.5.3. Uses of CLASP-2 Engineered Host Cells

In one embodiment of the invention, the CLASP-2 protein and/or cell lines that express CLASP-2 can be used to screen for antibodies, peptides, small molecules, natural and synthetic compounds or other cell bound or soluble molecules that bind to the CLASP-2 protein resulting in stimulation or inhibition of CLASP-2 function. For example, anti-CLASP-2 antibodies can be used to inhibit or stimulate CLASP-2 function and to detect its presence. Alternatively, screening of peptide libraries with recombinantly expressed soluble CLASP-2 protein or cell lines expressing CLASP-2 protein can be useful for identification of

therapeutic molecules that function by inhibiting or stimulating the biological activity of CLASP-2. The uses of the CLASP-2 protein and engineered cell lines, described in the subsections below, can be employed equally well for homologous CLASP-2 genes in various species.

5 In a specific embodiment of the invention, cell lines may be engineered to express the extracellular or intracellular domain of CLASP fused to another molecule such as GST. In addition, CLASP, its extracellular domain or its intracellular domain may be fused to an immunoglobulin constant region (Hollenbaugh and Aruffo, 1992, Current Protocols in Immunology, Unit 10.19; Aruffo *et al.*, 1990, Cell 61: 1303) to produce a soluble molecule
10 with increased half life. The soluble protein or fusion protein can be used in binding assays, affinity chromatography, immunoprecipitation, Western blot, and the like. Synthetic compounds, natural products, and other sources of potentially biologically active materials can be screened in assays that are well known in the art.

Random peptide libraries consisting of all possible combinations of amino
15 acids attached to a solid phase support can be used to identify peptides that are able to bind to a specific domain of CLASP-2 (Lam, K.S. *et al.*, 1991, Nature 354: 82-84). The screening of peptide libraries can have therapeutic value in the discovery of pharmaceutical agents that stimulate or inhibit the biological activity of CLASP-2.

Identification of molecules that are able to bind to the CLASP-2 protein can be
20 accomplished by screening a peptide library with recombinant soluble CLASP-2 protein. Methods for expression and purification of CLASP-2 are described in Section 5.7, *supra*, and can be used to express recombinant full length CLASP-2 or fragments of CLASP-2 depending on the functional domains of interest. Such domains include CLASP-2 extracellular domain, transmembrane domain, CLASP-2 intracellular domain, ITAM
25 containing domain, tyrosine phosphorylation site containing domain, cysteine cluster containing domain, cadherin motif containing domain, and coil/coil domain.

To identify and isolate the peptide/solid phase support that interacts and forms a complex with CLASP-2, it is necessary to label or "tag" the CLASP-2 molecule. The CLASP-2 protein can be conjugated to enzymes such as alkaline phosphatase or horseradish
30 peroxidase or to other reagents such as fluorescent labels which can include fluorescein isothiocyanate (FITC), phycoerythrin (PE) or rhodamine. Conjugation of any given label to CLASP-2 can be performed using techniques that are well known in the art. Alternatively, CLASP-2 expression vectors can be engineered to express a chimeric CLASP-2 protein containing an epitope for which a commercially available antibody exist. The epitope-

specific antibody can be tagged with a detectable label using methods well known in the art including an enzyme, a fluorescent dye or colored or magnetic beads.

The "tagged" CLASP-2 conjugate is incubated with the random peptide library for 30 minutes to one hour at 22°C to allow complex formation between CLASP-2 and peptide species within the library. The library is then washed to remove any unbound protein. If CLASP-2 has been conjugated to alkaline phosphatase or horseradish peroxidase the whole library is poured into a petri dish containing substrates for either alkaline phosphatase or peroxidase, for example, 5-bromo-4-chloro-3-indoyl phosphate (BCIP) or 3,3',4,4"-diaminobenzidine (DAB), respectively. After incubating for several minutes, the peptide/solid phase- CLASP-2 complex changes color, and can be easily identified and isolated physically under a dissecting microscope with a micromanipulator. If a fluorescent tagged CLASP-2 molecule has been used, complexes can be isolated by fluorescence activated sorting. If a chimeric CLASP-2 protein-expressing a heterologous epitope has been used, detection of the peptide/CLASP-2 complex can be accomplished by using a labeled epitope-specific antibody. Once isolated, the identity of the peptide attached to the solid phase support can be determined by peptide sequencing.

In addition to using soluble CLASP-2 molecules, in another embodiment, it is possible to detect peptides that bind to cell-associated CLASP-2 using intact cells. The use of intact cells is preferred for use with cell surface molecules. Methods for generating cell lines expressing CLASP-2 are described in Section 5.8. The cells used in this technique can be either live or fixed cells. The cells can be incubated with the random peptide library and bind to certain peptides in the library to form a "rosette" between the target cells and the relevant solid phase support/peptide. The rosette can thereafter be isolated by differential centrifugation or removed physically under a dissecting microscope. Techniques for screening combinatorial libraries are known in the art (Gallop *et al.*, 1994, J. Med. Chem., 37: 1233; Gordon, 1994, J. Med. Chem., 37: 1385).

As an alternative to whole cell assays for membrane bound receptors or receptors that require the lipid domain of the cell membrane to be functional, CLASP-2 molecules can be reconstituted into liposomes where label or "tag" can be attached.

30 **5.5.4. CLASP-2 Fusion Proteins**

In another embodiment of the invention, a CLASP-2 or a modified CLASP-2 sequence can be ligated to a heterologous sequence to encode a fusion protein. For example, for screening of peptide libraries for molecules that bind CLASP-2, it can be useful to

produce a chimeric CLASP-2 protein expressing a heterologous epitope that is recognized by a commercially available antibody. A fusion protein can also be engineered to contain a cleavage site located between a CLASP-2 sequence and the heterologous protein sequence, so that the CLASP-2 can be cleaved away from the heterologous moiety. In one embodiment, fusion proteins of the invention can contain the CLASP-2 extracellular domain comprising at least about residues 1 through 816 or fragment thereof. In another embodiment, fusion proteins can contain the CLASP-2 intracellular domain comprising at least about residue 843 through the end of the CLASP-2 sequence or fragment thereof.

5.6. Cloning Alleles, Variants, and Species Homologs of CLASP-2

In order to clone the full length cDNA sequence from any species encoding a CLASP-2 cDNA, or to clone variant forms of the molecule, labeled DNA probes made from nucleic acid fragments corresponding to any partial cDNA disclosed herein can be used to screen a cDNA library derived from lymphoid cells or brain cells. More specifically, oligonucleotides corresponding to either the 5' or 3' terminus of the cDNA sequence can be used to obtain longer nucleotide sequences. Briefly, the library can be plated out to yield a maximum of 30,000 pfu for each 150 mm plate. Approximately 40 plates can be screened. The plates are incubated at 37°C until the plaques reach a diameter of 0.25 mm or are just beginning to make contact with one another (3-8 hours). Nylon filters are placed onto the soft top agarose and after 60 seconds, the filters are peeled off and floated on a DNA denaturing solution consisting of 0.4N sodium hydroxide. The filters are then immersed in neutralizing solution consisting of 1M Tris-HCl, pH 7.5, before being allowed to air dry. The filters are prehybridized in hybridization buffer such as casein buffer containing 10% dextran sulfate, 0.5M NaCl, 50mM Tris-HCl, pH 7.5, 0.1% sodium pyrophosphate, 1% casein, 1% SDS, and denatured salmon sperm DNA at 0.5 mg/ml for 6 hours at 60°C. The radiolabeled probe is then denatured by heating to 95°C for 2 minutes and then added to the prehybridization solution containing the filters. The filters are hybridized at 60°C for 16 hours. The filters are then washed in 1X wash mix (10X wash mix contains 3M NaCl, 0.6M Tris base, and 0.02M EDTA) twice for 5 minutes each at room temperature, then in 1X wash mix containing 1% SDS at 60°C for 30 minutes, and finally in 0.3X wash mix containing 0.1% SDS at 60°C for 30 minutes. The filters are then air dried and exposed to x-ray film for autoradiography. After developing, the film is aligned with the filters to select a positive plaque. If a single, isolated positive plaque cannot be obtained, the agar plug containing the

plaques will be removed and placed in lambda dilution buffer containing 0.1M NaCl, 0.01M magnesium sulfate, 0.035M Tris HCl, pH 7.5, 0.01% gelatin. The phage can then be replated and rescreened to obtain single, well isolated positive plaques. Positive plaques can be isolated and the cDNA clones sequenced using primers based on the known cDNA sequence.

5 This step can be repeated until a full length cDNA is obtained.

It can be necessary to screen multiple cDNA libraries from different tissues to obtain a full length cDNA. In the event that it is difficult to identify cDNA clones encoding the complete 5' terminal coding region, an often encountered situation in cDNA cloning, the RACE (Rapid Amplification of cDNA Ends) technique can be used. RACE is a proven PCR-based strategy for amplifying the 5' end of incomplete cDNAs. 5'-RACE-Ready RNA
10 synthesized from human tissues containing a unique anchor sequence is commercially available (Clontech). To obtain the 5' end of the cDNA, PCR is carried out on 5'-RACE-Ready cDNA using the provided anchor primer and the 3' primer. A secondary PCR reaction is then carried out using the anchored primer and a nested 3' primer according to the
15 manufacturer's instructions. Once obtained, the full length cDNA sequence can be translated into amino acid sequence and examined for certain landmarks such as a continuous open reading frame flanked by translation initiation and termination sites, a cadherin-like domain, an ITAM domain, a tyrosine phosphorylation site, a cysteine cluster, a transmembrane domain, and finally overall structural similarity to the CLASP-2 genes disclosed herein. *See,*
20 *Ponassi et al.*, 1999, *Mech. Dev.* 80: 207-212; *Isakov*, 1998, *Receptor Channels* 5: 243-253; *Borroto et al.*, 1997, *Biopolymers* 42: 75-88; *Dimitratos et al.*, 1997, *Mech. Dev.* 63: 127-130; *Apperson et al.*, 1996, *J. Neurosci.* 16: 6839-6852; *Ozawa et al.*, 1990, *Mech. Dev.* 33: 49-56, which discuss protein domains and are incorporated herein by reference.

5.7. Modulating Expression of Endogenous CLASP-2 Genes

25 Alternatively, the expression characteristics of an endogenous CLASP-2 gene within a cell population can be modified by inserting a heterologous DNA regulatory element into the genome of the cell line such that the inserted regulatory element is operatively linked with the endogenous CLASP-2 gene. For example, an endogenous CLASP-2 gene which is normally "transcriptionally silent", *i.e.*, an *CLASP-2* gene which is normally not expressed, or
30 is expressed only at very low levels in a cell population, can be activated by inserting a regulatory element which is capable of promoting the expression of a normally expressed gene product in the cells. Alternatively, a transcriptionally silent, endogenous CLASP-2 gene

can be activated by insertion of a promiscuous regulatory element that works across cell types.

A heterologous regulatory element can be inserted into a cell line population, such that it is operatively linked with an endogenous CLASP-2 gene, using techniques, such as targeted homologous recombination, which are well known to those of skill in the art, (see 5 e.g. in Chappel, U.S. Patent No. 5,272,071; PCT publication No. WO 91/06667, published Jan 16, 1991).

5.8. Anti-CLASP-2 Antibodies

Various procedures known in the art can be used for the production of 10 antibodies to epitopes of the natural and recombinantly produced CLASP-2 protein. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, human or humanized, IgG, IgM, IgA, IgD or IgE, a complementarity determining region, Fab fragments, F(ab')₂ and fragments produced by an Fab expression library as well as anti-idiotypic antibodies. Antibodies which compete for CLASP-2 binding are especially 15 preferred for diagnostics and therapeutics.

Monoclonal antibodies that bind CLASP-2 can be radioactively labeled allowing one to follow their location and distribution in the body after injection. Radioisotope tagged antibodies can be used as a non-invasive diagnostic tool for imaging *de novo* lymphoid tumors and metastases that express CLASP-2.

Immunotoxins can also be designed which target cytotoxic agents to specific 20 sites in the body. For example, high affinity CLASP-2 specific monoclonal antibodies can be covalently complexed to bacterial or plant toxins, such as diphtheria toxin or ricin. A general method of preparation of antibody/hybrid molecules can involve use of thiol-crosslinking reagents such as SPDP, which attack the primary amino groups on the antibody and by 25 disulfide exchange, attach the toxin to the antibody. The hybrid antibodies can be used to specifically eliminate CLASP-2 expressing lymphocytes.

For the production of antibodies, various host animals can be immunized by injection with the recombinant or naturally purified CLASP-2 protein, fusion protein or peptides, including but not limited to goats, rabbits, mice, rats, hamsters, and the like 30 Various adjuvants can be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and poten-

tially useful human adjuvants such as BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum*.

Monoclonal antibodies to CLASP-2 can be prepared by using any technique which provides for the production of antibody molecules by continuous cell lines in culture.

5 These include, but are not limited to, the hybridoma technique originally described by Kohler and Milstein, (*Nature*, 1975, 256: 495-497), the human B-cell hybridoma technique (Kosbor *et al.*, 1983, *Immunology Today*, 4: 72; Cote *et al.*, 1983, *Proc. Natl. Acad. Sci. U.S.A.*, 80: 2026-2030) and the EBV-hybridoma technique (Cole *et al.*, 1985, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). In addition, techniques developed for the
10 production of "chimeric antibodies" (Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci. U.S.A.*, 81: 6851-6855; Neuberger *et al.*, 1984, *Nature*, 312: 604-608; Takeda *et al.*, 1985, *Nature*, 314: 452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. Alternatively, techniques described for the production of single chain
15 antibodies (U.S. Patent 4,946,778) can be adapted to produce CLASP-2 -specific single chain antibodies. In some embodiments, phage display technology is used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (see, *e.g.*, McCafferty *et al.*, *Nature* 348: 552-554 (1990); Marks *et al.*, *Biotechnology* 10: 779-783 (1992)).

20 Hybridomas can be screened using enzyme-linked immunosorbent assays (ELISA) in order to detect cultures secreting antibodies specific for refolded recombinant CLASP-2. Cultures can also be screened by ELISA to identify those cultures secreting antibodies specific for mammalian-produced CLASP-2. Confirmation of antibody specificity can be obtained by western blot using the same antigens. Subsequent ELISA testing can use
25 recombinant CLASP-2 fragments to identify the specific portion of the CLASP-2 molecule with which a monoclonal antibody binds. Additional testing can be used to identify monoclonal antibodies with desired functional characteristics such as staining of histological sections, immunoprecipitation of CLASP-2, inhibition of CLASP-2 binding or stimulation of CLASP-2 to transmit an intracellular signal. Determination of the monoclonal antibody
30 isotype can be accomplished by ELISA, thus providing additional information concerning purification or function.

Some anti-CLASP-2 monoclonal antibodies of the present invention are humanized, human or chimeric, in order to reduce their potential antigenicity, without reducing their affinity for their target. Humanized antibodies have been described in the art.

See, e.g., Queen, *et al.*, 1989, Proc. Natl Acad. Sci. U.S.A. 86: 10029; U.S. Patent Nos. 5,563,762; 5,693,761; 5,585,089 and 5,530,101. The human antibody sequences used for humanization can be the sequences of naturally occurring human antibodies or can be consensus sequences of several human antibodies. See Kettleborough *et al.*, 1991, *Protein Engineering* 4: 773; Kolbinger *et al.*, 1993, *Protein Engineering* 6: 971. Humanized monoclonal antibodies against CLASP-2 peptides can also be produced using transgenic animals having elements of a human immune system (see, e.g., U.S. Patent Nos. 5,569,825; 5,545,806; 5,693,762; 5,693,761; and 5,7124,350).

In some embodiments, an anti-CLASP-2 polypeptide monoclonal or polyclonal antiserum is produced that is specifically immunoreactive with a particular CLASP-2 polypeptide and is selected to have low cross-reactivity against other molecules (e.g., other CLASP polypeptides) and any such cross-reactivity is removed by immunoabsorption prior to use in the immunoassay. Methods for screening and characterizing monoclonal antibodies for specificity are well known in the art and are described generally in Harlow and Lane, *supra*. For example, polyclonal antibodies raised to hCLASP-2A, as shown in SEQ ID NO: 1, or splice variants, or immunogenic portions thereof, can be selected to obtain only those polyclonal or monoclonal antibodies that are specifically immunoreactive with the target protein not with other proteins. This selection may be achieved by subtracting out antibodies that cross-react with molecules. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (see, e.g., Harlow & Lane, *Antibodies, A Laboratory Manual* (1988) for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background. Alternatively, antibodies that cross-react with a selected set of polypeptides may be prepared.

Antibody fragments which contain specific binding sites of V can be generated by known techniques. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries can be constructed (Huse *et al.*, 1989, *Science*, 246: 1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity to CLASP-2.

Anti-CLASP-2 antibodies can also be used to identify, isolate, inhibit or eliminate CLASP-2-expressing cells. In one embodiment, the present invention includes a method of identifying an abnormal T cell profile of an immunocompromised subject relative to the T cell profile of a non-immunocompromised subject. The method includes (i) sorting a sample of peripheral blood mononuclear cells (PBMC) isolated from the immunocompromised subject into sets of T cell types, (ii) determining the ratio of CLASP-2⁺ cells relative to the total number of cells (CLASP-2⁺: total) in each set, and identifying an abnormal T cell profile in the immunocompromised subject by comparing the CLASP-2⁺: total ratios of sets from the immunocompromised subject with the CLASP-2⁺: total ratios of analogous sets from a non-immunocompromised subject.

In other embodiments, anti-CLASP-2 antibodies can be used for detection of hCLASP-2 protein in assays such as fluorescent activated cell sorting (FACS), ELISA, fluorescent or electron immunomicroscopy, Western blots, gel shift analyses. CLASP-2 expression in various cells, localization within cells, interactions with other proteins, and differentiation between CLASP-2 isoform expression can be determined by use of the techniques listed herein.

5.9. Screening Assays

The invention provides methods for identifying compounds or agents that modulate (*i.e.*, inhibit or enhance) CLASP-2 expression or activity. CLASP-2 expression or activity modulators are useful for treatment of disorders characterized by (or associated with) aberrant or abnormal CLASP-2 expression or activity. Aberrant expression of CLASP-2 mRNA or protein means expression in lymphocytes (*e.g.*, T lymphocytes or B lymphocytes) or other CLASP-2 expressing cells of at least 2-fold, preferably at least 5-fold greater than expression in control lymphocytes obtained from a healthy subject.

The CLASP-2 expression assays can include the steps of contacting a cell expressing CLASP-2 with a compound or agent and assaying CLASP-2 expression. CLASP-2 polypeptide expression is easily measured by ELISA using anti-CLASP-2 antibodies of the invention. CLASP-2 mRNA expression (including expression of specific species or splice variants of CLASP-2) can be measured by quantitative Northern analysis or quantitative PCR.

CLASP-2 activities include, for example, the CLASP-2 polypeptide binding to PDZ-domain containing molecules and CLASP-2 polypeptide involvement in signal transduction (*e.g.*, leading to T cell activation). Compounds or agents that modulate the

interaction of a CLASP-2 polypeptide and a target molecule, modulate CLASP-2 nucleic acid expression, or modulate CLASP-2 polypeptide activity are all contemplated by the methods of the present invention.

Test compounds include, for example, 1) peptides (*e.g.*, soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, *e.g.*, Lam, K. S. *et al.*, 1991, Nature 354: 82-84; Houghten, R. *et al.*, 1991, Nature 354: 84-86) and combinatorial chemistry-derived molecular libraries made of D- and/or L-configuration amino acids; 2) phosphopeptides (*e.g.*, members of random and partially degenerate, directed phosphopeptide libraries, see, *e.g.*, Songyang, Z. *et al.*, 1993, Cell 72: 767-778); 3) CLASP-2 antibodies (as described above); 4) small organic and inorganic molecules (*e.g.*, molecules obtained from combinatorial and natural product libraries); 5) antisense RNA and DNA molecules and ribozymes (described above).

The CLASP modulators can be any of a large variety of compounds, both naturally occurring and synthetic, organic and inorganic, and including polymers (*e.g.*, oligopeptides, polypeptides, oligonucleotides, and polynucleotides), small molecules, antibodies, sugars, fatty acids, nucleotides and nucleotide analogs, analogs of naturally occurring structures (*e.g.*, peptide mimetics, nucleic acid analogs, and the like), and numerous other compounds.

In one embodiment, the invention provides assays for screening test compounds which bind to CLASP-2 polypeptides. The assays can be recombinant cell based or cell-free assays. These assays can include the steps of combining a cell expressing a CLASP-2 polypeptide or a binding fragment thereof, and a compound or agent under conditions which allow binding of the compound or agent to the CLASP-2 polypeptide to form a complex. Complex formation can then be determined. The ability of the candidate compound or agent to bind to the CLASP-2 polypeptide or fragment thereof is indicated by the presence of the candidate compound in the complex. Formation of complexes between the CLASP-2 polypeptide and the candidate compound can be quantitated, for example, using standard immunoassays.

In another embodiment, the invention provides screening assays to identify test compounds which modulate the interaction (and most likely CLASP-2 activity as well) between a CLASP-2 polypeptide and a molecule (target molecule with which the CLASP-2 polypeptide normally interacts).

In one embodiment, these CLASP-2 target molecules can be tyrosine kinases (*e.g.*, lyn, lck, fyn, ZAP-70m SyK, and CSK). In another embodiment, these CLASP-2 target

molecules can be tyrosine phosphatases (*e.g.*, EZRIN, SHP-1, SHP-2 and PTP36). In another embodiment, these CLASP-2 target molecules can be adaptor proteins (*e.g.*, NCK, CBL, SHC, LNK, SLP-76, HS1, SIT, VAV, GrB2, and BRDG1). In another embodiment, these CLASP-2 target molecules can be cytoskeletal associated proteins such as ankyrin, spectrin, talin, ezrin, tropomyosin, myosin, plectin, syndecans, paralemmin, Band 3 protein, cytoskeletal protein 4.1, and PTP36. In a further embodiment, CLASP-2 target molecules can be members of the integrin family.

Typically, the assays are recombinant cell based or cell-free assay. These assays can include the steps of combining a cell expressing a CLASP-2 polypeptide or a binding fragment thereof, a CLASP-2 target molecule (*e.g.*, a CLASP-2 ligand) and a test compound, under conditions where but for the presence of the candidate compound, the CLASP-2 polypeptide or biologically active portion thereof binds to the target molecule. Detecting complex formation between the CLASP-2 polypeptide or the binding fragment thereof, the CLASP-2 target molecule and a test compound detecting the formation of a complex which includes the CLASP-2 polypeptide and the target molecule can be accomplished. Detection of complex formation can include direct quantitation of the complex by, for example, measuring inductive effects, such as T cell activation, of the CLASP-2 polypeptide. A significant change, such as a decrease, in the interaction of the CLASP-2 and target molecule (*e.g.*, in the formation of a complex between the CLASP-2 and the target molecule) in the presence of a candidate compound (relative to what is detected in the absence of the candidate compound) is indicative of a modulation of the interaction between the CLASP-2 polypeptide and the target molecule. Modulation of the formation of complexes between the CLASP-2 polypeptide and the target molecule can be quantitated using, for example, an immunoassay. To perform cell free drug screening assays, it is desirable to immobilize either CLASP-2 or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the polypeptides, as well as to accommodate automation of the assay. CLASP-2 binding to a target molecule, in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and microcentrifuge tubes.

In one embodiment, a fusion polypeptide can be provided which adds a domain that allows the polypeptide to be bound to a matrix. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of CLASP-2-binding

polypeptide found in the bead fraction quantitated from the gel using standard electrophoretic techniques.

Other techniques for immobilizing polypeptides on matrices can also be used in the drug screening assays of the invention. For example, either CLASP-2 or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated CLASP-2 molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with CLASP-2 but which do not interfere with binding of the polypeptide to its target molecule can be derivatized to the wells of the plate, and CLASP-2 trapped in the wells by antibody conjugation. As described above, preparations of a CLASP-2 -binding polypeptide and a candidate compound are incubated in the CLASP-2 -presenting wells of the plate, and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes include immunodetection of complexes using antibodies reactive with the CLASP-2 target molecule, or which are reactive with CLASP-2 polypeptide and compete with the target molecule; as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing the CLASP-2, *e.g.*, the protein having the sequence of SEQ ID NO: 2. Such cells, either in viable or fixed form, can be used for standard ligand/receptor binding assays (see, *e.g.*, Parce *et al.* (1989) Science 246: 243-247; and Owicki *et al.* (1990) Proc. Natl Acad. Sci. U.S.A. 87: 4007-4011, which describe sensitive methods to detect cellular responses. A test compound, often labeled, can be assayed for binding or for competition with another ligand for binding. Viable cells could also be used to screen for the effects of drugs on CLASP-2 mediated functions, *e.g.*, T cell activation, second messenger levels, and others).

In another embodiment, the invention provides a method for identifying a compound (*e.g.*, a screening assay) capable of use in the treatment of a disorder characterized by (or associated with) aberrant or abnormal CLASP-2 nucleic acid expression or CLASP-2 polypeptide activity. This method typically includes the step of assaying the ability of the compound or agent to modulate the expression of the CLASP-2 nucleic acid or the activity of the CLASP-2 polypeptide thereby identifying a compound for treating a disorder characterized by aberrant or abnormal CLASP-2 nucleic acid expression or CLASP-2 polypeptide activity.

Methods for assaying the ability of the compound or agent to modulate the expression of the CLASP-2 nucleic acid or activity of the CLASP-2 polypeptide are typically cell-based assays. For example, cells which are sensitive to ligands which transduce signals via a pathway involving CLASP-2 can be induced to overexpress a CLASP-2 polypeptide in the presence and absence of a candidate compound. Candidate compounds which produce a change in CLASP-2-dependent responses can be identified. In one embodiment, expression of the CLASP-2 nucleic acid or activity of a CLASP-2 polypeptide is modulated in cells and the effects of candidate compounds on the readout of interest (such as T cell activation) are measured. For example, the expression of genes which are up- or down-regulated in response to a CLASP-2-dependent signal cascade can be assayed.

Alternatively, modulators of CLASP-2 expression can be identified in a method where a cell is contacted with a candidate compound and the expression of CLASP-2 mRNA or polypeptide in the cell is determined. The level of expression of CLASP-2 mRNA or polypeptide in the presence of the candidate compound is compared to the level of expression of CLASP-2 mRNA or polypeptide in the absence of the candidate compound. The candidate compound can then be identified as a modulator of CLASP-2 nucleic acid expression based on this comparison. For example, when expression of CLASP-2 mRNA or polypeptide is greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of CLASP-2 nucleic acid expression. Alternatively, when CLASP-2 nucleic acid expression is less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of CLASP-2 nucleic acid expression. The level of CLASP-2 nucleic acid expression in the cells can be determined by methods described herein for detecting CLASP-2 mRNA or polypeptide.

Modulators of CLASP-2 polypeptide activity and CLASP-2 nucleic acid expression identified according to these drug screening assays can be used to treat, for example, immune disorders. These methods of treatment include the steps of administering the modulators of CLASP-2 polypeptide activity or nucleic acid expression, *e.g.*, in a pharmaceutical composition as described in §5.10.1 below, to a subject in need of such treatment, *e.g.*, a subject with a disorder described herein.

5.10. Therapeutic Administration of CLASP-2 Modulators

The CLASP-2 protein is expressed in lymphocytes and, as noted *supra*, play a role in regulating T cell and B cell interactions, thus making CLASP-2 activity (*e.g.*, CLASP-

2 binding of regulatory proteins) a target for diagnostic and treatment of immune disorders
and for modulation of immune function (e.g., T cell activation). Additionally, since CLASP-2
contains domains capable of transducing an intracellular signal, cell surface CLASP-2 can be
triggered by an anti- CLASP-2 antibody or soluble CLASP-2 or a fragment thereof in order
5 to enhance the activation state of a lymphocyte.

5.10.1. Formulation and Route of Administration

A CLASP-2 polypeptide, a fragment thereof, anti-CLASP-2 antibody,
CLASP-2 polynucleotide (e.g., antisense or ribozyme), or small molecule agonists or
antagonists can be administered to a subject *per se* or in the form of a pharmaceutical or
10 therapeutic composition. Pharmaceutical compositions comprising the proteins of the
invention can be manufactured by means of conventional mixing, dissolving, granulating,
dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.
Pharmaceutical compositions can be formulated in conventional manner using one or more
physiologically acceptable carriers, diluents, excipients or auxiliaries which facilitate
15 processing of the protein or active peptides into preparations which can be used
pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

Currently, there are three major classes of protein-derived cell-penetrating
peptides that have been used for delivering of proteins into cells and animals (Lindgren, M.;
et al., 2000, Trends Pharmacol Sci. 21: 99-103). In one embodiment, the CLASP-2 protein
20 or fragment (encoding a functional domain of CLASP-2) can be introduced into the cell as a
fusion protein tied to a transporter protein derived from homeoprotein transcription factors
such as ANTP. In another embodiment, the CLASP-2 protein or fragment (encoding a
functional domain of CLASP-2) can be introduced into the cell as a fusion protein tied to
other transcription factors such as the HIV Tat protein and the herpes simplex virus type 1
25 (HSV-1) VP22 protein. Members in this family have been widely used in different cellular
and animal systems (Schwarze, S.; *et al.*; 2000, Trends Pharmacol Sci. 21: 45-48). In another
embodiment, the CLASP-2 protein or fragment (encoding a functional domain of CLASP-2)
can be introduced into the cell as a fusion protein tied to peptides derived from signal-
sequences present in several proteins such as HIV-1 gp41. In other embodiments, there are
30 several synthetic and/or chimeric cell-penetrating peptides such as transportan and
Amphiphilic model peptide (Lindgren, M.; *et al.*, 2000, Trends Pharmacol Sci. 21: 99-103)
that can be used. In another embodiment, the CLASP-2 protein or fragment can be

introduced by using anti-DNA antibodies (see, *e.g.*, Zack, D. J., *et al.*, 1996, J. Immunol. 157: 2082-8

For topical administration the proteins of the invention can be formulated as solutions, gels, ointments, creams, suspensions, and the like. as are well-known in the art.

5 Systemic formulations include those designed for administration by injection, *e.g.*, subcutaneous, intravenous, intramuscular, intrathecal or intraperitoneal injection, as well as those designed for transdermal, transmucosal, oral or pulmonary administration.

For injection, the proteins of the invention can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's
10 solution, or physiological saline buffer. The solution can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the proteins can be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

15 For oral administration, a composition can be readily formulated by combining the proteins with pharmaceutically acceptable carriers well known in the art. Such carriers enable the proteins to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. For oral solid formulations such as, for example, powders, capsules and tablets, suitable excipients
20 include fillers such as sugars, such as lactose, sucrose, mannitol and sorbitol; cellulose preparations such as maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP); granulating agents; and binding agents. If desired, disintegrating agents can be added, such as the cross-linked
25 polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

If desired, solid dosage forms can be sugar-coated or enteric-coated using standard techniques.

For oral liquid preparations such as, for example, suspensions, elixirs and solutions, suitable carriers, excipients or diluents include water, glycols, oils, alcohols, and
30 the like. Additionally, flavoring agents, preservatives, coloring agents and the like can be added.

For buccal administration, the proteins can take the form of tablets, lozenges, and the like. formulated in conventional manner.

For administration by inhalation, the proteins for use according to the present invention are conveniently delivered in the form of an aerosol spray from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The proteins can also be formulated in rectal or vaginal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the proteins can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the proteins can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Alternatively, other pharmaceutical delivery systems can be employed. Liposomes and emulsions are well known examples of delivery vehicles that can be used to deliver the proteins or peptides of the invention. Certain organic solvents such as dimethylsulfoxide also can be employed, although usually at the cost of greater toxicity. Additionally, the proteins can be delivered using a sustained-release system, such as semipermeable matrices of solid polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules can, depending on their chemical nature, release the proteins for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization can be employed.

As the proteins and peptides of the invention can contain charged side chains or termini, they can be included in any of the above-described formulations as the free acids or bases or as pharmaceutically acceptable salts. Pharmaceutically acceptable salts are those salts which substantially retain the biologic activity of the free bases and which are prepared by reaction with inorganic acids. Pharmaceutical salts tend to be more soluble in aqueous and other protic solvents than are the corresponding free base forms.

5.10.2. Effective Dosages

CLASP-2 polypeptides, CLASP-2 fragments and anti-CLASP-2 antibodies will generally be used in an amount effective to achieve the intended purpose. For use to inhibit an immune response, the proteins of the invention, or pharmaceutical compositions thereof, are administered or applied in a therapeutically effective amount. By therapeutically effective amount is meant an amount effective ameliorate or prevent the symptoms, or prolong the survival of, the patient being treated. Determination of a therapeutically effective amount is well within the capabilities of those skilled in the art, especially in light of the detailed disclosure provided herein.

For systemic administration, a therapeutically effective dose can be estimated initially from *in vitro* assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture (*i.e.*, the concentration of test compound that inhibits 50% of CLASP-2 binding interactions). Such information can be used to more accurately determine useful doses in humans.

Initial dosages can also be estimated from *in vivo* data, *e.g.*, animal models, using techniques that are well known in the art. One having ordinary skill in the art could readily optimize administration to humans based on animal data.

Dosage amount and interval can be adjusted individually to provide plasma levels of the proteins which are sufficient to maintain therapeutic effect. Usual patient dosages for administration by injection range from about 0.1 to 5 mg/kg/day, preferably from about 0.5 to 1 mg/kg/day. Therapeutically effective serum levels can be achieved by administering multiple doses each day.

In cases of local administration or selective uptake, the effective local concentration of the proteins can not be related to plasma concentration. One having skill in the art will be able to optimize therapeutically effective local dosages without undue experimentation.

The amount of CLASP-2 administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

The therapy can be repeated intermittently while symptoms detectable or even when they are not detectable. The therapy can be provided alone or in combination with other drugs. In the case of autoimmune disorders, the drugs that can be used in combination

with CLASP-2 or fragments thereof include, but are not limited to, steroid and non-steroid immunosuppressive agents.

5.10.3. Toxicity

Preferably, a therapeutically effective dose of the proteins described herein will provide therapeutic benefit without causing substantial toxicity.

Toxicity of the proteins described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, by determining the LD₅₀ (the dose lethal to 50% of the population) or the LD₁₀₀ (the dose lethal to 100% of the population). The dose ratio between toxic and therapeutic effect is the therapeutic index.

The data obtained from these cell culture assays and animal studies can be used in formulating a dosage range that is not toxic for use in human. The dosage of the proteins described herein lies preferably within a range of circulating concentrations that include the effective dose with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (*See, e.g., Fingl et al., 1975, In: The Pharmacological Basis of Therapeutics, Ch.1, p.1*).

5.11. Binding Assays

CLASP-2 polypeptides can be used to screen for molecules that bind to CLASP-2 or for molecules to which CLASP-2 binds. The binding of CLASP-2 by the molecule can activate (agonist), increase, inhibit (antagonist), or decrease activity of the CLASP-2 or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (*e.g.*, receptors), or small molecules. Preferably, the molecule is closely related to the natural ligand of CLASP-2, *e.g.*, a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (*See, Coligan et al., Current Protocols in Immunology 1(2): Chapter 5 (1991).*) Similarly, the molecule can be closely-related to the natural receptor to which CLASP-2 binds, or at least, a fragment of the receptor capable of being bound by CLASP-2 (*e.g.*, active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express CLASP-2, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing CLASP-2

(or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either CLASP-2 or the molecule.

The assay can simply test binding of a candidate compound to CLASP-2, where binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay can test whether the candidate compound results in a signal generated by binding to CLASP-2.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide affixed to a solid support, chemical libraries, or natural product mixtures. The assay can also simply comprise the steps of mixing a candidate compound with a solution containing CLASP-2, measuring CLASP-2 activity or binding, and comparing the CLASP-2 activity or binding to a standard. Preferably, an ELISA assay can measure CLASP-2 level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure CLASP-2 level or activity by either binding, directly or indirectly, to CLASP-2 or by competing with CLASP-2 for a substrate.

In another aspect of the invention, the CLASP-2 polypeptides, or fragments thereof, can be used as "bait proteins" in a two-hybrid assay (see, e.g., U.S. Pat. No. 5,283,317; Zervos *et al.*, 1993, Cell 72: 223-232; Madura *et al.*, 1993, J. Biol. Chem. 268: 12046-12054; Bartel *et al.*, 1993, Biotechniques 14: 920-924; Iwabuchi *et al.*, 1993, Oncogene 8: 1693-1696; and Brent WO 94/10300), to identify other proteins, which bind to or interact with CLASP-2 ("CLASP-2-binding proteins" or "CLASP-2-bp") and modulate CLASP-2 polypeptide activity. Such CLASP-2-binding proteins are also likely to be involved in the propagation of signals by the CLASP-2 polypeptides as, for example, upstream or downstream elements of the CLASP-2 pathway.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient by activating or inhibiting the CLASP-2 molecule. Moreover, the assays can discover agents which can inhibit or enhance the production of CLASP-2 from suitably manipulated cells or tissues.

Therefore, the invention includes a method of identifying compounds or agents that bind to CLASP-2 polypeptides comprising the steps of: (a) contacting a CLASP-2 polypeptide with a compound or agent under conditions which allow binding of the compound to the CLASP-2 polypeptide to form a complex and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists or

antagonists comprising the steps of: (a) incubating a candidate compound with CLASP-2, (b) assaying a biological activity, and (b) determining if a biological activity of CLASP-2 has been altered.

Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, *e.g.*, Fodor *et al.*, 1991, Science 251: 767-773, and other descriptions of chemical diversity libraries, which describe means for testing of binding affinity by a plurality of compounds.

5.12. Other Uses of CLASP-2 Polynucleotides and Polypeptides

The polynucleotides, polypeptides, polypeptide homologues, modulators, and antibodies described herein can be used in one or more of the following methods: a) drug screening assays; b) diagnostic assays particularly in disease identification, allelic screening and pharmacogenetic testing; and c) pharmacogenomics. A CLASP-2 polypeptide of the invention has one or more of the activities described herein and can thus be used to, for example, modulate an immune response in an immune cell, for example by binding to a CLASP-2 binding partner making it unavailable for binding to the naturally present CLASP-2 polypeptide.

In one embodiment, these CLASP-2 binding partners can be tyrosine kinases (*e.g.*, lyn, lck, fyn, ZAP-70m SyK, and CSK). In another embodiment, these CLASP-2 binding partners can be tyrosine phosphatases (*e.g.*, EZRIN, SHP-1, SHP-2 and PTP36). In another embodiment, these CLASP-2 target molecules can be adaptor proteins (*e.g.*, NCK, CBL, SHC, LNK, SLP-76, HS1, SIT, VAV, GrB2, and BRDG1. In another embodiment, these CLASP-2 binding partners can be cytoskeletal associated proteins such as ankyrin, spectrin, talin, ezrin, tropomyosin, myosin, plectin, syndecans, paralemmin, Band 3 protein, cytoskeletal protein 4.1, and PTP36. In a further embodiment, CLASP-2 binding partners can be members of the integrin family. The isolated nucleic acid molecules of the invention can be used to express CLASP-2 polypeptide (*e.g.*, via a recombinant expression vector in a host cell or in gene therapy applications), to detect CLASP-2 mRNA (*e.g.*, in a biological sample) or a naturally occurring or recombinantly generated genetic mutation in an CLASP-2 gene, and to modulate CLASP-2 activity, as described further below. In addition, the CLASP-2 polypeptides can be used to screen drugs or compounds which modulate CLASP-2 polypeptide activity as well as to treat disorders characterized by insufficient production of CLASP-2 polypeptide or production of CLASP-2 polypeptide forms which have decreased activity compared to wild type CLASP-2. Moreover, the anti-CLASP-2 antibodies of the

invention can be used to detect and isolate an CLASP-2 polypeptide, particularly fragments of CLASP-2 present in a biological sample, and to modulate CLASP-2 polypeptide activity.

5.13. Diagnostic Assays

The invention further provides a method for detecting the presence of CLASP-2, or fragment thereof, in a biological sample. Usually the biological sample contains lymphocytes (*e.g.*, from blood). The method involves contacting the biological sample with a compound or an agent capable of detecting CLASP-2 polypeptide or mRNA such that the presence of CLASP-2 is detected in the biological sample.

A preferred agent for detecting CLASP-2 mRNA is a directly or indirectly labeled nucleic acid probe capable of hybridizing to CLASP-2 mRNA. The nucleic acid probe can be, for example, the full-length CLASP-2 cDNA of SEQ ID NO: 1, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to CLASP-2 mRNA.

A preferred agent for detecting CLASP-2 polypeptide is a directly or indirectly labeled antibody capable of binding to a CLASP-2 polypeptide. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab)₂) can be used. The term "directly or indirectly", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The detection method of the invention can be used to detect CLASP-2 mRNA or polypeptide in a biological sample in vitro as well as in vivo. For example, in vitro techniques for detection of CLASP-2 mRNA include Northern hybridizations and in situ hybridizations. In vitro techniques for detection of CLASP-2 polypeptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. Alternatively, CLASP-2 polypeptide can be detected in vivo in a subject by introducing into the subject a labeled anti-CLASP-2 antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods which detect the allelic variant of CLASP-2 expressed in a subject and methods which detect fragments of an CLASP-2 polypeptide in a sample.

The invention also encompasses kits for detecting the presence of CLASP-2 in a biological sample. For example, the kit can comprise a directly or indirectly labeled compound or agent capable of detecting CLASP-2 polypeptide or mRNA in a biological sample; means for determining the amount of CLASP-2 in the sample; and means for
5 comparing the amount of CLASP-2 in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect CLASP-2 mRNA or polypeptide.

The methods of the invention can also be used to detect naturally occurring genetic mutations in an CLASP-2 gene, thereby determining if a subject with the mutated
10 gene is at risk for a disorder characterized by aberrant or abnormal CLASP-2 nucleic acid expression or CLASP-2 polypeptide activity as described herein. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic mutation characterized by at least one of an alteration affecting the integrity of a gene encoding an CLASP-2 polypeptide, or the misexpression of the CLASP-2 gene.

15 **5.14. Biological Activities of CLASP-2**

As described herein, CLASP-2 mediates a variety of cell functions in lymphocytes and other cells. As described herein, a variety of assays are useful for detecting or quantitating CLASP-2 activity, or for identifying agents (including polynucleotides, polypeptides, and antibodies of the invention) that modulate CLASP-2 activity (*i.e.*,
20 biological activity, *e.g.*, binding) or expression. Such agents are useful for treatment of diseases and conditions associated with aberrant CLASP-2 expression or activity. Further, following the guidance provided herein, other CLASP-2-mediated activities can be identified by those of skill using routine assays, such as those described below.

Exemplary assays for CLASP-2 function (or modulation of function) include
25 assays for modulation of an *in vitro* or *in vivo* cell response (*e.g.*, an immune response such as lymphocyte activation, antibody production, inflammation) by detecting a change in an activity (*e.g.*, cytokine production, calcium flux, tyrosine phosphorylation, regulation of early activation markers, cell metabolism, proliferation, and the like, as described below) of cells *in vitro* or *in vivo*. In one embodiment, the cells are lymphocytes.

30 In one assay, for example, recombinant CLASP-2 protein, peptides, or antibodies corresponding to the CLASP-2 extracellular domain can be mixed directly with T and B cells. Cytokine production by these cells can then be measured and the degree of modulation of the immune response quantitated. Alternatively, antigen-presenting B cells are

mixed with untransfected T cells or T cells that have been transfected with CLASP-2 isoforms. Cytokine production (or calcium flux or other assays in §5.14.3) is to be measured at the appropriate time to determine the effect of CLASP-2 on such an immune response. In a similar assay, B cells transfected with CLASP-2 constructs are tested for their ability to stimulate a T cell to generate an immune response. Transfected constructs in any of these cases could encode, for example, full or partial length CLASP-2 sequences, or antisense constructs to inhibit translation of endogenous CLASP-2 gene. Any of the examples described herein can be used to stimulate an immune response in the presence or absence of CLASP-2 isoforms or antibodies and assay the resulting effects on immune response by the methods listed in §5.14.3.

5.14.1 Methods for Generating an Immune Response in vitro

In various assays, an effect of an agent on immune cells is detected using an *in vitro* assay. The degree of an immune response can be measured or quantitated by a number of standard assays including those described below.

In one assay, human peripheral blood mononuclear cells (PBMC), human T cell clones (*e.g.*, Jurkat E6, ATCC TIB-152), EBV-transformed B cell clones (*e.g.*, 9D10, ATCC CRL-8752), antigen-specific T cell clones or lines can be used to examine immune responses *in vitro*. Activation, enhanced activation or inhibition of activation of these cells or cell lines can be used for the evaluation of potential CLASP therapeutics. Standard methods by which hematopoietic cells are stimulated to undergo activation characteristic of an immune response are, for example:

A) Antigen specific stimulation of immune responses. Either pre-immunized or naïve mouse splenocytes can be generated by standard procedures. In addition, antigen-specific T cell clones and hybridomas (*e.g.*, MBP-specific), and numerous B cell lymphoma cell lines (*e.g.*, CH27), have been previously characterized and are available for the assays discussed below. Antigen specific splenocytes or B-cells can be mixed with specific T-cells in the presence of antigen to generate an immune response. This can be performed in the presence or absence of CLASP-2 to assay whether CLASP-2 modulates the immune response as measured by any of the assays in section 5.14.2.

B) Non-specific T cell activation. The following methods can be used to activate T cells in the absence of antigen: 1) cross-linking T cell receptor (TCR) by addition of antibodies against receptor activation molecules (*e.g.*, TCR, CD3, or CD2) together with antibodies against co-stimulator molecules, for example anti-CD28; 2) activating cell surface

receptors in a non-specific fashion using lectins such as concanavalin A (con A) and phytohemagglutinin (PHA); 3) mimicking cell surface receptor-mediated activation using pharmacological agents that activate protein kinase C (*e.g.*, phorbol esters) and increase cytoplasmic Ca^{2+} (*e.g.*, ionomycin).

5 C) Non-specific B cell activation: 1) application of antibodies against cell surface molecules such as IgM, CD20, or CD21. 2) Lipopolysaccharide (LPS), phorbol esters, calcium ionophores and ionomycin can also be used to by-pass receptor triggering.

D) Mixed lymphocyte reaction (MLR). Mix donor PBMC with recipient PBMC to activate lymphocytes by presentation of mismatched tissue antigens, which occurs
10 in all cases except identical twins.

E) Generation of a specific T cell clone or line that recognizes a particular antigen. A standard approach is to generate tetanus toxin-specific T cells from a donor that has recently been boosted with tetanus toxin. Major histocompatibility complex- (MHC-) matched antigen presenting cells and a source of tetanus toxin are used to maintain antigen
15 specificity of the cell line or T cell clone (Lanzavecchia, A., *et al.*, 1983, *Eur. J. Immun.* 13: 733-738).

The anticipated mechanism of action of a CLASP-2 polypeptide or polynucleotide should define the appropriate assay to use to investigate its potential enhancement or inhibition of lymphocyte activation. For example, soluble proteins
20 containing the CLASP extracellular domain may interfere with the interaction between T cells and antigen presenting cells. Such interaction plays a role in the MLR and in antigen-specific T cell activation, but not in non-specific T or B cell activation. The assays described above have the advantage of several possible detection methods for quantitation.

5.14.2. Methods for Generating an Immune Response in vivo

25 In various assays, an effect of an agent on immune cells is detected using an *in vivo* assay. The degree of an immune response can be measured or quantitated by a number of standard assays including those described below.

(A) Animal Model for Transplantation Rejection: Ectopic Heart
Transplantation

30 In one embodiment, a standard animal model for graft versus host rejection is ectopic heart transplantation (Fulmer *et al.*, 1963, *Am. J. Anat.* 113: 273-281). This method involves using BALB/C mice (either sex, and range from 1-9 months) for transplanting cardiac tissue into a surgically-created pocket on the dorsum for both ears made by slitting

the skin over the auricular artery at the base of the ear. Small curved forceps are forced into the slit, bluntly dissecting between the skin and the cartilage plate. Donor tissue is eased into the base of the pocket near the distal edge of the ear. The auricular artery is used to seal off the opening of the pocket. Within 10 to 14 days pulsatile activity of the transplant should be observed. Gross appearance of the graft, patterns of vacuolar supply to the graft area and pulsatile activity can be easily observed utilizing transilluminated light during the first three weeks post-transplantation. Follow-up can continue for for several months.

(B) Animal model for Autoimmune Disease: Induction of Collagen Induced Arthritis (CIA)

Collagen Induced Arthritis (CIA) is a standard model for studying progression and immune (Courtenay *et al.*, 1980, Nature 283: 666 and Wooley *et al.*, 1981, J. Exp. Med. 154: 688). DBA/a mice can be used as an assay for the in vivo relevance of CLASP-2 in vitro testing potential immune therapeutics. In vivo experiments will be performed to examine the ability of potential therapeutics to prevent CIA. We will use 3-5 mice per group to statistically justify our results.

Once a titer of the potency of collagen type II (CII) is obtained therapeutics can be tested. In one embodiment, three mice will be immunized with three different concentrations of CII 50, 200, and 400 µg per animal (Nabozny *et al.*, 1996, J. Exp. Med., 183: 27-37). To induce CIA, animals can be immunized with an appropriate concentration of CII, determined as described above. One half of a 1:1 ratio of antigen:CFA can be injected at the base of the tail and the remainder equally divided in each hind footpad. Mice can be carefully monitored every day for the onset and progression of CIA throughout the experiment until its termination 12 weeks post-immunization with CII. The pieces of heart transplanted can be approximately 3 X 3 mm in size. The severity of arthritis can be assessed following standard procedures known to one of skill in the art.

5.14.3 Assay Quantitation

(A) Tyrosine phosphorylation

Tyrosine phosphorylation of early response proteins such as HS1, PLC-r, ZAP-76, and Vav is an early biochemical event following T cell activation. The tyrosine phosphorylated proteins can be detected by Western blot using antibodies against phosphorylated tyrosine residues. Tyrosine phosphorylation of these early response proteins can be used as a standard assay for T cell activation (J. Biol. Chem., 1997, 272(23): 14562-14570). Any change in the phosphorylation pattern of these or related proteins when immune

responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

(B) Intracellular Calcium Flux

The kinetics of intracellular Ca^{2+} concentrations are measured over time after stimulation of cells preloaded with a calcium sensitive dye. Upon binding the Ca^{2+} indicator dye, Fluor-4 (Molecular Probes), exhibits an increase in fluorescence level using flow cytometry, solution fluorometry, and confocal microscopy. Any change in the level or timing of calcium flux when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response

(C) Regulation of early activation markers

Increased and diminished expression/regulation of early lymphocyte activation marker levels such as CD69, IL-2R, MHC class II, B7, and TCR are commonly measured with fluorescently labeled antibodies using flow cytometry. All antibodies are commercially available. Any change in the expression levels of lymphocyte activation markers when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

(D) Increased metabolic activity/acid release

Activation of most known signal transduction pathways trigger increases in acidic metabolites. This reproducible biological event is measured as the rate of acid release using a microphysiometer (Molecular Devices), can be used as an early activation marker when comparing the treatment of cells with potential biological therapeutics (McConnell, H.M. *et al.*, 1992, Science 257: 1906-1912 and McConnell, H.M., 1995, Proc. Natl. Acad. Sci. 92: 2750-2754). Any statistically significant increase or decrease in acid release of CLASP-2-treated sample, as compared to control sample (no treatment), suggest and effect of CLASP-2 on biological function.

(E) Cell proliferation/cell viability assays

(1) ^3H -thimidine incorporation

Exposure of lymphocytes to antigen or mitogen in vitro induces DNA synthesis and cellular proliferation. The measurement of mitotic activity by ^3H -thimidine incorporation into newly synthesized DNA is one of the most frequently used assays to quantitative T cell activation. Depending on the cell population and form of stimulation used to activate the T cells, mitotic activity can be measured within 24-72 hrs. in vitro, post ^3H -thimidine pulse (Mishell, B. B. and S. M. Shiigi, 1980, Selected Methods in Cellular Immunology, W. H. Freeman and Company and Dutton, R. W. and Pearce, J. D., 1962,

Nature 194: 93). Any statistically significant increase or decrease in CPM of CLASP-2-treated sample, as compared to control sample (no treatment), suggest an effect of CLASP-2 on biological function.

(2) MTS [5-(3-carboxymethoxyphenyl)-2-(4,5-dimethylthiazolyl)-3(4-sulfophenyl)tetrazolium, inner salt] is a colorimetric method for determining the number of viable cells in proliferation or cytotoxicity assays (Barltrop, J.A. *et al.*, 1991, Bioorg. & Med. Chem. Lett. 1: 611). 1-5 days after lymphocyte activation, MTS tetrazolium compound, Owen's reagent, is bio-reduced by cells into a colored formazan product that is soluble in tissue culture media. Color intensity is read at 490 nm minus 650 nm using a microplate reader. Any statistically significant increase or decrease in color intensity of CLASP-2-treated sample, as compared to control sample (no treatment), can suggest an effect of CLASP-2 on biological function (Mosmann, T., 1983, J. Immunol. Methods 65: 55 and Barltrop, J.A. *et al.* (1991)).

(3) Bromodeoxyuridine (BrdU), a thymidine analogue, readily incorporates into cells undergoing DNA synthesis. BrdU-pulsed cells are labeled with an enzyme-conjugated anti-BrdU antibody (Gratzner, H.G., 1982, Science 218: 474-475.). A colorimetric, soluble substrate is used to visualize proliferating cells that have incorporated BrdU. Reaction is stopped with sulfuric acid and plate is read at 450 nm using a microplate reader. Any statistically significant increase or decrease in color intensity of CLASP-2-treated sample, as compared to control sample (no treatment), suggest an effect of CLASP-2 on biological function.

(F) Apoptosis by Annexin V

Programmed cell death or apoptosis is an early event in a cascade of catabolic reactions leading to cell death. A loss in the integrity of the cell membrane allows for the binding of fluorescently conjugated phosphatidylserine. Stained cells can be measured by fluorescence microscopy and flow cytometry (Vermes, I., 1995, J. Immunol. Methods. 180: 39-52). In one embodiment, any statistically significant increase or decrease in apoptotic cell number of CLASP-2-treated sample, as compared to control sample (no treatment), suggest an effect of CLASP-2 on biological function. For evaluating apoptosis in situ, assays for evaluating cell death in tissue samples can also be used in vivo studies.

(G) Quantitation of cytokine production

Cell supernatants harvested after cell stimulation for 16-48 hrs are stored at -80°C until assayed or directly tested for cytokine production. Multiple cytokine assays can be performed on each sample. IL-2, IL-3, IFN- γ and other cytokine ELISA Assays are

available for mouse, rat, and human (Endogen, Inc. and BioSource). Cytokine production is measured using a standard two-antibody sandwich ELISA protocol as described by the manufacturer. The presence of horseradish peroxidase is detected with 3, 3', 5' tertamethyl benziidine (TMB) substrate and the reaction is stopped with sulfuric acid. The absorbency at 450 nm is measured using a microplate reader. Any statistically significant increase or decrease in color intensity of CLASP-2-treated sample, as compared to control sample (no treatment), suggest an effect of CLASP-2 on biological function.

(H) NF-AT can be visualized by Immunostaining

T cell activation requires the import of nuclear factor of activated T cells (NFAT) to the nucleus. This translocation of NF-AT can be visualized by immunostaining with anti-NF-AT antibody (Cell 1998, 93: 851-861). Therefore, NF-AT nuclear translocation has been used to assay T cell activation. Similarly, NF-AT/luciferase reporter assays have been used as a standard measurement of T cell activation (MCB 1996, 12: 7151-7160).

(I) ELISA for collagen type II (CII)-specific antibodies (see above for related in vivo assay)

C(II) titers from serum of animals immunized with CLASP-2 can be measured and compared. Both TH1-dependent IgG2a and TH2-dependent IgG1 and IgE CII-specific antibody isotypes will be measured by ELISA. Mouse blood will be obtained by orbital bleed one and two months post-immunization with CII. Samples will be allowed to coagulate and centrifuge to obtain sera, and stored at -80°C until assayed by ELISA. Coat ELISA plates with CII and dilute sera. HRP conjugated goat, isotype specific antibody. Plates are then expose to TMB substrate and read at 450 nm using a microplate reader (Nabozny *et al.*, 1996, J. Exp. Med. 183: 27-37). Any change in the levels of Collagen specific antibodies by colorimetric test when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

(J) Antibody Production by ELISPOT Assay

A solid-phase enzyme-linked immunospot (ELISPOT) assay for the quantification of isotype-specific antibody secreting cells (Czerkinsky *et al.*, 1983, J Immunol. Methods. 65: 109-121). Both human and mouse B cells can be tested for isotype and antigen specific antibody production. Although based on a standard ELISA, this technique becomes more sensitive by detecting antibody secretion from single cells. Any change in ELISPOT levels when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

(K) Cellular degranulation following IgE cross-linking.

Two cell lines have been obtained from ATCC (MEG01 and HEL-17.92), both of which express the human FCεR1 receptor. FCεR1 is the high affinity receptor for IgE complexes, which when coupled to biotin can be cross-linked with avidin to induce degranulation and histamine release of lymphocytes. Following acylation of the sample, histamine is quantified with an enzyme immunoassay competition assay (Immunotech). Histamine release. A statistically significant increase or decrease in histamine concentration of a CLASP-2 treated sample, as compared to control sample (no treatment), suggest an effect of CLASP-2 on biological function. Any change in frequency of degranulation or histamine levels when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

(L) Cellular phenotyping of lymphocytes by flow cytometry and Immunocytochemistry

Determining the tissue distribution of lymphocytes following a pathological disorder can aid in identifying specific organ, tissue and lymphocyte involved in an immune response. Cellular phenotyping of lymphocyte trafficking is generally performed with by flow cytometry and Immunocytochemistry. There are several cluster determination (CD) molecules that are routinely used to identify phenotype, activation kinetics, and regulation events of cells. Any change in levels or distribution of CD molecules when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

(M) Structure/Function Assays: Homotypic and/or Heterotypic, Calcium-dependant Cell Adhesion

L929 cells can be transfected with CLASP-2 and Neomycin. G418-resistant clones can be screened for CLASP-expression with anti-CLASP peptide-specific antibodies. These CLASP-expressing clones can then be used to test for homotypic and/or heterotypic calcium dependent cell adhesion using the "cell aggregation assay" described for cadherin molecules (Murphy-Erdosh, C. *et al.*, 1995, J. Cell Biol. 129: 1379-1390). Any change in the levels of cellular aggregation when immune responses are generated in the presence of CLASP-2 is indicative of a CLASP-2 modulation of this response.

The following cDNA clones described in the Specification and further described in the Examples below have been deposited with the American Type Culture

Collection, 10801 University Boulevard, Manassas, VA 20110-2209 under the Budapest Treaty on March 24, 2000 and given the Accession Nos. indicated:

hCLASP-2A 3' clone (AVC-PD1) ATCC accession number PTA-1563

hCLASP-2A 5' clone (AVC-PD2) ATCC accession number PTA-1562

5 hCLASP-2B clone (AVC-PD12) ATCC accession number PTA-1573

The following cDNA clones described in the Specification and further described in the Examples below have been deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209 under the Budapest Treaty on _____ and given the Accession Nos. indicated:

10 hCLASP-2 clone hC2GR3.3 (AVC-PD14) ATCC Accession No. _____

hCLASP-2 clone hC2RT (AVC-PD19) ATCC Accession No. _____

6. EXAMPLES

EXAMPLE 1

Cloning of CLASP-2

The cloning of the CLASP gene family has not been a straightforward process.

5 The cloning of each CLASP family member required the use of multiple techniques and resources. CLASP-2 was cloned in the following manner: an expressed sequence tag or EST clone (IMAGE clone 815795, derived from human germinal B cells) was identified based on a BLAST search of human GenBank human EST database using CLASP-1 sequences. IMAGE clone 815795 was sequenced completely. A polynucleotide probe prepared from
10 815795 sequence was labeled with ^{32}P -dCTP and used to screen human cDNA libraries including Jurkat (Stratagene) and Ramos B cell cDNA library (James Boulter, UCLA). The screening methods employed were as described in Maniatis *et al.*, 1989, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory, New York. Several clones were identified and clone C9, with an insert of 3,752 base pairs, was sequenced (ABI dye-
15 sequencing system, PE Applied Biosystems; Perkin-Elmer Corporation, 761 Main Avenue, Norwalk, CT, U.S.A.). A 5' probe was prepared from C9 sequence and used to rescreen the cDNA libraries. Several clones were isolated, but could not be excised from the phage (Stratagene, CA) without deleting the insert. To circumvent this problem, anchor PCR was performed using M13F primer and CLASP-2 primer (C96AS). The PCR fragment was
20 cloned using the pGEM-T system (Promega), although initial attempts were unsuccessful. The isolated sequence encompassed additional but incomplete cDNA sequence and was determined to carry at least one mutation that may have allowed it to be propagated in bacteria. Commercial libraries from multiple tissue sources including human placenta, B cell, T cell and peripheral blood were exhaustively screened and re-screened resulting in the
25 acquisition of only partial cDNAs. Generation of cDNA libraries using oligo dT or CLASP-specific primers also resulted in the acquisition of partial cDNAs. Genomic libraries were screened to obtain a portion of the genomic locus for each of the CLASP genes, and a genomic walk was initiated to obtain 5' exons and extend the cDNA sequence.

To obtain additional 5' CLASP-2 sequence, portions of the cDNA and
30 genomic sequence from a BAC (Bacterial Artificial Chromosome) genomic library were compared to the NCBI database by BLAST. A genomic clone (Genbank identifier: gi9988160) comprising random, shotgun genomic sequence was identified. Using TFASTX (Pearson and Lipman, PNAS (1988) 85:2444-2448), the amino-terminal sequence of human

CLASP4 was compared to 6 frame translation of gi9988160. Areas of gi9988160 that encoded amino acids with high similarity to CLASP4 amino acid sequence were used to design CLASP-2-specific oligonucleotides for RTPCR (reverse transcriptase polymerase chain reaction according to manufacturers instructions: Reverse transcriptase Gibco/BRL, Taq Polymerase from Sigma). Using oligonucleotides hC2gS5 (nucleotides -66 to -44 of FIG. 11) and C2AS18 (reverse complement of nucleotides 2120 to 2140 of FIG. 11) an RTPCR product of approximately 2.2kb was generated, sequenced (dideoxynucleotide termination sequencing, Beckman Coulter CEQ2000) and shown to be additional human CLASP-2 5' sequence. Further complicating the cloning full-length CLASP cDNA products was the difficulty to clone (and subclone) certain CLASP cDNA products. Standard isolation of some of the CLASP cDNAs from a pure phage population following screening of commercially available cDNA libraries ("ZAP-out" procedure, Stratagene) resulted in no bacterial colonies. Similarly, certain RT-PCR products could not be cloned into standard plasmid vectors. No colonies were isolated by cloning these fragments into vectors lacking promoters, reverse orientations, low copy vectors, or by growth at altered temperatures or levels of antibiotic for plasmid selection (examples: CLASP-7 - HC7gS6 to HC7gAS1 and HC7gS3 to HC7AS14; CLASP-4 - C4P2 to hC4ASTM and C4P2 to HC4AS3'; CLASP-1 - hC1S5' to hC1AS3'Kpn and C1S7 to hC1AS3'Kpn; see Primer Table below). One possibility is that sequences contained within certain regions of CLASP cDNAs are bacteriicidal and therefore not amenable to cloning. To circumvent these problems direct sequencing of RT-PCR products was performed.

Primer Table

CLASP gene	Sense Primer	Sense sequence	Antisense Primer	Antisense sequence
CLASP-7	HC7gS5	AGGCCTTGCTCTGTTTACCTG	HC7gAS1	TGTCATGTACTGCACTCGCACAGC
CLASP-7	HC7gS3	ACAGGAACCTGCTGTACGTGTAC	HC7AS14	TCGTGGCTGCACAGGATGCGGGTG
CLASP-4	C4P2	GACCCATTAGGAGGTCTAC	HC4AS3'	CGGGATCCATTGTCACCGTACATCTGC
CLASP-4	C4P2	GACCCATTAGGAGGTCTAC	HC4AS3'	CGGGATCCATTGTCACCGTACATCTGC
CLASP-1	hC1S5'	TATGTCTCAGTCACCTACCTG	HC1AS3'Kpn	CTTGGTACCACTTCAGCACTAGATGAGATG
CLASP-1	C1S7	TCAAGACCAGGGCATGCAAG	HC1AS3'Kpn	CTTGGTACCACTTCAGCACTAGATGAGATG

In-frame stop codons were not present suggesting that the cDNA was not full length. To obtain the 5' terminus of CLASP-2 5' RACE was employed. Antisense
5 oligonucleotides directed against the 5' end of the longest CLASP-2 sequence were generated:

Primers used for human CLASP-2 5' RACE

	<u>primer</u>	<u>sequence(5' TO 3')</u>	<u>nucleotide position</u>
	HC2RACE1		
10		AAGAGCAGCATCTCCCGTAAACAGTC	-15 to 11
	HC2RACE2		
		TAACAAGCTCTGTGCTTCCTCTTCCG	414 to 443
	HC2RACE3		
		ACCACTTTGTTCGGAAGCTGTCGAAACTC	512 to 540
15	HC2RACE4		
		TTTGTACAGCCAGCCATGCTTGGTGATC	634 to 661

RACE was carried out using Generacer kit (Invitrogen) according to manufacturers specifications using polyA selected mRNA from 9D10 B cell tissue culture line. The sequence of the oligonucleotides presented is the reverse complement (*i.e.*,

antisense) of the the CLASP1 cDNA at the indicated position based upon numbering in FIG. 11.

The full length cDNA (presented in FIG. 11) is therefore a compilation of cDNA from cDNA libraries, RTPCR products and 5' RACE products. The sequence of the CLASP-2 cDNA is shown in FIG. 11.

EXAMPLE 2

Tissue and Cell Line Expression of the CLASP-2 gene

Multiple Tissue Northern blots were purchased from Clontech; hybridization procedures were followed according to manufacturer's procedures and recommendations.

Human T cell line (Jurkat), human myelomonocyte cells (MV4-11), B cells (9D10), monocytes (THP-1), mouse T cells (3A9), mouse B cells (CH27), human promyelocyte (HL60) and human kidney epithelial cells (293 cell line) were maintained as cultured cell lines. For Multiple Cell Northern, RNA was prepared from cell suspensions using the GIBCO-BRL Trizol system. All steps were performed according to the manufacturer's procedures and recommendations. RNA concentrations were determined by the 260nm/280nm light absorption of the RNA solution. 20 µg RNA was ethanol precipitated and resuspended in formamide/formaldehyde buffer and incubated for 15' at 65°C to eliminate putative secondary structures. RNA samples were run over night on a 1.1% agarose gel containing 1.5% formaldehyde (both gel and running buffer were 20 mM sodium phosphate, pH 7.5). To visualize RNA after gel migration, approx. 0.5 µg ethidium bromide was added to each sample prior to the run together with RNA loading buffer. RNA in the gel was then visualized by 260nm wavelength light. After soaking the gel for 15' in deionized water to reduce the concentration of ethidium bromide in the gel, the RNA was transferred onto Amersham Hybond-N plus membrane by capillary blotting in 20 x SSC buffer for 5 hours. Subsequent to blotting, the membrane was washed in 5 x SSC for 3' and RNA was crosslinked to the membrane by UV light (Stratagene Stratalinker).

A probe which recognizes CLASP-2 isoforms A, B, C, and D (probe HC2.2) was used. Probe HC2.2 encompasses to nucleotides 3920 to 4650 (731 bp long) of CLASP-2A. The HC2.2 probe was prepared using standard labeling kits and desalted using pasteur pipette G-50 Sephadex column in TEN (10 mM Tris-HCl, pH 8.0, 1 mM EDTA, and 100 mM NaCl).

Hybridizations of ³²P dCTP labeled DNA probes to the membrane bound RNAs (multiple tissue and multiple cells) were carried out in CLONTECH EXPRESSHYB

solution, at 68°C and for 1-2 hours. Blots were washed 2 times in 2x SSC 0.1% SDS for 10' each at 50°C and then twice in 0.2 x SSC 0.1% SDS for 10' each at 50°C, followed by a 5' wash in 2xSSC at 50°C. Exposure to KODAK BIOMAX MS film was carried out at minus 80°C using amplifying screens. Typical exposure times were 10 to 36 hours.

EXAMPLE 3

Southern Analysis of CLASP-2

BAC DNA was prepared from E. coli over night cultures using the QIAGEN DNA preparation system. All preps were performed according to the manufacturer's procedures, including the modifications for low copy number DNA constructs. Genomic DNA was prepared from HeLa cells (ATCC #CCL-17) using the methods described by Sambrook, Fritsch and Maniatis (1989); DNA concentrations were determined by the 260nm light absorption of the DNA solution, and aliquots corresponding to 20 microgram (µg) genomic DNA or 2 µg for BAC DNA were used for restriction enzyme digests with Eco RI or Hind III (genomic DNA) or Eco RI and Pst I (BAC DNA). Digests were carried out in 150 microliter volume for 4 hours at 37°C. Digested DNA was ethanol precipitated and the pellet was resuspended in 20 microliter deionized water prior to migration over a 1.2 % agarose gel at 35 V over night. Running buffer was TAE, and the gel contained 0.1 µg ethidium bromide/ml to visualize DNA.

Subsequent to gel separation, DNA was visualized by 260 nm wavelength light. The gel was then washed twice for 20' in denaturing buffer (0.5M NaCl, 0.4 N NaOH) and twice in neutralization buffer (1.5 M NaCl, 0.5 M TRIS pH 8.0). DNA was transferred from the gel onto AMERSHAM HYBOND N membrane by capillary blotting in 20 x SSC for 5 hours. The DNA was crosslinked to the membrane by UV light using a Stratagene Stratalinker.

A probe, HC2.1, which recognizes CLASP-2, was used. Probe HC2.1 encompasses nucleotides 325 to 1126 (802 bp long) of CLASP-2A. The HC2.1 probe was prepared using standard labeling kits and desalted using pasteur pipette G-50 Sephadex column in TEN (10 mM Tris-HCl, pH 8.0, 1 mM EDTA, and 100 mM NaCl). Hybridizations of ³²P dCTP labeled DNA against DNA immobilized onto the membrane were carried out at 65°C overnight in modified CHURCH hybridization solution (7% SDS, 0.5 M sodiumphosphate, 1mM EDTA). Membranes were then exposed to KODAK BIOMAX MS film at minus 80°C. Typical exposure times were 12 hours for genomic DNA southern analysis and 3 hours for BAC DNA Southern analysis.

The genomic DNA southern analysis revealed two fragments (~4.5 kb and 1.85 kb) in the Eco RI digested DNA but three fragments in BACs 4 and 6 DNA. The two major bands are identical in both genomic and BAC DNA (FIG. 7).

EXAMPLE 4

CLASP-2 Genomic Cloning

Genomic clones of human CLASP-2 were obtained using the Release I high density filters from Genome Systems Inc (cat # FBAC-4434). Two rounds of screening were completed. The first round of screening was carried out using a probe corresponding to nucleotides 3830 to 4558 of the human CLASP-2 cDNA by standard protocols specific by Genome Systems. This screen identified two genomic clones, referred to as AVC BAC4 and 7. A second round of screening using a probe that corresponded to nucleotides 1208 to 1604 of human CLASP-2 cDNA identified clone AVC BAC26. All the clones were partially sequenced to authenticate that they were indeed CLASP-2 genomic clones, to verify exon sequences, and to identify exon/intron boundaries. Oligonucleotides for sequencing the BACs were based upon human CLASP-2 cDNA sequence. Sense and antisense sequencing oligonucleotides were designed along the length of the human CLASP-2 cDNA spaced approximately every 200 nucleotides to ensure a high density of coverage of the corresponding genomic regions. Sequencing reactions with primers and BAC DNA were carried out by standard PCR sequencing using Big Dye termination sequencing mix (ABI). Results from sequence reactions were analyzed using Sequencher software (Genecodes). The results are summarized in FIG. 6.

EXAMPLE 5

Expression of Recombinant CLASP-2A Polypeptide in Bacterial Cells

Portions of hCLASP-2 were cloned into the GST expression vector pGEX (Pharmacia). These include the region spanning the potential Cadherin processing site through 200 amino acids of the predicted extracellular domain (nucleotide 866 – 1459; GST-EC12; 55 kD fusion) and a portion of the intracellular domain (nucleotide 3230 – 4065; GST-cyto; 57 kD fusion). These regions were amplified using primers at the limits of these sequences on either cDNA clones or cDNA generated from Jurkat or Human Peripheral Blood RNA. Amplified DNA sequences were digested with restriction enzymes for cloning in-frame into GST expression vectors. Fusion proteins were expressed by IPTG induction in DH5 α and purified according to instructions from Pharmacia using glutathione-Sepharose

(Pharmacia). SDS-PAGE gel stained with Coomassie Blue showing induced and uninduced expression of the GST-CLASP-2-cyto construct is shown in FIG. 8. These recombinant proteins were expressed in DH5 α and purified according to instructions from Pharmacia using glutathione-Sepharose. Such recombinant proteins were used to generate antibodies (Josman laboratories) using a AVC Rapid Immunization Protocol.

The full length CLASP can easily be expressed from either the beginning of the hCLASP-2 sequence (in frame with nucleotide 2) or from the first or second methionine (nucleotide 278 or nucleotide 476, underlined in FIG. 1) through to the stop codon (nucleotide 4058). Assuming that the GST moiety has a weight of 26 kD, the total predicted sizes are 180, 168, and 164.5 kD respectively. Alternatively, other bacterial expression systems such as 6CLASP HIS tags, Calmodulin binding protein, maltose binding protein can also be used in a similar manner.

EXAMPLE 6

Expression of Recombinant CLASP-2A Polypeptide in Mammalian Cells

Example 6A. Secreted fusions

Several portions of the predicted extracellular domain were constructed as hIgG fusions using the CD5gamma-1 expression vector (kindly provided by B. Seed, Harvard University). Polypeptides were cloned into this vector in frame with a CD5 leader sequence that directs the fusion protein into the secretory pathway and in frame with a C-terminal hIgG(Fc) protein. This fusion can be secreted from cell lines such as 293 (Hsieh, J-C., 1999, Nature 398: 431-436). Sense primers with hCLASP-2 sequences beginning at nucleotide 866 and antisense primers at nucleotide 1459 (EC12-IgG), nucleotide 2389 (ECC-IgG) and nucleotide 2857 (ECM-IgG) were used to amplify portions of the extracellular domain for insertion into this vector. Recombinant vectors were purified by Maxiprep (Qiagen) and transfected into 293 EBNA- T cells (kindly provided by B. Seed, Harvard University) by calcium phosphate techniques (Sambrook and Maniatis). After 2-7 days, secreted expression was analyzed by an ELISA against the hIgG fusion using a goat F(ab')₂ anti human IgG(Fc) antibody (Jackson Immunolabs) and Protein-A-HRP (Pierce). Intracellular expression was monitored by immunofluorescence microscopy with a FITC labeled goat anti Human IgG(Fc) antibody (Caltag).

Example 6B. Intracellular fusions

Similar methods have been used to construct fusions for expression of full length hCLASP-2 isoforms as well as truncated C-terminal forms in other cell lines such as Jurkat. Recombinant hCLASP-2 fragments were either isolated by digestion of cDNA clones or amplified by primers flanking specific regions (Please provide some specific regions). These can be cloned into expression vectors such as pBJ1-neo (Mark Davis, Stanford University), Peak12 (B. Seed, Harvard University), and pDsRed1-N1 (Clontech). pBJ1-neo and Peak12 allow untagged expression of recombinant proteins and pDsRed1-N1 will allow either untagged or a C-terminal Red fluorescent protein tag. These can be used to generate protein or for expression of various forms for functional analyses.

EXAMPLE 7

Antisense Inhibition of CLASP-2 Expression

Example 7A. Inhibition of CLASP-2 expression *in vitro*

In this example, inhibition of CLASP-2 expression is examined using an *in vitro* cell-free expression system. To identify the useful antisense oligonucleotides, a series of antisense phosphorothioate oligonucleotides (PS-ODNs), which span portion CLASP-2 sequence, can be systematically assayed for the ability to block CLASP-2 expression *in vitro*.

For inhibition of CLASP-2 expression *in vitro*, a CLASP-2 transcription/expression plasmid can be used according to standard methodology for *in vitro* transcription and translation of sense CLASP-2 RNA. Coupled transcription-translation reactions can be performed with a reticulocyte lysate system (Promega TNTTM) according to standard conditions. Each coupled transcription/translation reaction can include CLASP-2 RNA transcribed from the expression plasmid, and a test antisense polynucleotide at a range of standard test concentrations, as well as the luciferase transcription/translation internal control to normalize each reaction (see, *e.g.*, Sambrook *et al.*, *supra*, Ausubel *et al.*, *supra*). The translation reaction can also be performed with sense CLASP-2 RNA that is synthesized *in vitro* in a separate reaction and then added to the translation reaction. ³⁵S-Met is included in the reaction to label the translation products. The negative control is performed without added PS-ODN or a sense PS-ODN.

The labeled translation products can be separated by gel electrophoresis and quantitated after exposing the gel to a phosphorimager screen. The amount of CLASP-2 protein expressed in the presence of CLASP-2 specific PS-ODNs can be normalized to the co-expressed luciferase control.

Example 7B. Inhibition of CLASP-2 expression *ex vivo*

A. Reagents

Cells: Jurkat, Clone E6-1 ATCC TIB-152; 9D10 ATCC CRL8752; additional cells from the ATCC or NCI.

5 Media and solutions: RPMI 1640 medium, BioWhitaker; DMEM/M199 medium, BioWhitaker; EMEM, BioWhitaker; Fetal Bovine Serum, Summit (stored frozen at -20°C, stored thawed at 4°C); Trypsin-EDTA, GIBCO (catalogue #25300-054) (stored frozen at -20°C, stored thawed 4°C); Isoton II (stored at RT); DMSO (stored at RT); oligonucleotides (see Table 1 and FIG. 3, stored in solution at -20°C); PBS (Ca²⁺/Mg²⁺ free); TE; 10 mM Tris-HCL, pH 8.0; 1 mM EDTA.

To prepare oligonucleotide stocks: Oligonucleotide nucleotides (PS-ODNs) can dissolved in the appropriate amount of TE to make a concentrated stock solution (1 - 20 mM).

B. Treatment of cells *ex vivo* with antisense CLASP-2 oligonucleotides

15 Stock cultures of cells in log-phase growth (in T75 flask) can be used. Jurkat, and 9D10 cells are used in this assay. Jurkat and 9D10 are suspension cultures and are passed through dilutions in media. Cell density is measured using a Coulter counter or hemacytometer.

For 6-well dishes, 1.1×10^5 cells total per well, 2 ml/well is added. The
20 amount of cells can be scaled up or down proportionally for 12-well, 100 mm, or 150 mm dishes. For example, for 12-well dishes, use 4.6×10^4 cells in 2 ml media; for 100 mm dishes use 6×10^5 cells in 10 ml media; for 150 mm dishes use 1.7×10^6 cells in 35 ml media.

An appropriate number of cells (as described in step 2 above) are collected, centrifuged and resuspended in media containing a range of ODN concentrations. The cells
25 are treated in single, duplicate, or triplicate wells. Control wells are treated with TE or sense ODNs diluted in media.

The suspension cultures are washed and resuspended daily with PS-ODN media.

Suspension cultures are grown for 2-4 days. Cells are washed with PBS and
30 density measured using a Coulter counter or a hemocytometer. If necessary, the cells are replated at 1.1×10^5 cells per well, 2 ml media per well, and fed with PS-ODN as described above.

Samples of the cells can also be harvested for analysis to determine the effects of CLASP-2 antisense ODNs. Samples are harvested for RNA and analyzed by either Northern analysis or RT-PCR for the presence of CLASP-2 mRNA. Functional consequences of CLASP-2 antisense ODNs can be analyzed by measuring the ability of Jurkat and 9D10 cells to be activated. Jurkat cells are activated by exposure to anti-CD3 and anti-CD28 crosslinking antibodies, and 9D10 cells are activated by exposure to anti-IgM crosslinking antibody or *P. aeruginosa* lipopolysaccharide. A hallmark of activation, calcium influx, can be measured by flow cytometry. Additionally, ELISA assays can be used to measure Interleukin-2 production from Jurkat cells and secreted IgM can be measured using standard assays from 9D10.

Table 5 below shows exemplary oligonucleotides for this assay:

Table 5

Oligo	Sequence 5'-3'	length	notes/comments
1	GAAGGCGATCATCACGT GGCCTTCCATCGC	30-mer	encompasses nucleotides 473-502 and spans the putative initiator methionine (<u>underlined</u>). The function of HC2A, 2B, 2C, and 2E isoforms can be eliminated by this oligonucleotide.
2	GCTTCAAGTAATGACTGG TGCAGAACATCTG	31-mer	Oligonucleotide that should recognize HC2A, 2B, 2D, 2E, and 2F. Encompasses nucleotides 2121-2151. Can be eliminate function of these CLASP-2 isoforms.
3	GCTCCTCCTCAGGCAGGC GCTATGGCTGTGG	34-mer	oligonucleotide specific for HC2C based upon a specific exon found at nucleotide 2927. Can eliminate only HC2D function.
4	GTAGGCCCGGTGCAGCGT GTCATACAGATGG	31-mer	oligonucleotide specific for HC2B, 2C, 2D and 2E based upon specific exon sequence found at nucleotide 3153. Can eliminate function of these CLASP-2 isoforms.
5	GCAATGTCTGAGACTTTC GATCATGAACTATG	32-mer	oligonucleotide specific for HC2A, 2B, 2E, and 2F. Encompasses nucleotides 1987-2018. Can eliminate function of these CLASP-2 isoforms.
6	CAGGAGCTGGTTCTTAAA	18-mer	oligonucleotide specific for HC2A, 2D and 2E. Encompasses nucleotides 2219-2224. Can eliminate function of these CLASP-2 isoforms

Table 5 legend. All nucleotide numeration are relative to Human CLASP-2A (HC2A). See FIG. 2A.

EXAMPLE 8

Example 8A. Synthesis of carboxyl-termini PDZ-ligand peptides

The GST-PDZ fusion proteins are made following standard procedures. An exemplary GST-PDZ fusion protein was constructed as follows: A 572 bp fragment encoding

two PDZ domains of the human neDLG gene (Genbank Accession No. U49089.1) was amplified from total Jurkat RNA by RT-PCR according to standard protocols (Sambrook, Fritsch, and Maniatis, 1989, Molecular Cloning – A Laboratory Manual. Cold Spring Harbor Press.) using primers flanked by restriction endonuclease sites for cloning. Fragments were
5 purified by Sephaglas (Pharmacia), digested with the appropriate enzymes, and ligated into the GST expression vector pGEX-3X (Pharmacia) cut with similar enzymes. Recombinant constructs were confirmed by sequencing. Fusion proteins were expressed by IPTG induction in DH5 α and purified using glutathione-Sepharose (Pharmacia) according to instructions from Pharmacia. Excess glutathione was removed using a PD10 desalting
10 column (Pharmacia) and samples were diaconcentrated by placing the protein in dialysis tubing (14,000 MW cutoff) and laying the tubing on polyethylene glycol (3350; Sigma) until volume had been reduced by approximately 50%. Glycerol was then added to 35% final concentration and samples were stored at –20°C. These recombinant proteins have been used to generate antibodies (Josman laboratories) by standard protocols and for biochemical
15 studies describe herein.

Synthetic peptides corresponding to the carboxyl-terminus of a protein of interest are synthesized by standard resin-based chemistry (*e.g.*, Fmoc), labeled with biotin at the amino-terminus when indicated, and cleaved from the resin using a halide containing acid (*e.g.*, trifluoroacetic acid). The synthetic peptides are then purified by reverse phase
20 high performance liquid chromatography (HPLC) and the identity of the peptides are confirmed by mass spectrometry.

Example 8B. Measurement of CLASP-2 peptide binding to PDZ Domain-containing proteins

The binding of a biotinylated carboxyl-terminal peptide to a GST-PDZ fusion
25 protein is measured as follows:

- (1) GST fusion protein containing one or more PDZ domain(s) is coated onto a protein-binding surface. The protein-binding surface is the surface of a polystyrene plate, which in some cases has been pre-treated by coating with 5 μ g/ml of goat-anti-GST polyclonal antibody followed by blocking with excess bovine serum
30 albumin (BSA). The concentration of GST fusion protein used is 5–10 μ g/ml and the reaction of the GST fusion protein with the plate is carried out in PBS for 1 – 16 hours at

4°C. If not already blocked, the plate is then blocked with BSA (2% in PBS, 2 hours, 4°C)

(2) The plate is washed with PBS.

5 (3) The biotinylated peptide (generally 0.2–20 µM) is then added to the plate and allowed to react in PBS/2% BSA buffer with the GST fusion protein for 10 minutes at 4°C followed by 20 minutes at 25°C. In cases where competition between a labeled (biotinylated) and unlabeled (non-biotinylated) peptide is performed, the unlabeled peptide is added immediately prior to adding the labeled peptide. .

(4) The plate is washed with PBS.

10 (1) 0.5 µg/ml streptavidin-HRP conjugate is added to the plate in PBS/2 % BSA buffer and allowed to react for 20 minutes at 4°C.

(6) The plate is washed 5 X with detergent (tween 20) containing solution.

(7) The plate is developed by addition of HRP-substrate solution for 20 minutes at room temperature.

15 (8) The reaction of the HRP and its substrate is terminated by addition of 1 M sulfuric acid.

(9) The optical density of each well of the plate is read at 450 nm.

In cases where measurement of the apparent affinity of PDZ-ligand interaction is desired, the above procedure is carried out with multiple concentrations of the labeled
20 peptide being used in a single experiment. A plot of binding versus peptide concentration added is then fit to the equation:

$$\text{Binding [peptide]} = \text{Saturation Binding} \times ([\text{peptide}] / ([\text{peptide}] + K_d))$$

where "Binding [peptide]" is the binding of a given concentration of peptide to the GST-PDZ fusion protein minus binding to the GST alone control, "K_d" is the apparent
25 affinity of the binding reaction, and "Saturation Binding" is computed to allow the best fit of the data to the above equation. The term apparent affinity is used because the reaction may

not reach equilibrium during the duration of the binding reaction in which case the apparent affinity would underestimate the actual affinity (*i.e.*, actual K_d < observed K_d).

EXAMPLE 9

Expression of human CLASP -2 in activated T-cells

General experimental design

The expression profiles of human CLASP-2 in T cells upon T cell activation was determined by Northern analysis. Jurkat E6 lymphoblasts were activated by treatment with anti-CD28, PMA, and Ionomycin. Subsequently, total RNA was extracted from cell aliquots harvested at 0, 1, 2, 4, 8, and 14 hours post activation. The RNA concentration of each preparation was determined by the absorption at 260 nm using a spectrophotometer and concentrations of the different RNA preparations were adjusted such that equal quantities of each RNA preparation could be subjected to Northern analysis. Even gel loading was monitored by ethidium bromide staining of the formaldehyde-agarose gel. Northern membranes were hybridized to radioactively labeled probes corresponding to portions of human CLASP-2 and human beta-actin. Expression levels of CLASP-2 at different time points post T-cell activation are proportional to the radioactive signal generated by hybridization by the CLASP-2 specific radioactively labeled probe that remained bound to the Northern membrane under stringent washing conditions. The entire experiment was done in duplicate.

Jurkat E6 cell activation

Jurkat E6 cells were maintained and tested in complete IMDM medium supplemented with 2 mM glutamine, 10 mM HEPES, 100 u/mL penicillin, 100 µg/mL streptomycin, 0.1 mM nonessential amino acids, 1 mM sodium pyruvate (Gibco/BRL), 50 µM beta mercaptoethanol (Sigma), and 10% fetal calf serum (Gemini). T cells were activated as described per Fraser et al., using 0.1 g/mL mouse anti-human CD28 monoclonal antibody (PharMingen International catalog number 33741A), 50 ng/mL PMA (Sigma), and 1 µM ionomycin (Calbiochem). Following incubation at 37°C and 5.0% v/v CO₂, 0.5 x 10⁶ cells were harvested by centrifugation at 500 x g for 10 minutes (min) at room temperature at 0, 1, 2, 4, 8 and 14 hours post activation and subjected to RNA extraction.

For RNA preparation, probe labelling and Northern analysis protocols, see methods and procedures described in Example 2 above. The CLASP-2 specific probe

encompassing nucleotides 5352 to 5922 was generated by PCR from a plasmid containing cloned CLASP-2 cDNA sequences using primers C2S12 and C2AS21.

Hybridization, Washing, and Exposure

Blots were washed twice in 2x SSC 0.1% SDS for 10 min each at 60° C and then twice in 0.2x SSC 0.1% SDS for 10 min each at 60° C, followed by a 5' wash in 2xSSC at 60° C. Exposure to KODAK BIOMAX MS film was carried out at minus 80° C using amplifying screens. Typically, exposure times were 10 to 36 hours. Signal intensities on Northern membranes were quantified by the use of a phosphor imager system (STORM, Molecular Dynamics). Signals were counted in the "volume report" mode.

Results

CLASP-2 expression levels as determined by Northern analysis (FIG. 14) slightly decrease at 1 hour post activation. The maximum decrease of approximately 36 % is seen at 2 hours post activation. Expression levels augment again at 4 hours post activation but do not attain the level that is seen before activation (0 hours). Intensities of CLASP-2-specific signals on the Northern blot were quantified by phosphor imager analysis. Rectangles were drawn around the areas of CLASP-2-specific signal and total quantity of signal was determined by the "volume report" mode; phosphor imager quantification results of two entirely independent experiments are shown in the diagram (green bars corresponds to Northern blot shown). The above result suggests, that transcriptional control of CLASP-2 expression and T-cell activation are functionally linked to each other.

EXAMPLE 10

Chromosomal location of CLASP-2 and possible disease associations
CLASP-2 cDNA sequences have been mapped to the genomic clone (GI:9926440, GI:9988160) by use of sequence homology bioinformatics tools BLAST.

Clone (GI:9926440, GI:9988160) has previously been mapped to the chromosomal location 13q12-q13. The literature research reports that the mutations, deletions, rearrangements, disomies and/or breakpoints (in general: chromosomal aberrations) in below listed genes make the genes strong candidates for the onset of the listed diseases/disorders. Because the CLASP-2 gene is localized in the chromosome location 13q12-q13, abnormal CLASP-2 gene regulation or deletion, rearrangement and/or mutations in CLASP-2 locus might be directly or indirectly associated with the onset of the listed diseases. Further, CLASP-2 gene can be used as a genetic probe to detect the abnormality in

regions of these below listed genes and as a diagnostic marker for the related disease/disorders.

<i>CANDIDATE GENES</i>	<i>LOCUS</i>	<i>RELATED DISEASE/DISORDERS</i>
<u>IPF1:Insulin promoter factor1</u>	13q12.1	MODY4: non insulin-dependent juvenile type, Defect in pancreatic islet development and insulin transcription.
BRCA2	13q12.3	BCLL2: B cell lymphoma, deletion encompassing BRCA2 causes B cell lymphoma. BRCA2 is one of the responsible genes for DNA repairing in S phase.
	13q13.1-q14.3	Deletion of these locus causes MDS6: Myelo dysplastic syndrome type 6 including AML.

5

The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and any clones, DNA or amino acid sequences which are functionally equivalent are within the scope of the invention. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims. It is also to be understood that all base pair sizes given for nucleotides are approximate and are used for purposes of description.

10

15

All publications and patent documents cited above are hereby incorporated by reference in their entirety for all purposes to the same extent as if each were so individually denoted.

WHAT IS CLAIMED IS:

- 1 1. An isolated CLASP-2 polynucleotide, wherein said polynucleotide is
2 (a) a polynucleotide that has the sequence of SEQ ID NO: 1, 3, 5 or 9; or
3 (b) a polynucleotide that hybridizes under stringent hybridization conditions to
4 (a) and encodes a polypeptide having the sequence of SEQ ID NO: 2, 4, 6 or 10 or an allelic
5 variant or homologue of a polypeptide having the sequence of SEQ ID NO: 2, 4, 6 or 10; or
6 (c) a polynucleotide that hybridizes under stringent hybridization conditions to
7 (a) and encodes a polypeptide with at 25 contiguous residues of the polypeptide of SEQ ID
8 NO: 2, 4, 6 or 10; or
9 (d) a polynucleotide that hybridizes under stringent hybridization conditions to
10 (a) and has at least 12 contiguous bases identical to or exactly complementary to SEQ ID NO:
11 1, 3, 5 or 9.
- 1 2. The polynucleotide of claim 1, wherein said polypeptide specifically
2 binds to a PDZ domain of PSD95, DLG1 or neDLG.
- 1 3. The polynucleotide of claim 2, wherein said polypeptide has a binding
2 affinity of at least 10^4 M^{-1} for binding PSD95, DLG1 or neDLG.
- 1 4. The polynucleotide of claim 1 that encodes a polypeptide having the
2 full-length sequence of SEQ ID NO: 2, 4, 6 or 10.
- 1 5. The isolated polynucleotide of claim 1, comprising the cDNA coding
2 sequence of ATCC Deposit Nos. PTA-1562 and PTA-1563 and PTA-1573.
- 1 6. An isolated CLASP-2 polynucleotide comprising a nucleotide
2 sequence that has at least 90% percent identity to SEQ ID NO: 1, 3, 5 or 9.
- 1 7. An isolated polypeptide comprising a nucleotide sequence that has at
2 least 90% sequence identity to SEQ ID NO: 2, 4, 6 or 10 and is immunologically
3 crossreactive with SEQ ID NO: 2, 4, 6 or 10 or shares a biological function with native
4 CLASP-2.
- 1 8. A vector comprising the polynucleotide of claim 1.

1 9. An expression vector comprising the polynucleotide of claim 1 in
2 which the nucleotide sequence of the polynucleotide is operatively linked with a regulatory
3 sequence that controls expression of the polynucleotide in a host cell.

1 10. A host cell comprising the polynucleotide of claim 1, or progeny of the
2 cell.

1 11. A host cell comprising the polynucleotide of claim 1, wherein the
2 nucleotide sequence of the polynucleotide is operatively linked with a regulatory sequence
3 that controls expression of the polynucleotide in a host cell, or progeny of the cell.

1 12. The host cell of claim 10 which is a eukaryote.

1 13. The polynucleotide of claim 1 that is an antisense polynucleotide less
2 than about 200 bases in length.

1 14. An antisense oligonucleotide complementary to a messenger RNA
2 comprising SEQ ID NO: 1, 3, 5 or 9 and encoding CLASP-2, wherein the oligonucleotide
3 inhibits the expression of CLASP-2.

1 15. An isolated DNA that encodes a CLASP-2 protein as shown in SEQ ID
2 NO: 2, 4, 6 or 10.

1 16. The polynucleotide of claim 1 that is RNA.

1 17. A method for producing a polypeptide comprising:
2 (a) culturing the host cell of claim 10 under conditions such that the
3 polypeptide is expressed; and

4 (b) recovering the polypeptide from the cultured host cell or its cultured
5 medium.

1 18. An isolated polypeptide encoded by a polynucleotide of claim 1 (a) or
2 (b).

1 19. The polypeptide of claim 18 that has the amino acid sequence of SEQ
2 ID NO: 2, 4, 6 or 10, or a fragment thereof.

1 20. The isolated polypeptide of claim 18, wherein the polypeptide is cell-
2 membrane associated.

1 21. The isolated polypeptide of claim 18, wherein the polypeptide is
2 soluble.

1 22. The polypeptide of claim 19, wherein the polypeptide is fused with a
2 heterologous polypeptide.

1 23. An isolated CLASP-2 protein having the sequence as shown in SEQ
2 ID NO: 2, 4, 6 or 10.

1 24. A protein comprising the sequence as shown in SEQ. ID. NO: 1 and
2 variants thereof that are at least 95% identical to SEQ ID. NO: 2 and specifically binds
3 spectrin.

1 25. An isolated antibody that specifically binds to a polypeptide having the
2 amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 10, or a binding fragment thereof.

1 26. The antibody of claim 25, that is monoclonal.

1 27. A hybridoma capable of secreting the antibody of claim 26

1 28. A method for identifying a compound or agent that binds a CLASP-2
2 polypeptide comprising:

3 i) contacting a CLASP-2 polypeptide of claim 19 with the compound or agent
4 under conditions which allow binding of the compound to the CLASP-2 polypeptide to form
5 a complex and

6 ii) detecting the presence of the complex.

1 29. A method of detecting a CLASP-2 polypeptide in a sample,
2 comprising:

3 (a) contacting the sample with an antibody or binding fragment of claim 26
4 and (b) determining whether a complex has been formed between the antibody and with
5 CLASP-2 polypeptide.

1 30. A method of detecting a CLASP-2 polypeptide in a sample,
2 comprising:

3 (a) contacting the sample with a polynucleotide of claim 1 or a polynucleotide
4 that comprises a sequence of at least 12 nucleotides and is complementary to a contiguous
5 sequence of the polynucleotide of section (a) of claim 1, and (b) determining whether a
6 hybridization complex has been formed.

1 31. A method of detecting a CLASP-2 nucleotide in a sample, comprising:
2 (a) using a polynucleotide that comprises a sequence of at least 12 nucleotides
3 and is complementary to a contiguous sequence of the polynucleotide of section (a) of claim
4 1, in an amplification process; and

5 (b) determining whether a specific amplification product has been formed.

1 32. A pharmaceutical composition comprising a polynucleotide of claim 1,
2 a polypeptide of claim 18, or an antibody of claim 25 and a pharmaceutically acceptable
3 carrier.

1 33. A method of inhibiting an immune response in a subject comprising:
2 (a) interfering with the expression of a CLASP-2 gene;
3 (b) interfering with the ability of a CLASP-2 protein to bind to another cell;
4 (c) interfering with the ability of a CLASP-2 protein to bind to another protein.

1 34. The method of claim 33, wherein the cell is a T cell or a B cell.

1 35. The method of claim 33 comprising contacting the cell with an
2 effective amount of a polypeptide which comprises the amino acid sequence of SEQ ID NO:
3 2, 4, 6 or 10 or a fragment thereof.

1 36. A method of inhibiting an immune response in a subject, comprising
2 administering to the subject a therapeutically effective amount of an antibody which
3 specifically binds a polypeptide having the sequence of SEQ ID NO: 2, 4, 6 or 10.

1 37. A method of preventing or treating a CLASP-2-mediated disease
2 comprising administering to a subject in need thereof a therapeutically effective amount of a
3 pharmaceutical composition of claim 32.

1 38. The method claim 37, wherein the CLASP-2-mediated disease is an
2 autoimmune disease.

1 39. A method of treating an autoimmune disease in a subject caused or
2 exacerbated by increased activity of T_H1 cells consisting of administering a therapeutically
3 effective amount of a pharmaceutical composition of claim 32 to the subject.

1
A

2 32
 GTT TTA CAC CAT CAC CAA AAC CCA GAA TTT TAT GAT GAG ATT AAA ATA GAG TTG CCC ACT
 val leu his his his gln asn pro glu phe tyr asp glu ile lys ile glu leu pro thr

62 92
 CAG CTG CAT GAA AAG CAC CAC CTG TTG CTC ACA TTC TTC CAT GTC AGC TGT GAC AAC TCA
 gln leu his glu lys his his leu leu leu thr phe phe his val ser cys asp asn ser

122 152
 AGT AAA GGA AGC ACG AAG AAG AGG GAT GTC GTT GAA ACC CAA GTT GGC TAC TCC TGG CTT
 ser lys gly ser thr lys lys arg asp val val glu thr gln val gly tyr ser trp leu

182 212
 CCC CTC CTG AAA GAC GGA AGG GTG GTG ACA AGC GAG CAG CAC ATC CCG GTC TCG GCG AAC
 pro leu leu lys asp gly arg val val thr ser glu gln his ile pro val ser ala asn

242 272
 CTT CCT TCG GGC TAT CTT GGC TAC CAA GAG CTT GGG ATG GGC AGG CAT TAT GGT CCG GAA
 leu pro ser gly tyr leu gly tyr gln glu leu gly met gly arg his tyr gly pro glu

302 332
 ATT AAA TGG GTA GAT GGA GGC AAG CCA CTG CTG AAA ATT TCC ACT CAT CTG GTT TCT ACA
 ile lys trp val asp gly gly lys pro leu leu lys ile ser thr his leu val ser thr

362 392
 GTG TAT ACT CAG GAT CAG CAT TTA CAT AAT TTT TTC CAG TAC TGT CAG AAA ACC GAA TCT
 val tyr thr gln asp gln his leu his asn phe phe gln tyr cys gln lys thr glu ser

422 452
 GGA GCC CAA GCC TTA GGA AAC GAA CTT GTA AAG TAC CTT AAG AGT CTG CAT GCG ATG GAA
 gly ala gln ala leu gly asn glu leu val lys tyr leu lys ser leu his ala met glu

482 512
 GGC CAC GTG ATG ATC GCC TTC TTG CCC ACT ATC CTA AAC CAG CTG TTC CGA GTC CTC ACC
 gly his val met ile ala phe leu pro thr ile leu asn gln leu phe arg val leu thr

542 572
 AGA GCC ACA CAG GAA GAA GTC GCG GTT AAC GTG ACT CGG GTC ATT ATT CAT GTG GTT GCC
 arg ala thr gln glu glu val ala val asn val thr arg val ile ile his val val ala

602 632
 CAG TGC CAT GAG GAA GGA TTG GAG AGC CAC TTG AGG TCA TAT GTT AAG TAC GCG TAT AAG
 gln cys his glu glu gly leu glu ser his leu arg ser tyr val lys tyr ala tyr lys

662 692
 GCT GAG CCA TAT GTT GCC TCT GAA TAC AAG ACA GTG CAT GAA GAA CTG ACC AAA TCC ATG
 ala glu pro tyr val ala ser glu tyr lys thr val his glu glu leu thr lys ser met

FIG. 1


```

722                               752
ACC ACG ATT CTC AAG CCT TCT GCC GAT TTC CTC ACC AGC AAC AAA CTA CTG AGG TAC TCA
thr thr ile leu lys pro ser ala asp phe leu thr ser asn lys leu leu arg tyr ser

782                               812
TGG TTT TTC TTT GAT GTA CTG ATC AAA TCT ATG GCT CAG CAT TTG ATA GAG AAC TCC AAA
trp phe phe phe asp val leu ile lys ser met ala gln his leu ile glu asn ser lys

842                               872
|Cadherin Cleavage|
GTT AAG TTG CTG CGA AAC CAG AGA TTT CCT GCA TCC TAT CAT CAT GCA GCG GAA ACC GTT
val lys leu leu arg asn gln arg phe pro ala ser tyr his his ala ala glu thr val

902                               932
GTA AAT ATG CTG ATG CCA CAC ATC ACT CAG AAG TTT GGA GAT AAT CCA GAG GCA TCT AAG
val asn met leu met pro his ile thr gln lys phe gly asp asn pro glu ala ser lys

962                               992
AAC GCG AAT CAT AGC CTT GCT GTC TTC ATC AAG AGA TGT TTC ACC TTC ATG GAC AGG GGC
asn ala asn his ser leu ala val phe ile lys arg cys phe thr phe met asp arg gly

1022                              1052
TTT GTC TTC AAG CAG ATC AAC AAC TAC ATT AGC TGT TTT GCT CCT GGA GAC CCA AAG ACC
phe val phe lys gln ile asn asn tyr ile ser cys phe ala pro gly asp pro lys thr

1082                              1112
CTC TTT GAA TAC AAG TTT GAA TTT CTC CGT GTA GTG TGC AAC CAT GAA CAT TAT ATT CCG
leu phe glu tyr lys phe glu phe leu arg val val cys asn his glu his tyr ile pro

1142                              1172
TTG AAC TTA CCA ATG CCA TTT GGA AAA GGC AGG ATT CAA AGA TAC CAA GAC CTC CAG CTT
leu asn leu pro met pro phe gly lys gly arg ile gln arg tyr gln asp leu gln leu

1202                              1232
|Cadherin EC
GAC TAC TCA TTA ACA GAT GAG TTC TGC AGA AAC CAC TTC TTG GTG GGA CTG TTA CTG AGG
asp tyr ser leu thr asp glu phe cys arg asn his phe leu val gly leu leu leu arg

1292
GAG GTG GGG ACA GCC CTC CAG GAG TTC CGG GAG GTC CGT CTG ATC GCC ATC AGT GTG CTC
glu val gly thr ala leu gln glu phe arg glu val arg leu ile ala ile ser val leu

1322                              1352
AAG AAC CTG CTG ATA AAG CAT TCT TTT GAT GAC AGA TAT GCT TCA AGG AGC CAT CAG GCA
lys asn leu leu ile lys his ser phe asp asp arg tyr ala ser arg ser his gln ala

1382                              1412/471
AGG ATA GCC ACC CTC TAC CTG CCT CTG TTT GGT CTG CTG ATT GAA AAC GTC CAG CGG ATC
arg ile ala thr leu tyr leu pro leu phe gly leu leu ile glu asn val gln arg ile

1442                              1472
AAT GTG AGG GAT GTG TCA CCC TTC CCT GTG AAC GCG GGC ATG ACC GTG AAG GAT GAA TCC
asn val arg asp val ser pro phe pro val asn ala gly met thr val lys asp glu ser

```

FIG. 1 (cont.)

1502 1532
CTG GCT CTA CCA GCT GTG AAT CCG CTG GTG ACG CCG CAG AAG GGA AGC ACC CTG GAC AAC
leu ala leu pro ala val asn pro leu val thr pro gln lys gly ser thr leu asp asn

1562 1592
AGC CTG CAC AAC CAC CTG CTG GGC GCC ATC TCC GGC ATT GCT TCT CCA TAT ACA ACC TCA
ser leu his lys asp leu leu gly ala ile ser gly ile ala ser pro tyr thr thr ser

1622 1652
ACT CCA AAC ATC AAC AGT GTG AGA AAT GCT GAT TCG AGA GGA TCT CTC ATA AGC ACA GAT
thr pro asn ile asn ser val arg asn ala asp ser arg gly ser leu ile ser thr asp

1682 1712
TCG GGT AAC AGC CTT CCA GAA AGG AAT AGT GAG AAG AGC AAT TCC CTG GAT AAG CAC CAA
ser gly asn ser leu pro glu arg asn ser glu lys ser asn ser leu asp lys his gln

1742 1772
CAA AGT AOC ACA TTG GGA AAT TCC GTG GTT CGC TGT GAT AAA CTT GAC CAG TCT GAG ATT
gln ser ser thr leu gly asn ser val val arg cys asp lys leu asp gln ser glu ile

1802 1832
AAG AGC CTA CTG ATG TGT TTC CTC TAC ATC TTA AAG AGC ATG TCT GAT GAT GCT TTG TTT
lys ser leu leu met cys phe leu tyr ile leu lys ser met ser asp asp ala leu phe

1862 1892
ACA TAT TCG AAC AAG GCT TCA ACA TCT GAA CTT ATG GAT TTT TTT ACA ATA TCT GAA GTC
thr tyr trp asn lys ala ser thr ser glu leu met asp phe phe thr ile ser glu val

1922 1952
TGC CTG CAC CAG TTC CAG TAC ATG GGC AAG CGA TAC ATA GCC AGG AAC CAG GAG GGC TTG
cys leu his gln phe gln tyr met gly lys arg tyr ile ala arg asn gln glu gly leu

1982 2012
GGA CCC ATA CTT CAT GAT CGA AAG TCT CAG ACA TTG CCT GTT TCC CGT AAC AGA ACA GGA
gly pro ile val his asp arg lys ser gln thr leu pro val ser arg asn arg thr gly

2042 2072
ATG ATG CAT GCC AGA TTG CAG CAG CTG GGC AGC CTG GAT AAC TCT CTC ACT TTT AAC CAC
met met his ala arg leu gln gln leu gly ser leu asp asn ser leu thr phe asn his

2102 2132
AGC TAT GGC CAC TCG GAC GCA GAT GTT CTG CAC CAG TCA TTA CTT GAA GCC AAC ATT GCT
ser tyr gly his ser asp ala asp val leu his gln ser leu leu glu ala asn ile ala

2162 2192
ACT GAG GTT TGC CTG ACA GCT CTG GAC ACG CTT TCT CTA TTT ACA TTG GCG TTT AAG AAC
thr glu val cys leu thr ala leu asp thr leu ser leu phe thr leu ala phe lys asn

2222 2252
CAG CTC CTG GCC GAC CAT GGA CAT AAT CCT CTC ATG AAA AAA GTT TTT GAT GTC TAC CTG
gln leu leu ala asp his gly his asn pro leu met lys lys val phe asp val tyr leu

FIG. 1 (cont.)

```

2282                2312
TGT TTT CTT CAA AAA CAT CAG TCT GAA ACG GCT TTA AAA AAT GTC TTC ACT GCC TTA AGG
cys phe leu gln lys his gln ser glu thr ala leu lys asn val phe thr ala leu arg

2342                2372
TCC TTA ATT TAT AAG TTT CCC TCA ACA TTC TAT GAA GGG AGA GCG GAC ATG TGT GCG GCT
ser leu ile tyr lys phe pro ser thr phe tyr glu gly arg ala asp met cys ala ala

2402                2432
CTG TGT TAC GAG ATT CTC AAG TGC TGT AAC TCC AAG CTG AGC TCC ATC AGG ACG GAG GCC
leu cys tyr glu ile leu lys cys cys asn ser lys leu ser ser ile arg thr glu ala

2462                2492
TCC CAG CTG CTC TAC TTC CTG ATG AGG AAC AAC TTT GAT TAC ACT GGA AAG AAG TCC TTT
ser gln leu leu tyr phe leu met arg asn asn phe asp tyr thr gly lys lys ser phe

2522                2552
GTC CGG ACA CAT TTG CAA GTC ATC ATA TCT GTC AGC CAG CTG ATA GCA GAC GTT GTT GGC
val arg thr his leu gln val ile ile ser val ser gln leu ile ala asp val val gly

2582                2612
ATT GGG GAA ACC AGA TTC CAG CAG TCC CTG TCC ATC ATC AAC AAC TGT GCC AAC AGT GAC
ile gly glu thr arg phe gln gln ser leu ser ile ile asn asn cys ala asn ser asp

2642                2672
CGG CTT ATT AAG CAC ACC AGC TTC TCC TCT GAT GTG AAG GAC TTA ACC AAA AGG ATA CGC
arg leu ile lys his thr ser phe ser ser asp val lys asp leu thr lys arg ile arg

2702                2732
ACG GTG CTA ATG GCC ACC GCC CAG ATG AAG GAG CAT GAG AAC GAC CCA GAG ATG CTG GTG
thr val leu met ala thr ala gln met lys glu his glu asn asp pro glu met leu val

2762                2792
GAC CTC CAG TAC AGC CTG GCC AAA TCC TAT GCC AGC ACG CCC GAG CTC AGG AAG ACG TGG
asp leu gln tyr ser leu ala lys ser tyr ala ser thr pro glu leu arg lys thr trp

2822                2852
CTC GAC AGC ATG GCC AGG ATC CAT GTC AAA AAT GGC GAT CTC TCA GAG GCA GCA ATG TGC
leu asp ser met ala arg ile his val lys asn gly asp leu ser glu ala ala met cys

Transmembrane Domain |XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX|
TAT GTC CAC GTA ACA GCC CTA GTG GCA GAA TAT CTC ACA CGG AAA GGC GTG TTT AGA CAA
tyr val his val thr ala leu val ala glu tyr leu thr arg lys gly val phe arg gln

2942                2972
GGA TGC ACC GCC TTC AGG GTC ATT ACC CCA AAC ATC GAC GAG GAG GCC TCC ATG ATG GAA
gly cys thr ala phe arg val ile thr pro asn ile asp glu glu ala ser met met glu

3002                3032
GAC GTG GGG ATG CAG GAT GTC CAT TTC AAC GAG GAT GTG CTG ATG GAG CTC CTT GAG CAG
asp val gly met gln asp val his phe asn glu asp val leu met glu leu leu glu gln

3062                3092
TGC GCA GAT GGA CTC TGG AAA GCC GAG CGC TAC GAG CTC ATC GCC GAC ATC TAC AAA CTT
cys ala asp gly leu trp lys ala glu arg tyr glu leu ile ala asp ile tyr lys leu

```

FIG. 1 (cont.)

```

3122                                     3155
ATC ATC CCC ATT TAT GAG AAG CGG AGG GAT TTC TTT GAA GAT GAA GAT GGA AAG GAG TAT
ile ile pro ile tyr glu lys arg arg asp phe phe glu asp glu asp gly lys glu tyr

3182                                     3212
ATT TAC AAG GAA CCC AAA CTC ACA CCG CTG TCG GAA ATT TCT CAG AGA CTC CTT AAA CTG
ile tyr lys glu pro lys leu thr pro leu ser glu ile ser gln arg leu leu lys leu

3242                                     3272
TAC TCG GAT AAA TTT GGT TCT GAA AAT GTC AAA ATG ATA CAG GAT TCT GGC AAG GTC AAC
tyr ser asp lys phe gly ser glu asn val lys met ile gln asp ser gly lys val asn

3302                                     3332
CCT AAG GAT CTG GAT TCT AAG TAT GCA TAC ATC CAG GTG ACT CAC GTC ATC CCC TTC TTT
pro lys asp leu asp ser lys tyr ala tyr ile gln val thr his val ile pro phe phe

3362                                     3392
GAC GAA AAA GAG TTG CAA GAA AGG AAA ACA GAG TTT GAG AGA TCC CAC AAC ATC CGC CGC
asp glu lys glu leu gln glu arg lys thr glu phe glu arg ser his asn ile arg arg

3422                                     3452
TTC ATG TTT GAG ATG CCA TTT ACG CAG ACC GGG AAG AGG CAG GGC GGG GTG GAA GAG CAG
phe met phe glu met pro phe thr gln thr gly lys arg gln gly gly val glu glu gln

3482                                     3512
TGC AAA CGG CGC ACC ATC CTG ACA GCC ATA CAC TGC TTC CCT TAT GTG AAG AAG CGC ATC
cys lys arg arg thr ile leu thr ala ile his cys phe pro tyr val lys lys arg ile

3542                                     3572 |XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX
CCT GTC ATG TAC CAG CAC CAC ACT GAC CTG AAC CCC ATC GAG GTG GCC ATT GAC GAG ATG
pro val met tyr gln his his thr asp leu asn pro ile glu val ala ile asp glu met

3602 XXX Coiled-Coil 1 XXXXXXXXXXXXXXXX 3632 XXX Coiled-Coil 1 XXXXXXXXXXXXXXXX
AGT AAG AAG GTG GCG GAG CTC CGG CAG CTG TGC TCC TCG GCC GAG GTG GAC ATG ATC AAA
ser lys lys val ala glu leu arg gln leu cys ser ser ala glu val asp met ile lys

3662 XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX | 3692
CTG CAG CTC AAA CTC CAG GGC AGC GTG AGT GTT CAG GTC AAT GCT GGC CCA CTA GCA TAT
leu gln leu lys leu gln gly ser val ser val gln val asn ala gly pro leu ala tyr

3722                                     3752
GCG CGA GCT TTC TTA GAT GAT ACA AAC ACA AAG CGA TAT CCT GAC AAT AAA GTG AAG CTG
ala arg ala phe leu asp asp thr asn thr lys arg tyr pro asp asn lys val lys leu

3782                                     3812 |XXXXXXXXXXXXXXXXXXXXXXXXXXXXX
CTT AAG GAA GTT TTC AGG CAA TTT GTG GAA GCT TGC GGT CAA GCC TTA GCG GTA AAC GAA
leu lys glu val phe arg gln phe val glu ala cys gly gln ala leu ala val asn glu

```

FIG. 1 (cont.)

```

3842 xxx Coiled-Coil 2 xxxxxxxxxxxxxxxx 3872 xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx
CGT CTG ATT AAA GAA GAC CAG CTC GAG TAT CAG GAA GAA ATG AAA GCC AAC TAC AGG GAA
arg leu ile lys glu asp gln leu glu tyr gln glu glu met lys ala asn tyr arg glu

3902 xxx Coiled-Coil 2 xxxxxxxxxxxxxxxx 3932 xxxxxx|
ATG GCG AAG GAG CTT TCT GAA ATC ATG CAT GAG CAG ATC TGC CCC CTG GAG GAG AAG ACG
met ala lys glu leu ser glu ile met his glu gln ile cys pro leu glu glu lys thr

3962
AGC GTC TTA CCG AAT TCC CTT CAC ATC TTC AAC GCC ATC AGT GGG ACT CCA ACA AGC ACA
ser val leu pro asn ser leu his ile phe asn ala ile ser gly thr pro thr ser thr

4022 |xxxxx PBM xxxxxx|
ATG GTT CAC GGG ATG ACC AGC TCG TCT TCG GTC GTG TGA TTA CAT CTC ATG GCC CGT GTG
met val his gly met thr ser ser ser ser val val STP

4082
TGG GGA CTT GCT TTG TCA TTT GCA AAC TCA GGA TGC TTT CCA AAG CCA ATC ACT GGG GAG

4142
ACC GAG CAC AGG GAG GAC CAA GGG GAA GGG GAG AGA AAG GAA ATA AAG AAC AAC GTT ATT

4202
TCT TAA CAG ACT TTC TAT AGG AGT TGT AAG AAG GTG CAC ATA TTT TTT TAA ATC TCA CTG

4262
GCA ATA TTC AAA GTT TTC ATT GTG TCT TAA CAA AGG TGT GGT AGA CAC TCT TGA GCT GGA

4322
CTT AGA TTT TAT TCT TCC TTG CAG AGT AGT GTT AGA ATA GAT GGC CTA CAG AAA AAA AAG

4382
GTT CTG GGA TCT ACA TGG CAG GGA GGG CTG CAC TGA CAT TGA TGC CTG GGG GAC CTT TTG

4442
CCT CGA CTC GTG CCG GAA ATC TGA TCG TAA TCA GGG TAC AGA ACT TAC TAG TTT TGT CTA

4502
GGA GTA TGT TGT ATG ACT AGG ATT TGT GCT ATT ATC TCA TTC AAC AAC ATA GAG CAA GAA

4562
TAG TGA GCT AAC TGA GCT AGA CAC TCA ATT AAT CCG CTA CTG GCT TCA AGT CAG AAC TTT

4622
GTC ATT AAT CAT CGA CTC CGG GAC GGT CAT ATA TGT ATT ACA TTT CTA CAT TTT TAA TAC


4682
TCA CAT GGG CTT ATG CAT TAA GTT TAA TTG TGA TAA ATT TGT GCT GGT CCA GTA TAT GCA

4742
ATA CAC TTT AAT GGT TTA TTC TTG TCA TAA AAA TGT GCA ATA TGG AGA TGT ATA CAA GTC

4802
TTT ACT

```

FIG. 1 (cont.)



1
A

```

2                               32
GTT TTA CAC CAT CAC CAA AAC CCA GAA TTT TAT GAT GAG ATT AAA ATA GAG TTG CCC ACT
val leu his his his gln asn pro glu phe tyr asp glu ile lys ile glu leu pro thr

62                               92
CAG CTG CAT GAA AAG CAC CAC CTG TTG CTC ACA TTC TTC CAT GTC AGC TGT GAC AAC TCA
gln leu his glu lys his his leu leu leu thr phe phe his val ser cys asp asn ser

122                             152
AGT AAA GGA AGC ACG AAG AAG AGG GAT GTC GTT GAA ACC CAA GTT GGC TAC TCC TGG CTT
ser lys gly ser thr lys lys arg asp val val glu thr gln val gly tyr ser trp leu

182                             212
CCC CTC CTG AAA GAC GGA AGG GTG GTG ACA AGC GAG CAG CAC ATC CCG GTC TCG GCG AAC
pro leu leu lys asp gly arg val val thr ser glu gln his ile pro val ser ala asn

242                             272
CTT CCT TCG GGC TAT CTT GGC TAC CAA GAG CTT GGG ATG GGC AGG CAT TAT GGT CCG GAA
leu pro ser gly tyr leu gly tyr gln glu leu gly met gly arg his tyr gly pro glu

302                             332
ATT AAA TGG GTA GAT GGA GGC AAG CCA CTG CTG AAA ATT TCC ACT CAT CTG GTT TCT ACA
ile lys trp val asp gly gly lys pro leu leu lys ile ser thr his leu val ser thr

362                             392
GTG TAT ACT CAG GAT CAG CAT TTA CAT AAT TTT TTC CAG TAC TGT CAG AAA ACC GAA TCT
val tyr thr gln asp gln his leu his asn phe phe gln tyr cys gln lys thr glu ser

422                             452
GGA GCC CAA GCC TTA GGA AAC GAA CTT GTA AAG TAC CTT AAG AGT CTG CAT GCG ATG GAA
gly ala gln ala leu gly asn glu leu val lys tyr leu lys ser leu his ala met glu

482                             512
GGC CAC GTG ATG ATC GCC TTC TTG CCC ACT ATC CTA AAC CAG CTG TTC CGA GTC CTC ACC
gly his val met ile ala phe leu pro thr ile leu asn gln leu phe arg val leu thr

542                             572
AGA GCC ACA CAG GAA GAA GTC GCG GTT AAC GTG ACT CGG GTC ATT ATT CAT GTG GTT GCC
arg ala thr gln glu glu val ala val asn val thr arg val ile ile his val val ala

602                             632
CAG TGC CAT GAG GAA GGA TTG GAG AGC CAC TTG AGG TCA TAT GTT AAG TAC GCG TAT AAG
gln cys his glu glu gly leu glu ser his leu arg ser tyr val lys tyr ala tyr lys

662                             692
GCT GAG CCA TAT GTT GCC TCT GAA TAC AAG ACA GTG CAT GAA GAA CTG ACC AAA TCC ATG
ala glu pro tyr val ala ser glu tyr lys thr val his glu glu leu thr lys ser met

```

FIG. 2B

```

722                               752
ACC ACG ATT CTC AAG CCT TCT GCC GAT TTC CTC ACC AGC AAC AAA CTA CTG AGG TAC TCA
thr thr ile leu lys pro ser ala asp phe leu thr ser asn lys leu leu arg tyr ser

782                               812
TGG TTT TTC TTT GAT GTA CTG ATC AAA TCT ATG GCT CAG CAT TTG ATA GAG AAC TCC AAA
trp phe phe phe asp val leu ile lys ser met ala gln his leu ile glu asn ser lys

842                               872
      |Cadherin Cleavage|
GTT AAG TTG CTG CGA AAC CAG AGA TTT CCT GCA TCC TAT CAT CAT GCA GCG GAA ACC GTT
val lys leu leu arg asn gln arg phe pro ala ser tyr his his ala ala glu thr val

902                               932
GTA AAT ATG CTG ATG CCA CAC ATC ACT CAG AAG TTT GGA GAT AAT CCA GAG GCA TCT AAG
val asn met leu met pro his ile thr gln lys phe gly asp asn pro glu ala ser lys

962                               992
AAC GCG AAT CAT AGC CTT GCT GTC TTC ATC AAG AGA TGT TTC ACC TTC ATG GAC AGG GGC
asn ala asn his ser leu ala val phe ile lys arg cys phe thr phe met asp arg gly

1022                              1052
TTT GTC TTC AAG CAG ATC AAC AAC TAC ATT AGC TGT TTT GCT CCT GGA GAC CCA AAG ACC
phe val phe lys gln ile asn asn tyr ile ser cys phe ala pro gly asp pro lys thr

1082                              1112
CTC TTT GAA TAC AAG TTT GAA TTT CTC CGT GTA GTG TGC AAC CAT GAA CAT TAT ATT CCG
leu phe glu tyr lys phe glu phe leu arg val val cys asn his glu his tyr ile pro

1142                              1172
TTG AAC TTA CCA ATG CCA TTT GGA AAA GGC AGG ATT CAA AGA TAC CAA GAC CTC CAG CTT
leu asn leu pro met pro phe gly lys gly arg ile gln arg tyr gln asp leu gln leu

1202                              1232
      |Cadherin EC
GAC TAC TCA TTA ACA GAT GAG TTC TGC AGA AAC CAC TTC TTG GTG GGA CTG TTA CTG AGG
asp tyr ser leu thr asp glu phe cys arg asn his phe leu val gly leu leu leu arg

xxx|                               1292
GAG GTG GGG ACA GCC CTC CAG GAG TTC CGG GAG GTC CGT CTG ATC GCC ATC AGT GTG CTC
glu val gly thr ala leu gln glu phe arg glu val arg leu ile ala ile ser val leu

1322                              1352
AAG AAC CTG CTG ATA AAG CAT TCT TTT GAT GAC AGA TAT GCT TCA AGG AGC CAT CAG GCA
lys asn leu leu ile lys his ser phe asp asp arg tyr ala ser arg ser his gln ala

1382                              1412
AGG ATA GCC ACC CTC TAC CTG CCT CTG TTT GGT CTG CTG ATT GAA AAC GTC CAG CGG ATC
arg ile ala thr leu tyr leu pro leu phe gly leu leu ile glu asn val gln arg ile

1442                              1472
AAT GTG AGG GAT GTG TCA CCC TTC CCT GTG AAC GCG GGC ATG ACC GTG AAG GAT GAA TCC
asn val arg asp val ser pro phe pro val asn ala gly met thr val lys asp glu ser

```

FIG. 2B (cont.)


```

1502                               1532
CTG GCT CTA CCA GCT GTG AAT CCG CTG GTG ACG CCG CAG AAG GGA AGC ACC CTG GAC AAC
leu ala leu pro ala val asn pro leu val thr pro gln lys gly ser thr leu asp asn

1562                               1592
AGC CTG CAC AAG GAC CTG CTG GGC GCC ATC TCC GGC ATT GCT TCT CCA TAT ACA ACC TCA
ser leu his lys asp leu leu gly ala ile ser gly ile ala ser pro tyr thr thr ser

1622                               1652
ACT CCA AAC ATC AAC AGT GTG AGA AAT GCT GAT TCG AGA GGA TCT CTC ATA AGC ACA GAT
thr pro asn ile asn ser val arg asn ala asp ser arg gly ser leu ile ser thr asp

1682                               1712
TCG GGT AAC AGC CTT CCA GAA AGG AAT AGT GAG AAG AGC AAT TCC CTG GAT AAG CAC CAA
ser gly asn ser leu pro glu arg asn ser glu lys ser asn ser leu asp lys his gln

1742                               1772
CAA AGT AGC ACA TTG GGA AAT TCC GTG GTT CGC TGT GAT AAA CTT GAC CAG TCT GAG ATT
gln ser ser thr leu gly asn ser val val arg cys asp lys leu asp gln ser glu ile

1802                               1832
AAG AGC CTA CTG ATG TGT TTC CTC TAC ATC TTA AAG AGC ATG TCT GAT GAT GCT TTG TTT
lys ser leu leu met cys phe leu tyr ile leu lys ser met ser asp asp ala leu phe

1862                               1892
ACA TAT TGG AAC AAG GCT TCA ACA TCT GAA CTT ATG GAT TTT TTT ACA ATA TCT GAA GTC
thr tyr trp asn lys ala ser thr ser glu leu met asp phe phe thr ile ser glu val

1922                               1952 |xxxxxxxxxxxxxxxxxxxxxxxxxxxx
TGC CTG CAC CAG TTC CAG TAC ATG GGG AAG CGA TAC ATA GCC AGG AAC CAG GAG GGG TTG
cys leu his gln phe gln tyr met gly lys arg tyr ile ala arg asn gln glu gly leu

1982 xxxxxxxxxxxx deleted in CLASP-2D(KIAA1058) xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx|
GGA CCC ATA GTT CAT GAT CGA AAG TCT CAG ACA TTG CCT GTT TCC CGT AAC AGA ACA GGA
gly pro ile val his asp arg lys ser gln thr leu pro val ser arg asn arg thr gly

2042                               2072
ATG ATG CAT GCC AGA TTG CAG CAG CTG GGC AGC CTG GAT AAC TCT CTC ACT TTT AAC CAC
met met his ala arg leu gln gln leu gly ser leu asp asn ser leu thr phe asn his

2102                               2132
AGC TAT GGC CAC TCG GAC GCA GAT GTT CTG CAC CAG TCA TTA CTT GAA GCC AAC ATT GCT
ser tyr gly his ser asp ala asp val leu his gln ser leu leu glu ala asn ile ala

                                                                                               Deleted
2162                               2192 |xxx
ACT GAG GTT TGC CTG ACA GCT CTG GAC ACG CTT TCT CTA TTT ACA TTG GCG TTT AAG AAC
thr glu val cys leu thr ala leu asp thr leu ser leu phe thr leu ala phe lys asn

in HC2B
xxx|
CAG CTC CTG GCC GAC CAT GGA CAT AAT CCT CTC ATG AAA AAA GTT TTT GAT GTC TAC CTG
gln leu leu ala asp his gly his asn pro leu met lys lys val phe asp val tyr leu

```

FIG. 2B (cont.)

2282 2312
TGT TTT CTT CAA AAA CAT CAG TCT GAA ACG GCT TTA AAA AAT GTC TTC ACT GCC TTA AGG
cys phe leu gln lys his gln ser glu thr ala leu lys asn val phe thr ala leu arg

2342 2372
TCC TTA ATT TAT AAG TTT CCC TCA ACA TTC TAT GAA GGG AGA GCG GAC ATG TGT GCG GCT
ser leu ile tyr lys phe pro ser thr phe tyr glu gly arg ala asp met cys ala ala

2402 2432
CTG TGT TAC GAG ATT CTC AAG TGC TGT AAC TCC AAG CTG AGC TCC ATC AGG ACG GAG GCC
leu cys tyr glu ile leu lys cys cys asn ser lys leu ser ser ile arg thr glu ala

2462 2492
TCC CAG CTG CTC TAC TTC CTG ATG AGG AAC AAC TTT GAT TAC ACT GGA AAG AAG TCC TTT
ser gln leu leu tyr phe leu met arg asn asn phe asp tyr thr gly lys lys ser phe

2522 2552
GTC CGG ACA CAT TTG CAA GTC ATC ATA TCT GTC AGC CAG CTG ATA GCA GAC GTT GTT GGC
val arg thr his leu gln val ile ile ser val ser gln leu ile ala asp val val gly

2582 2612
ATT GGG GAA ACC AGA TTC CAG CAG TCC CTG TCC ATC ATC AAC AAC TGT GCC AAC AGT GAC
ile gly glu thr arg phe gln gln ser leu ser ile ile asn asn cys ala asn ser asp

2642 2672
CGG CTT ATT AAG CAC ACC AGC TTC TCC TCT GAT GTG AAG GAC TTA ACC AAA AGG ATA CGC
arg leu ile lys his thr ser phe ser ser asp val lys asp leu thr lys arg ile arg

2702 2732
ACG GTG CTA ATG GCC ACC GCC CAG ATG AAG GAG CAT GAG AAC GAC CCA GAG ATG CTG GTG
thr val leu met ala thr ala gln met lys glu his glu asn asp pro glu met leu val

2762 2792
GAC CTC CAG TAC AGC CTG GCC AAA TCC TAT GCC AGC ACG CCC GAG CTC AGG AAG ACG TGG
asp leu gln tyr ser leu ala lys ser tyr ala ser thr pro glu leu arg lys thr trp

2822 2852 |xxxxxxxxxxxxxxxxxxxx Predicted
CTC GAC AGC ATG GCC AGG ATC CAT GTC AAA AAT GGC GAT CTC TCA GAG GCA GCA ATG TGC
leu asp ser met ala arg ile his val lys asn gly asp leu ser glu ala ala met cys

[Additional and differential exon usage found at position 2927 consisting
of 69 nucleotides. This entire sequence is found in Human CLASP-2D
(KIAA1058) and not other isoforms of CLASP-2. It has a sequence of:
AAGCAGTCCAGTGGGAGCCGCCCTTCTCCCCACAGCCATAGCGCCTGCCTGAGGAGGAGCCGGGGAG]

Transmembrane Domain xx|
TAT GTC CAC GTA ACA GCC CTA GTG GCA GAA TAT CTC ACA CGG AAA GGC GTG TTT AGA CAA
tyr val his val thr ala leu val ala glu tyr leu thr arg lys gly val phe arg gln

2942 2972
GGA TGC ACC GCC TTC AGG GTC ATT ACC CCA AAC ATC GAC GAG GAG GCC TCC ATG ATG GAA
gly cys thr ala phe arg val ile thr pro asn ile asp glu glu ala ser met met glu

FIG. 2B (cont.)

3002 |xx Sequence deleted in CLASP-2E xxxxx
 GAC GTG GGG ATG CAG GAT GTC CAT TTC AAC GAG GAT GTG CTG ATG GAG CTC CTT GAG CAG
 asp val gly met gln asp val his phe asn glu asp val leu met glu leu leu glu gln

3062 xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx 3092
 TGC GCA GAT GGA CTC TGG AAA GCC GAG CGC TAC GAG CTC ATC GCC GAC ATC TAC AAA CTT
 cys ala asp gly leu trp lys ala glu arg tyr glu leu ile ala asp ile tyr lys leu

[Additional and differential exon usage found at position 3153. The entire sequence below is found in Human CLASP-2D. Underlined sequence is found in Human CLASP-2B, 2C and 2E.

TGAGAGGCTGCCCCATCTGTATGACACGCTGCACCGGGCTACAGCAAAGTGACCGAGGTCAT
GCACTCGGGCCGCAGGCTTCTGGGGACCTACTTCCGGGTAGCCTTCTTCGGGCAGGCAGCGCAATACCAGTTT
ACAGACAGTGAAACAGATGTGAGGGATT]

3122 3155
 ATC ATC CCC ATT TAT GAG AAG CGG AGG GAT TTC TTT GAA GAT GAA GAT GGA AAG GAG TAT
 ile ile pro ile tyr glu lys arg arg asp phe phe glu asp glu asp gly lys glu tyr

3182 3212
 ATT TAC AAG GAA CCC AAA CTC ACA CCG CTG TCG GAA ATT TCT CAG AGA CTC CTT AAA CTG
 ile tyr lys glu pro lys leu thr pro leu ser glu ile ser gln arg leu leu lys leu

3242 3272
 TAC TCG GAT AAA TTT GGT TCT GAA AAT GTC AAA ATG ATA CAG GAT TCT GGC AAG GTC AAC
 tyr ser asp lys phe gly ser glu asn val lys met ile gln asp ser gly lys val asn

3302 3332
 CCT AAG GAT CTG GAT TCT AAG TAT GCA TAC ATC CAG GTG ACT CAC GTC ATC CCC TTC TTT
 pro lys asp leu asp ser lys tyr ala tyr ile gln val thr his val ile pro phe phe

3362 3392
 GAC GAA AAA GAG TTG CAA GAA AGG AAA ACA GAG TTT GAG AGA TCC CAC AAC ATC CGC CGC
 asp glu lys glu leu gln glu arg lys thr glu phe glu arg ser his asn ile arg arg

3422 3452
 TTC ATG TTT GAG ATG CCA TTT ACG CAG ACC GGG AAG AGG CAG GGC GGG GTG GAA GAG CAG
 phe met phe glu met pro phe thr gln thr gly lys arg gln gly gly val glu glu gln

3482 3512
 TGC AAA CGG CGC ACC ATC CTG ACA GCC ATA CAC TGC TTC CCT TAT GTG AAG AAG CGC ATC
 cys lys arg arg thr ile leu thr ala ile his cys phe pro tyr val lys lys arg ile

Two nucleotide deletion (nts 3586 and 3587) found in Human CLASP-2C

3542 3572 |xxx|
 CCT GTC ATG TAC CAG CAC CAC ACT GAC CTG AAC CCC ATC GAG GTG GCC ATT GAC GAG ATG
 pro val met tyr gln his his thr asp leu asn pro ile glu val ala ile asp glu met

FIG. 2B (cont.)

```

3602                               3632
AGT AAG AAG GTG GCG GAG CTC CGG CAG CTG TGC TCC TCG GCC GAG GTG GAC ATG ATC AAA
ser lys lys val ala glu leu arg gln leu cys ser ser ala glu val asp met ile lys

3662                               3692
CTG CAG CTC AAA CTC CAG GGC AGC GTG AGT GTT CAG GTC AAT GCT GGC CCA CTA GCA TAT
leu gln leu lys leu gln gly ser val ser val gln val asn ala gly pro leu ala tyr

3722                               3752
GCG CGA GCT TTC TTA GAT GAT ACA AAC ACA AAG CGA TAT CCT GAC AAT AAA GTG AAG CTG
ala arg ala phe leu asp asp thr asn thr lys arg tyr pro asp asn lys val lys leu

3782                               3812
CTT AAG GAA GTT TTC AGG CAA TTT GTG GAA GCT TGC GGT CAA GCC TTA GCG GTA AAC GAA
leu lys glu val phe arg gln phe val glu ala cys gly gln ala leu ala val asn glu

3842                               3872
CGT CTG ATT AAA GAA GAC CAG CTC GAG TAT CAG GAA GAA ATG AAA GCC AAC TAC AGG GAA
arg leu ile lys glu asp gln leu glu tyr gln glu glu met lys ala asn tyr arg glu

Insertion of 8 nucleotides found only in Human CLASP-2D with sequence: CTGGGATG
                                     ↓
3902                               3932
ATG GCG AAG GAG CTT TCT GAA ATC ATG CAT GAG CAG ATC TGC CCC CTG GAG GAG AAG ACG
met ala lys glu leu ser glu ile met his glu gln ile cys pro leu glu glu lys thr

3962                               3992
AGC GTC TTA CCG AAT TCC CTT CAC ATC TTC AAC GCC ATC AGT GGG ACT CCA ACA AGC ACA
ser val leu pro asn ser leu his ile phe asn ala ile ser gly thr pro thr ser thr

4022                               |XXXX PRM XXXX|
ATG GTT CAC GGG ATG ACC AGC TCG TCT TCG GTC GTG TGA TTA CAT CTC ATG GCC CGT GTG
met val his gly met thr ser ser ser ser val val STP

4082                               4112
TGG GGA CTT GCT TTG TCA TTT GCA AAC TCA GGA TGC TTT CCA AAG CCA ATC ACT GGG GAG

4142                               4172
ACC GAG CAC AGG GAG GAC CAA GGG GAA GGG GAG AGA AAG GAA ATA AAG AAC AAC GTT ATT

4202                               4232
TCT TAA CAG ACT TTC TAT AGG AGT TGT AAG AAG GTG CAC ATA TTT TTT TAA ATC TCA CTG

4262                               4292
GCA ATA TTC AAA GTT TTC ATT GTG TCT TAA CAA AGG TGT GGT AGA CAC TCT TGA GCT GGA

4322                               4352
CTT AGA TTT TAT TCT TCC TTG CAG AGT AGT GTT AGA ATA GAT GGC CTA CAG AAA AAA AAG

4382                               4412
GTT CTG GGA TCT ACA TGG CAG GGA GGG CTG CAC TGA CAT TGA TGC CTG GGG GAC CTT TTG

```

FIG. 2B (cont.)

4442
CCT CGA CTC GTG CCG GAA ATC TGA TCG TAA TCA GGG TAC AGA ACT TAC TAG TTT TGT CTA

4472
4502
GGA GTA TGT TGT ATG ACT AGG ATT TGT GCT ATT ATC TCA TTC AAC AAC ATA GAG CAA GAA

4532
4562
TAG TGA GCT AAC TCA GCT AGA CAC TCA ATT AAT CCG CTA CTG GCT TCA AGT CAG AAC TTT

4592
4622
GTC ATT AAT CAT CGA CTC CGG GAC GGT CAT ATA TGT ATT ACA TTT CTA CAT TTT TAA TAC

4652
4682
TCA CAT CGC CTT ATC CAT TAA GTT TAA TTG TGA TAA ATT TGT GCT GGT CCA GTA TAT GCA

4712
4742
ATA CAC TTT AAT GGT TTA TTC TTG TCA TAA AAA TGT GCA ATA TGG AGA TGT ATA CAA GTC

4772
4802
TTT ACT

FIG. 2B (cont.)

HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GCATCTGGAAATCTTGACAAAAATGCCAGATTTTCTGCCATCTACAGGCAAGACAGCAAT
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AAGCTATCCAATGATGACATGCTCAAGTTACTTGCAGACTTTCGGAAACCTGAGAAGATG
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GCTAAGCTCCCACTGATTTTAGGCAATCTAGACATTACAATTGATAATGTTTCCTCAGAC
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TTCCCTAATTATGTTAATTCATCATACATTCCCACAAAACAATTTGAAACCTGCAGTAAA
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	ACTCCCATCACGTTTGAAGTGGAGGAATTTGTGCCCTGCATACCAAAACACACTCAGCCT
HC2E	-----
HC2F	-----

FIG. 3A

HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TACACCATCTACACCAATCACCTTTACGTTTATCCTAAGTACTTGAAATACGACAGTCAG
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AAGTCTTTTGCCAAGGCTAGAAATATTCCGATTTCGATTGAATTCAAAGATTCAGATGAG
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GAAGACTCTCAGCCCCCTTAAGTGCATTTATGGCAGACCTGGTGGGCCAGTTTTCACAAGA
HC2E	-----
HC2F	-----
HC2A	-----AGTTTACACCATCACCAAACCCAGAATTTTATGATGAGATTAAA
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AGCGCCTTTGCTGCAGTTTACACCATCACCAAACCCAGAATTTTATGATGAGATTAAA
HC2E	-----
HC2F	-----
HC2A	ATAGAGTTGCCCACTCAGCTGCATGAAAAGCACCACTGTTGCTCACATTCTTCCATGTC
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	ATAGAGTTGCCCACTCAGCTGCATGAAAAGCACCACTGTTGCTCACATTCTTCCATGTC
HC2E	-----
HC2F	-----
HC2A	AGCTGTGACAACTCAAGTAAAGGAAGCACGAAGAAGAGGGATGTCGTTGAAACCCAAGTT
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AGCTGTGACAACTCAAGTAAAGGAAGCACGAAGAAGAGGGATGTCGTTGAAACCCAAGTT
HC2E	-----
HC2F	-----

FIG. 3A (cont.)

HC2A	GGCTACTCCTGGCTTCCCCCTCCTGAAAGACGGAAGGGTGGTGACAAGCGAGCAGCACATC
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GGCTACTCCTGGCTTCCCCCTCCTGAAAGACGGAAGGGTGGTGACAAGCGAGCAGCACATC
HC2E	-----
HC2F	-----
HC2A	CCGGTCTCGGCGAACCTTCCTTCGGGCTATCTTGGCTACCAAGAGCTTGGGATGGGCAGG
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	CCGGTCTCGGCGAACCTTCCTTCGGGCTATCTTGGCTACCAAGAGCTTGGGATGGGCAGG
HC2E	-----
HC2F	-----
HC2A	CATTATGGTCCGGAAATTAAATGGGTAGATGGAGGCAAGCCACTGCTGAAAATTTCCACT
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	CATTATGGTCCGGAAATTAAATGGGTAGATGGAGGCAAGCCACTGCTGAAAATTTCCACT
HC2E	-----
HC2F	-----
HC2A	CATCTGGTTTCTACAGTGTATACTCAGGATCAGCATTACATAATTTTTTCCAGTACTGT
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	CATCTGGTTTCTACAGTGTATACTCAGGATCAGCATTACATAATTTTTTCCAGTACTGT
HC2E	-----
HC2F	-----
HC2A	CAGAAAACCGAATCTGGAGCCCAAGCCTTAGGAAACGAACTTGTAAGTACCTTAAGAGT
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	CAGAAAACCGAATCTGGAGCCCAAGCCTTAGGAAACGAACTTGTAAGTACCTTAAGAGT
HC2E	-----
HC2F	-----
HC2A	CTGCATGCGATGGAAGGCCACGTGATGATCGCCTTCTTGCCCACTATCCTAAACCAGCTG
HC2-80	-----
HC2B	-----GCGATGGAAGGCCACGTGATGATCGCCTTCTTGCCCACTATCCTAAACCAGCTG
HC2C	-----
HC2D-KIAA1058	CTGCATGCGATGGAAGGCCACGTGATGATCGCCTTCTTGCCCACTATCCTAAACCAGCTG
HC2E	-----GCGATGGAAGGCCACGTGATGATCGCCTTCTTGCCCACTATCCTAAACCAGCTG
HC2F	-----

FIG. 3A (cont.)

HC2A	TTCCGAGTCCTCACCAGAGCCACACAGGAAGAAGTCGCGGTTAACGTGACTCGGGTCATT
HC2-80	-----
HC2B	TTCCGAGTCCTCACCAGAGCCACACAGGAAGAAGTCGCGGTTAACGTGACTCGGGTCATT
HC2C	-----
HC2D-KIAA1058	TTCCGAGTCCTCACCAGAGCCACACAGGAAGAAGTCGCGGTTAACGTGACTCGGGTCATT
HC2E	TTCCGAGTCCTCACCAGAGCCACACAGGAAGAAGTCGCGGTTAACGTGACTCGGGTCATT
HC2F	-----
HC2A	ATTCATGTGGTTGCCCAGTGCCATGAGGAAGGATTGGAGAGCCACTTGAGGTCATATGTT
HC2-80	-----
HC2B	ATTCATGTGGTTGCCCAGTGCCATGAGGAAGGATTGGAGAGCCACTTGAGGTCATATGTT
HC2C	-----
HC2D-KIAA1058	ATTCATGTGGTTGCCCAGTGCCATGAGGAAGGATTGGAGAGCCACTTGAGGTCATATGTT
HC2E	ATTCATGTGGTTGCCCAGTGCCATGAGGAAGGATTGGAGAGCCACTTGAGGTCATATGTT
HC2F	-----
HC2A	AAGTACGCGTATAAGGCTGAGCCATATGTTGCCTCTGAATACAAGACAGTGCATGAAGAA
HC2-80	-----
HC2B	AAGTACGCGTATAAGGCTGAGCCATATGTTGCCTCTGAATACAAGACAGTGCATGAAGAA
HC2C	-----
HC2D-KIAA1058	AAGTACGCGTATAAGGCTGAGCCATATGTTGCCTCTGAATACAAGACAGTGCATGAAGAA
HC2E	AAGTACGCGTATAAGGCTGAGCCATATGTTGCCTCTGAATACAAGACAGTGCATGAAGAA
HC2F	-----
HC2A	CTGACCAAATCCATGACCACGATTCTCAAGCCTTCTGCCGATTTCCTCACCAGCAACAAA
HC2-80	-----
HC2B	CTGACCAAATCCATGACCACGATTCTCAAGCCTTCTGCCGATTTCCTCACCAGCAACAAA
HC2C	-----
HC2D-KIAA1058	CTGACCAAATCCATGACCACGATTCTCAAGCCTTCTGCCGATTTCCTCACCAGCAACAAA
HC2E	CTGACCAAATCCATGACCACGATTCTCAAGCCTTCTGCCGATTTCCTCACCAGCAACAAA
HC2F	-----
HC2A	CTACTGAGGTACTCATGGTTTTTCTTTGATGTACTGATCAAATCTATGGCTCAGCATTG
HC2-80	-----
HC2B	CTACTGAGGTACTCATGGTTTTTCTTTGATGTACTGATCAAATCTATGGCTCAGCATTG
HC2C	-----
HC2D-KIAA1058	CTACTGAAGTACTCATGGTTTTTCTTTGATGTACTGATCAAATCTATGGCTCAGCATTG
HC2E	CTACTGAGGTACTCATGGTTTTTCTTTGATGTACTGATCAAATCTATGGCTCAGCATTG
HC2F	-----
HC2A	ATAGAGAACTCCAAAGTTAAGTTGCTGCGAAACCAGAGATTTCTGCATCCTATCATCAT
HC2-80	-----
HC2B	ATAGAGAACTCCAAAGTTAAGTTGCTGCGAAACCAGAGATTTCTGCATCCTATCATCAT
HC2C	-----
HC2D-KIAA1058	ATAGAGAACTCCAAAGTTAAGTTGCTGCGAAACCAGAGATTTCTGCATCCTATCATCAT
HC2E	ATAGAGAACTCCAAAGTTAAGTTGCTGCGAAACCAGAGATTTCTGCATCCTATCATCAT
HC2F	-----

FIG. 3A (cont.)

HC2A	GCAGCGGAAACCGTTGTAAATATGCTGATGCCACACATCACTCAGAAGTTTGGAGATAAT
HC2-80	-----
HC2B	GCAGCGGAAACCGTTGTAAATATGCTGATGCCACACATCACTCAGAAGTTTGGAGATAAT
HC2C	-----
HC2D-KIAA1058	GCAGTGGAACCGTTGTAAATATGCTGATGCCACACATCACTCAGAAGTTTGGAGATAAT
HC2E	GCAGCGGAAACCGTTGTAAATATGCTGATGCCACACATCACTCAGAAGTTTGGAGATAAT
HC2F	-----
HC2A	CCAGAGGCATCTAAGAACGCGAATCATAGCCTTGCTGTCTTCATCAAGAGATGTTTCACC
HC2-80	-----
HC2B	CCAGAGGCATCTAAGAACGCGAATCATAGCCTTGCTGTCTTCATCAAGAGATGTTTCACC
HC2C	-----
HC2D-KIAA1058	CCAGAGGCATCTAAGAACGCGAATCATAGCCTTGCTGTCTTCATCAAGAGATGTTTCACC
HC2E	CCAGAGGCATCTAAGAACGCGAATCATAGCCTTGCTGTCTTCATCAAGAGATGTTTCACC
HC2F	-----
HC2A	TTCATGGACAGGGGCTTTGTCTTCAAGCAGATCAACAACACTACATTAGCTGTTTTGCTCCT
HC2-80	-----
HC2B	TTCATGGACAGGGGCTTTGTCTTCAAGCAGATCAACAACACTACATTAGCTGTTTTGCTCCT
HC2C	-----
HC2D-KIAA1058	TTCATGGACAGGGGCTTTGTCTTCAAGCAGATCAACAACACTACATTAGCTGTTTTGCTCCT
HC2E	TTCATGGACAGGGGCTTTGTCTTCAAGCAGATCAACAACACTACATTAGCTGTTTTGCTCCT
HC2F	-----
HC2A	GGAGACCCAAAGACCCCTCTTTGAATACAAGTTTGAATTTCTCCGTGTAGTGTGCAACCAT
HC2-80	-----
HC2B	GGAGACCCAAAGACCCCTCTTTGAATACAAGTTTGAATTTCTCCGTGTAGTGTGCAACCAT
HC2C	-----
HC2D-KIAA1058	GGAGACCCAAAGACCCCTCTTTGAATACAAGTTTGAATTTCTCCGTGTAGTGTGCAACCAT
HC2E	GGAGACCCAAAGACCCCTCTTTGAATACAAGTTTGAATTTCTCCGTGTAGTGTGCAACCAT
HC2F	-----
HC2A	GAACATTATATTCCGTTGAACTTACCAATGCCATTGGAAAAGGCAGGATTCAAAGATAC
HC2-80	-----
HC2B	GAACATTATATTCCGTTGAACTTACCAATGCCATTGGAAAAGGCAGGATTCAAAGATAC
HC2C	-----
HC2D-KIAA1058	GAACATTATATTCCGTTGAACTTACCAATGCCATTGGAAAAGGCAGGATTCAAAGATAC
HC2E	GAACATTATATTCCGTTGAACTTACCAATGCCATTGGAAAAGGCAGGATTCAAAGATAC
HC2F	-----
HC2A	CAAGACCTCCAGCTTGACTACTCATTAAACAGATGAGTTCTGCAGAAACCACTTCTTGGTG
HC2-80	-----TCCAGCTTGACTACTCATTAAACAGATGAGTTCTGCAGAAACCACTTCTTGGTG
HC2B	CAAGACCTCCAGCTTGACTACTCATTAAACAGATGAGTTCTGCAGAAACCACTTCTTGGTG
HC2C	-----
HC2D-KIAA1058	CAAGACCTCCAGCTTGACTACTCATTAAACAGATGAGTTCTGCAGAAACCACTTCTTGGTG
HC2E	CAAGACCTCCAGCTTGACTACTCATTAAACAGATGAGTTCTGCAGAAACCACTTCTTGGTG
HC2F	-----

FIG. 3A (cont.)

HC2A	GGACTGTTACTGAGGGAGGTGGGGACAGCCCTCCAGGAGTTCCGGGAGGTCCGTCTGATC
HC2-80	GGACTGTTACTGAGGGAGGTGGGGACAGCCCTCCAGGAGTTCCGGGAGGTCCGTCTGATC
HC2B	GGACTGTTACTGAGGGAGGTGGGGACAGCCCTCCAGGAGTTCCGGGAGGTCCGTCTGATC
HC2C	-----
HC2D-KIAA1058	GGACTGTTACTGAGGGAGGTGGGGACAGCCCTCCAGGAGTTCCGGGAGGTCCGTCTGATC
HC2E	GGACTGTTACTGAGGGAGGTGGGGACAGCCCTCCAGGAGTTCCGGGAGGTCCGTCTGATC
HC2F	-----
HC2A	GCCATCAGTGTGCTCAAGAACCCTGCTGATAAAGCATTCCTTTTGATGACAGATATGCTTCA
HC2-80	GCCATCAGTGTGCTCAAGAACCCTGCTGATAAAGCATTCCTTTTGATGACAGATATGCTTCA
HC2B	GCCATCAGTGTGCTCAAGAACCCTGCTGATAAAGCATTCCTTTTGATGACAGATATGCTTCA
HC2C	-----
HC2D-KIAA1058	GCCATCAGTGTGCTCAAGAACCCTGCTGATAAAGCATTCCTTTTGATGACAGATATGCTTCA
HC2E	GCCATCAGTGTGCTCAAGAACCCTGCTGATAAAGCATTCCTTTTGATGACAGATATGCTTCA
HC2F	-----
HC2A	AGGAGCCATCAGGCAAGGATAGCCACCCTCTACCTGCCTCTGTTTGGTCTGCTGATTGAA
HC2-80	AGGAGCCATCAGGCAAGGATAGCCACCCTCTACCTGCCTCTGTTTGGTCTGCTGATTGAA
HC2B	AGGAGCCATCAGGCAAGGATAGCCACCCTCTACCTGCCTCTGTTTGGTCTGCTGATTGAA
HC2C	-----
HC2D-KIAA1058	AGGAGCCATCAGGCAAGGATAGCCACCCTCTACCTGCCTCTGTTTGGTCTGCTGATTGAA
HC2E	AGGAGCCATCAGGCAAGGATAGCCACCCTCTACCTGCCTCTGTTTGGTCTGCTGATTGAA
HC2F	-----
HC2A	AACGTCCAGCGGATCAATGTGAGGGATGTGTCAACCCTTCCTCTGTGAACGCGGGCATGACC
HC2-80	AACGTCCAGCGGATCAATGTGAGGGATGTGTCAACCCTTCCTCTGTGAACGCGGGCATGACC
HC2B	AACGTCCAGCGGATCAATGTGAGGGATGTGTCAACCCTTCCTCTGTGAACGCGGGCATGACC
HC2C	-----
HC2D-KIAA1058	AACGTCCAGCGGATCAATGTGAGGGATGTGTCAACCCTTCCTCTGTGAACGCGGGCATGACT
HC2E	AACGTCCAGCGGATCAATGTGAGGGATGTGTCAACCCTTCCTCTGTGAACGCGGGCATGACC
HC2F	-----
HC2A	GTGAAGGATGAATCCCTGGCTCTACCAGCTGTGAATCCGCTGGTGACGCCGAGAAGGGA
HC2-80	GTGAAGGATGAATCCCTGGCTCTACCAGCTGTGAATCCGCTGGTGACGCCGAGAAGGGA
HC2B	GTGAAGGATGAATCCCTGGCTCTACCAGCTGTGAATCCGCTGGTGACGCCGAGAAGGGA
HC2C	-----
HC2D-KIAA1058	GTGAAGGATGAATCCCTGGCTCTACCAGCTGTGAATCCGCTGGTGACGCCGAGAAGGGA
HC2E	GTGAAGGATGAATCCCTGGCTCTACCAGCTGTGAATCCGCTGGTGACGCCGAGAAGGGA
HC2F	-----
HC2A	AGCACCTGGACAACAGCCTGCACAAGGACCTGCTGGGCGCCATCTCCGGCATTGCTTCT
HC2-80	AGCACCTGGACAACAGCCTGCACAAGGACCTGCTGGGCGCCATCTCCGGCATTGCTTCT
HC2B	AGCACCTGGACAACAGCCTGCACAAGGACCTGCTGGGCGCCATCTCCGGCATTGCTTCT
HC2C	-----
HC2D-KIAA1058	AGCACCTGGACAACAGCCTGCACAAGGACCTGCTGGGCGCCATCTCCGGCATTGCTTCT
HC2E	AGCACCTGGACAACAGCCTGCACAAGGACCTGCTGGGCGCCATCTCCGGCATTGCTTCT
HC2F	-----

FIG. 3A (cont.)

HC2A	CCATATACAACCTCAACTCCAAACATCAACAGTGTGAGAAATGCTGATTCGAGAGGATCT
HC2-80	CCATATACAACCTCAACTCCAAACATCAACAGTGTGAGAAATGCTGATTCGAGAGGATCT
HC2B	CCATATACAACCTCAACTCCAAACATCAACAGTGTGAGAAATGCTGATTCGAGAGGATCT
HC2C	-----
HC2D-KIAA1058	CCATATACAACCTCAACTCCAAACATCAACAGTGTGAGAAATGCTGATTCGAGAGGATCT
HC2E	CCATATACAACCTCAACTCCAAACATCAACAGTGTGAGAAATGCTGATTCGAGAGGATCT
HC2F	-----GCTGATTCGAGAGGATCT
HC2A	CTCATAAGCACAGATTCGGGTAACAGCCTTCCAGAAAGGAATAGTGAGAAGAGCAATTCC
HC2-80	CTCATAAGCACAGATTCGGGTAACAGCCTTCCAGAAAGGAATAGTGAGAAGAGCAATTCC
HC2B	CTCATAAGCACAGATTCGGGTAACAGCCTTCCAGAAAGGAATAGTGAGAAGAGCAATTCC
HC2C	-----
HC2D-KIAA1058	CTCATAAGCACAGATTCGGGTAACAGCCTTCCAGAAAGGAATAGTGAGAAGAGCAATTCC
HC2E	CTCATAAGCACAGATTCGGGTAACAGCCTTCCAGAAAGGAATAGTGAGAAGAGCAATTCC
HC2F	CTCATAAGCACAGATTCGGGTAACAGCCTTCCAGAAAGGAATAGTGAGAAGAGCAATTCC
HC2A	CTGGATAAGCACCAACAAAGTAGCACATTGGGAAATTCGGTGGTTCGCTGTGATAAACTT
HC2-80	CTGGATAAGCACCAACAAAGTAGCACATTGGGAAATTCGGTGGTTCGCTGTGATAAACTT
HC2B	CTGGATAAGCACCAACAAAGTAGCACATTGGGAAATTCGGTGGTTCGCTGTGATAAACTT
HC2C	-----
HC2D-KIAA1058	CTGGATAAGCACCAACAAAGTAGCACATTGGGAAATTCGGTGGTTCGCTGTGATAAACTT
HC2E	CTGGATAAGCACCAACAAAGTAGCACATTGGGAAATTCGGTGGTTCGCTGTGATAAACTT
HC2F	CTGGATAAGCACCAACAAAGTAGCACATTGGGAAATTCGGTGGTTCGCTGTGATAAACTT
HC2A	GACCAGTCTGAGATTAAGAGCCTACTGATGTGTTTCCTCTACATCTTAAAGAGCATGTCT
HC2-80	GACCAGTCTGAGATTAAGAGCCTACTGATGTGTTTCCTCTACATCTTAAAGAGCATGTCT
HC2B	GACCAGTCTGAGATTAAGAGCCTACTGATGTGTTTCCTCTACATCTTAAAGAGCATGTCT
HC2C	-----
HC2D-KIAA1058	GACCAGTCTGAGATTAAGAGCCTACTGATGTGTTTCCTCTACATCTTAAAGAGCATGTCT
HC2E	GACCAGTCTGAGATTAAGAGCCTACTGATGTGTTTCCTCTACATCTTAAAGAGCATGTCT
HC2F	GACCAGTCTGAGATTAAGAGCCTACTGATGTGTTTCCTCTACATCTTAAAGAGCATGTCT
HC2A	GATGATGCTTTGTTTACATATTGGAACAAGGCTTCAACATCTGAACTTATGGATTTTTTT
HC2-80	GATGATGCTTTGTTTACATATTGGAACAAGGCTTCAACATCTGAACTTATGGATTTTTTT
HC2B	GATGATGCTTTGTTTACATATTGGAACAAGGCTTCAACATCTGAACTTATGGATTTTTTT
HC2C	-----
HC2D-KIAA1058	GATGATGCTTTGTTTACATATTGGAACAAGGCTTCAACATCTGAACTTATGGATTTTTTT
HC2E	GATGATGCTTTGTTTACATATTGGAACAAGGCTTCAACATCTGAACTTATGGATTTTTTT
HC2F	GATGATGCTTTGTTTACATATTGGAACAAGGCTTCAACATCTGAACTTATGGATTTTTTT
HC2A	ACAATATCTGAAGTCTGCCTGCACCAAGTTCCAGTACATGGGGAAGCGATACATAGCCAGG
HC2-80	ACAATATCTGAAGTCTGCCTGCACCAAGTTCCAGTACATGGGGAAGCGATACATAGCCAGG
HC2B	ACAATATCTGAAGTCTGCCTGCACCAAGTTCCAGTACATGGGGAAGCGATACATAGCCAGG
HC2C	-----
HC2D-KIAA1058	ACAATATCTGAAGTCTGCCTGCACCAAGTTCCAGTACATGGGGAAGCGATACATAGCCAG-
HC2E	ACAATATCTGAAGTCTGCCTGCACCAAGTTCCAGTACATGGGGAAGCGATACATAGCCAGG
HC2F	ACAATATCTGAAGTCTGCCTGCACCAAGTTCCAGTACATGGGGAAGCGATACATAGCCAG-

FIG. 3A (cont.)

HC2A AACCAGGAGGGGTTGGGACCCATAGTTCATGATCGAAAGTCTCAGACATTGCCTGTTTCC
HC2-80 AACCAGGAGGGGTTGGGACCCATAGTTCATGATCGAAAGTCTCAGACATTGCCTGTTTCC
HC2B AACCAGGAGGGGTTGGGACCCATAGTTCATGATCGAAAGTCTCAGACATTGCCTGTTTCC
HC2C -----
HC2D-KIAA1058 -----AA-----
HC2E AACCAGGAGGGGTTGGGACCCATAGTTCATGATCGAAAGTCTCAGACATTGCCTGTTTCC
HC2F -----TGTGA-----GAAAG-----ATATCAAGTGT-----

HC2A CGTAACAGAACAGGAATGATGCATGCCAGATTGCAGCAGCTGGGCAGCCTGGATAA ACTCT
HC2-80 CGTAACAGAACAGGAATGATGCATGCCAGATTGCAGCAGCTGGGCAGCCTGGATAA ACTCT
HC2B CGTAACAGAACAGGAATGATGCATGCCAGATTGCAGCAGCTGGGCAGCCTGGATAA ACTCT
HC2C -----
HC2D-KIAA1058 -----CAGGAATGATGCATGCCAGATTGCAGCAGCTGGGCAGCCTGGATAA ACTCT
HC2E CGTAACAGAACAGGAATGATGCATGCCAGATTGCAGCAGCTGGGCAGCCTGGATAA ACTCT
HC2F -----GCTTGGAA-----

HC2A CTCAC TTTTAACCACAGCTATGGCCACTCGGACGCAGATGTTCTGCACCAGTCATTACTT
HC2-80 CTCAC TTTTAACCACAGCTATGGCCACTCGGACGCAGATGTTCTGCACCAGTCATTACTT
HC2B CTCAC TTTTAACCACAGCTATGGCCACTCGGACGCAGATGTTCTGCACCAGTCATTACTT
HC2C -----
HC2D-KIAA1058 CTCAC TTTTAACCACAGCTATGGCCACTCGGACGCAGATGTTCTGCACCAGTCATTACTT
HC2E CTCAC TTTTAACCACAGCTATGGCCACTCGGACGCAGATGTTCTGCACCAGTCATTACTT
HC2F -TTTCTGTAGACAATGGCTATGGCCACTCGGACGCAGATGTTCTGCACCAGTCATTACTT

HC2A GAAGCCAACATTGCTACTGAGGTTTGCCTGACAGCTCTGGACACGCTTTCTCTATT TACA
HC2-80 GAAGCCAACATTGCTACTGAGGTTTGCCTGACAGCTCTGGACACGCTTTCTCTATT TACA
HC2B GAAGCCAACATTGCTACTGAGGTTTGCCTGACAGCTCTGGACACGCTTTCTCTATT TACA
HC2C -----
HC2D-KIAA1058 GAAGCCAACATTGCTACTGAGGTTTGCCTGACAGCTCTGGACACGCTTTCTCTATT TACA
HC2E GAAGCCAACATTGCTACTGAGGTTTGCCTGACAGCTCTGGACACGCTTTCTCTATT TACA
HC2F GAAGCCAACATTGCTACTGAGGTTTGCCTGACAGCTCTGGACACGCTTTCTCTATT TACA

HC2A TTGGCGTTTAAAGAACAGCTCCTGGCCGACCATGGACATAATCCTCTCATGAAAAAAGTT
HC2-80 TTGGCGTTTAAAGAACAGCTCCTGGCCGACCATGGACATAATCCTCTCATGAAAAAAGTT
HC2B TTGGCGTTTAAAG-----CTCCTGGCCGACCATGGACATAATCCTCTCATGAAAAAAGTT
HC2C -----
HC2D-KIAA1058 TTGGCGTTTAAAGAACAGCTCCTGGCCGACCATGGACATAATCCTCTCATGAAAAAAGTT
HC2E TTGGCGTTTAAAGAACAGCTCCTGGCCGACCATGGACATAATCCTCTCATGAAAAAAGTT
HC2F TTGGCGTTTAAAGAACAGCTCCTGGCCGACCATGGACATAATCCTCTCATGAAAAAAGTT

HC2A TTTGATGTCTACCTGTGTTTTCTTCAAAAACATCAGTCTGAAACGGCTTTAAAAAATGTC
HC2-80 TTTGATGTCTACCTGTGTTTTCTTCAAAAACATCAGTCTGAAACGGCTTTAAAAAATGTC
HC2B TTTGATGTCTACCTGTGTTTTCTTCAAAAACATCAGTCTGAAACGGCTTTAAAAAATGTC
HC2C -----
HC2D-KIAA1058 TTTGATGTCTACCTGTGTTTTCTTCAAAAACATCAGTCTGAAACGGCTTTAAAAAATGTC
HC2E TTTGATGTCTACCTGTGTTTTCTTCAAAAACATCAGTCTGAAACGGCTTTAAAAAATGTC
HC2F A-----

FIG. 3A (cont.)

HC2A	TTCAC	TGCCTTAAGGTCCTTAATTTATAAGTTTCCCTCAACATTCTATGAAGGGAGAGCG
HC2-80	TTCAC	TGCCTTAAGGTCCTTAATTTATAAGTTTCCCTCAACATTCTATGAAGGGAGAGCG
HC2B	TTCAC	TGCCTTAAGGTCCTTAATTTATAAGTTTCCCTCAACATTCTATGAAGGGAGAGCG
HC2C	-----	-----
HC2D-KIAA1058	TTCAC	TGCCTTAAGGTCCTTAATTTATAAGTTTCCCTCAACATTCTATGAAGGGAGAGCG
HC2E	TTCAC	TGCCTTAAGGTCCTTAATTTATAAGTTTCCCTCAACATTCTATGAAGGGAGAGCG
HC2F	-----	-----
HC2A	GACATG	TGTCGGCTCTGTGTTACGAGATTCTCAAGTGCTGTAACCTCAAGCTGAGCTCC
HC2-80	GACATG	TGTCGGCTCTGTGTTACGAGATTCTCAAGTGCTGTAACCTCAAGCTGAGCTCC
HC2B	GACATG	TGTCGGCTCTGTGTTACGAGATTCTCAAGTGCTGTAACCTCAAGCTGAGCTCC
HC2C	-----	-----
HC2D-KIAA1058	GACATG	TGTCGGCTCTGTGTTACGAGATTCTCAAGTGCTGTAACCTCAAGCTGAGCTCC
HC2E	GACATG	TGTCGGCTCTGTGTTACGAGATTCTCAAGTGCTGTAACCTCAAGCTGAGCTCC
HC2F	-----	-----
HC2A	ATCAGG	ACGGAGGCCTCCAGCTGCTCTACTTCCTGATGAGGAACAACTTTGATTACACT
HC2-80	ATCAGG	ACGGAGGCCTCCAGCTGCTCTACTTCCTGATGAGGAACAACTTTGATTACACT
HC2B	ATCAGG	ACGGAGGCCTCCAGCTGCTCTACTTCCTGATGAGGAACAACTTTGATTACACT
HC2C	-----	-----
HC2D-KIAA1058	ATCAGG	ACGGAGGCCTCCAGCTGCTCTACTTCCTGATGAGGAACAACTTTGATTACACT
HC2E	ATCAGG	ACGGAGGCCTCCAGCTGCTCTACTTCCTGATGAGGAACAACTTTGATTACACT
HC2F	-----	-----
HC2A	GGAAAGA	AGTCCTTTGTCCGGACACATTTGCAAGTCATCATATCTGTGTCAGCCAGCTGATA
HC2-80	GGAAAGA	AGTCCTTTGTCCGGACACATTTGCAAGTCATCATATCTGTGTCAGCCAGCTGATA
HC2B	GGAAAGA	AGTCCTTTGTCCGGACACATTTGCAAGTCATCATATCTGTGTCAGCCAGCTGATA
HC2C	-----	-----
HC2D-KIAA1058	GGAAAGA	AGTCCTTTGTCCGGACACATTTGCAAGTCATCATATCTGTGTCAGCCAGCTGATA
HC2E	GGAAAGA	AGTCCTTTGTCCGGACACATTTGCAAGTCATCATATCTGTGTCAGCCAGCTGATA
HC2F	-----	-----
HC2A	GCAGAC	GTTGTTGGCATTGGGGAAACCAGATTCCAGCAGTCCCTGTCCATCATCAACAAC
HC2-80	GCAGAC	GTTGTTGGCATTGGGGAAACCAGATTCCAGCAGTCCCTGTCCATCATCAACAAC
HC2B	GCAGAC	GTTGTTGGCATTGGGGAAACCAGATTCCAGCAGTCCCTGTCCATCATCAACAAC
HC2C	-----	-----
HC2D-KIAA1058	GCAGAC	GTTGTTGGCATTGGGGAAACCAGATTCCAGCAGTCCCTGTCCATCATCAACAAC
HC2E	GCAGAC	GTTGTTGGCATTGGGGAAACCAGATTCCAGCAGTCCCTGTCCATCATCAACAAC
HC2F	-----	-----
HC2A	TGTGCC	AACAGTGACCGGCTTATTAAGCACACCAGCTTCTCCTCTGATGTGAAGGACTTA
HC2-80	TGTGCC	AACAGTGACCGGCTTATTAAGCACACCAGCTTCTCCTCTGATGTGAAGGACTTA
HC2B	TGTGCC	AACAGTGACCGGCTTATTAAGCACACCAGCTTCTCCTCTGATGTGAAGGACTTA
HC2C	-----	-----
HC2D-KIAA1058	TGTGCC	AACAGTGACCGGCTTATTAAGCACACCAGCTTCTCCTCTGATGTGAAGGACTTA
HC2E	TGTGCC	AACAGTGACCGGCTTATTAAGCACACCAGCTTCTCCTCTGATGTGAAGGACTTA
HC2F	-----	-----

FIG. 3A (cont.)

HC2A	ACCAAAAGGATACGCACGGTGCTAATGGCCACCGCCAGATGAAGGAGCATGAGAACGAC
HC2-80	ACCAAAAGGATACGCACGGTGCTAATGGCCACCGCCAGATGAAGGAGCATGAGAACGAC
HC2B	ACCAAAAGGATACGCACGGTGCTAATGGCCACCGCCAGATGAAGGAGCATGAGAACGAC
HC2C	-----
HC2D-KIAA1058	ACCAAAAGGATACGCACGGTGCTAATGGCCACCGCCAGATGAAGGAGCATGAGAACGAC
HC2E	ACCAAAAGGATACGCACGGTGCTAATGGCCACCGCCAGATGAAGGAGCATGAGAACGAC
HC2F	-----
HC2A	CCAGAGATGCTGGTGGACCTCCAGTACAGCCTGGCCAAATCCTATGCCAGCAGCGCCGAG
HC2-80	CCAGAGATGCTGGTGGACCTCCAGTACAGCCTGGCCAAATCCTATGCCAGCAGCGCCGAG
HC2B	CCAGAGATGCTGGTGGACCTCCAGTACAGCCTGGCCAAATCCTATGCCAGCAGCGCCGAG
HC2C	-----
HC2D-KIAA1058	CCAGAGATGCTGGTGGACCTCCAGTACAGCCTGGCCAAATCCTATGCCAGCAGCGCCGAG
HC2E	CCAGAGATGCTGGTGGACCTCCAGTACAGCCTGGCCAAATCCTATGCCAGCAGCGCCGAG
HC2F	-----
HC2A	CTCAGGAAGACGTGGCTCGACAGCATGGCCAGGATCCATGTCAAAAATGGCGATCTCTCA
HC2-80	CTCAGGAAGACGTGGCTCGACAGCATGGCCAGGATCCATGTCAAAAATGGCGATCTCTCA
HC2B	CTCAGGAAGACGTGGCTCGACAGCATGGCCAGGATCCATGTCAAAAATGGCGATCTCTCA
HC2C	-----
HC2D-KIAA1058	CTCAGGAAGACGTGGCTCGACAGCATGGCCAGGATCCATGTCAAAAATGGCGATCTCTCA
HC2E	CTCAGGAAGACGTGGCTCGACAGCATGGCCAGGATCCATGTCAAAAATGGCGATCTCTCA
HC2F	-----
HC2A	GAGGCAGCAATGTGCTATGTCCACGTAACAGCCCTAGTGGCAGAATATCTCACACGGAAA
HC2-80	GAGGCAGCAATGTGCTATGTCCACGTAACAGCCCTAGTGGCAGAATATCTCACACGGAAA
HC2B	GAGGCAGCAATGTGCTATGTCCACGTAACAGCCCTAGTGGCAGAATATCTCACACGGAAA
HC2C	-----
HC2D-KIAA1058	GAGGCAGCAATGTGCTATGTCCACGTAACAGCCCTAGTGGCAGAATATCTCACACGGAAA
HC2E	GAGGCAGCAATGTGCTATGTCCACGTAACAGCCCTAGTGGCAGAATATCTCACACGGAAA
HC2F	-----
HC2A	G-----
HC2-80	G-----
HC2B	G-----
HC2C	-----
HC2D-KIAA1058	GAAGCAGTCCAGTGGGAGCCGCCCTTCTCCCCACAGCCATAGCGCCTGCCTGAGGAGG
HC2E	G-----
HC2F	-----
HC2A	-----GCGTGTTTAGACAAGGATGCACCGCCTTCAGGGTCATTACCCCAAACATC
HC2-80	-----GCGTGTTTAGACAAGGATGCACCGCCTTCAGGGTCATTACCCCAAACATC
HC2B	-----GCGTGTTTAGACAAGGATGCACCGCCTTCAGGGTCATTACCCCAAACATC
HC2C	-----GTTTAGACAAGGATGCACCGCCTTCAGGGTCATTACCCCAAACATC
HC2D-KIAA1058	AGCCGGGGAGGCGTGTTTAGACAAGGATGCACCGCCTTCAGGGTCATTACCCCAAACATC
HC2E	-----GCGTGTTTAGACAAGGATGCACCGCCTTCAGGGTCATTACCCCAAACATC
HC2F	-----

FIG. 3A (cont.)

```
HC2A      GACGAGGAGGCCTCCATGATGGAAGACGTGGGGATGCAGGATGTCCATTTC AACGAGGAT
HC2-80    GACGAGGAGGCCTCCATGATGGAAGACGTGGGGATGCAGGATGTCCATTTC AACGAGGAT
HC2B      GACGAGGAGGCCTCCATGATGGAAGACGTGGGGATGCAGGATGTCCATTTC AACGAGGAT
HC2C      GACGAGGAGGCCTCCATGATGGAAGACGTGGGGATGCAGGATGTCCATTTC AACGAGGAT
HC2D-KIAA1058 GACGAGGAGGCCTCCATGATGGAAGACGTGGGGATGCAGGATGTCCATTTC AACGAGGAT
HC2E      GACGAGGAGGCCTCCATGATGGAAGACGTGGGGA-----
HC2F      -----

HC2A      GTGCTGATGGAGCTCCTTGAGCAGTGCGCAGATGGACTCTGGAAGCCGAGCGCTACGAG
HC2-80    GTGCTGATGGAGCTCCTTGAGCAGTGCGCAGATGGACTCTGGAAGCCGAGCGCTACGAG
HC2B      GTGCTGATGGAGCTCCTTGAGCAGTGCGCAGATGGACTCTGGAAGCCGAGCGCTACGAG
HC2C      GTGCTGATGGAGCTCCTTGAGCAGTGCGCAGATGGACTCTGGAAGCCGAGCGCTACGAG
HC2D-KIAA1058 GTGCTGATGGAGCTCCTTGAGCAGTGCGCAGATGGACTCTGGAAGCCGAGCGCTACGAG
HC2E      -----AAGCCGAGCGCTACGAG
HC2F      -----

HC2A      CTCATCGCCGACATCTACAACTTATCATCCCCATTTATGAGAAGCGGAGGGATTT----
HC2-80    CTCATCGCCGACATCTACAACTTATCATCCCCATTTATGAGAAGCGGAGGGATTT----
HC2B      CTCATCGCCGACATCTACAACTTATCATCCCCATTTATGAGAAGCGGAGGGATTTTGAG
HC2C      CTCATCGCCGACATCTACAACTTATCATCCCCATTTATGAGAAGCGGAGGGATTTTGAG
HC2D-KIAA1058 CTCATTGCCGACATCTACAACTTATCATCCCCATTTATGAGAAGCGGAGGGATTTTGAG
HC2E      CTCATCGCCGACATCTACAACTTATCATCCCCATTTATGAGAAGCGGAGGGATTTTGAG
HC2F      -----

HC2A      -----
HC2-80    -----
HC2B      AGGCTGGCCCATCTGTATGACACGCTGCACCGGGCCTACAGCAAAGTGACCGAGGTCATG
HC2C      AGGCTGGCCCATCTGTATGACACGCTGCACCGGGCCTACAGCAAAGTGACCGAGGTCATG
HC2D-KIAA1058 AGGCTGGCCCATCTGTATGACACGCTGCACCGGGCCTACAGCAAAGTGACCGAGGTCATG
HC2E      AGGCTGGCCCATCTGTATGACACGCTGCACCGGGCCTACAGCAAAGTGACCGAGGTCATG
HC2F      -----

HC2A      -----
HC2-80    -----
HC2B      CACTCGGGCCGACAGGCTTCTGGGGACCTACTTCCGGGTAGCCTTCTTCGGGCAGG----
HC2C      CACTCGGGCCGACAGGCTTCTGGGGACCTACTTCCGGGTAGCCTTCTTCGGGCAGG----
HC2D-KIAA1058 CACTCGGGCCGACAGGCTTCTGGGGACCTACTTCCGGGTAGCCTTCTTCGGGCAGGCAGCG
HC2E      CACTCGGGCCGACAGGCTTCTGGGGACCTACTTCCGGGTAGCCTTCTTCGGGCAGG----
HC2F      -----

HC2A      -----CTTTGAAGATGAAGATGGA
HC2-80    -----CTTTGAAGATGAAGATGGA
HC2B      -----GATTCTTTGAAGATGAAGATGGA
HC2C      -----GATTCTTTGAAGATGAAGATGGA
HC2D-KIAA1058 CAATACCAGTTTACAGACAGTGAAACAGATGTGGAGGGATTCTTTGAAGATGAAGATGGA
HC2E      -----GATTCTTTGAAGATGAAGATGGA
HC2F      -----
```

FIG. 3A (cont.)

HC2A	AAGGAGTATATTTACAAGGAACCCAAACTCACACCGCTGTCTCGGAAATTTCTCAGAGACTC
HC2-80	AAGGAGTATATTTACAAGGAACCCAAACTCACACCGCTGTCTCGGAAATTTCTCAGAGACTC
HC2B	AAGGAGTATATTTACAAGGAACCCAAACTCACACCGCTGTCTCGGAAATTTCTCAGAGACTC
HC2C	AAGGAGTATATTTACAAGGAACCCAAACTCACACCGCTGTCTCGGAAATTTCTCAGAGACTC
HC2D-KIAA1058	AAGGAGTATATTTACAAGGAACCCAAACTCACACCGCTGTCTCGGAAATTTCTCAGAGACTC
HC2E	AAGGAGTATATTTACAAGGAACCCAAACTCACACCGCTGTCTCGGAAATTTCTCAGAGACTC
HC2F	-----
HC2A	CTTAAACTGTACTCGGATAAAATTTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGC
HC2-80	CTTAAACTGTACTCGGATAAAATTTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGC
HC2B	CTTAAACTGTACTCGGATAAAATTTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGC
HC2C	CTTAAACTGTACTCGGATAAAATTTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGC
HC2D-KIAA1058	CTTAAACTGTACTCGGATAAAATTTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGC
HC2E	CTTAAACTGTACTCGGATAAAATTTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGC
HC2F	-----
HC2A	AAGGTCAACCCCTAAGGATCTGGATTCTAAGTATGCATACATCCAGGTGACTCACGTCATC
HC2-80	AAGGTCAACCCCTAAGGATCTGGATTCTAAGTATGCATACATCCAGGTGACTCACGTCATC
HC2B	AAGGTCAACCCCTAAGGATCTGGATTCTAAGTATGCATACATCCAGGTGACTCACGTCATC
HC2C	AAGGTCAACCCCTAAGGATCTGGATTCTAAGTATGCATACATCCAGGTGACTCACGTCATC
HC2D-KIAA1058	AAGGTCAACCCCTAAGGATCTGGATTCTAAGTATGCCTACATCCAGGTGACTCACGTCATC
HC2E	AAGGTCAACCCCTAAGGATCTGGATTCTAAGTATGCATACATCCAGGTGACTCACGTCATC
HC2F	-----
HC2A	CCCTTCTTTGACGAAAAAGAGTTGCAAGAAAGGAAAAACAGAGTTTGAGAGATCCCACAAC
HC2-80	CCCTTCTTTGACGAAAAAGAGTTGCAAGAAAGGAAAAACAGAGTTTGAGAGATCCCACAAC
HC2B	CCCTTCTTTGACGAAAAAGAGTTGCAAGAAAGGAAAAACAGAGTTTGAGAGATCCCACAAC
HC2C	CCCTTCTTTGACGAAAAAGAGTTGCAAGAAAGGAAAAACAGAGTTTGAGAGATCCCACAAC
HC2D-KIAA1058	CCCTTCTTTGACGAAAAAGAGTTGCAAGAAAGGAAAAACAGAGTTTGAGAGATCCCACAAC
HC2E	CCCTTCTTTGACGAAAAAGAGTTGCAAGAAAGGAAAAACAGAGTTTGAGAGATCCCACAAC
HC2F	-----
HC2A	ATCCGCCGCTTCATGTTTGAGATGCCATTTACGCAGACCGGGAAGAGGCAGGGCGGGGTG
HC2-80	ATCCGCCGCTTCATGTTTGAGATGCCATTTACGCAGACCGGGAAGAGGCAGGGCGGGGTG
HC2B	ATCCGCCGCTTCATGTTTGAGATGCCATTTACGCAGACCGGGAAGAGGCAGGGCGGGGTG
HC2C	ATCCGCCGCTTCATGTTTGAGATGCCATTTACGCAGACCGGGAAGAGGCAGGGCGGGGTG
HC2D-KIAA1058	ATCCGCCGCTTCATGTTTGAGATGCCATTTACGCAGACCGGGAAGAGGCAGGGCGGGGTG
HC2E	ATCCGCCGCTTCATGTTTGAGATGCCATTTACGCAGACCGGGAAGAGGCAGGGCGGGGTG
HC2F	-----
HC2A	GAAGAGCAGTGCAAACGGCGCACCATCCTGACAGCCATACACTGCTTCCCTTATGTGAAG
HC2-80	GAAGAGCAGTGCAAACGGCGCACCATCCTGACAGCCATACACTGCTTCCCTTATGTGAAG
HC2B	GAAGAGCAGTGCAAACGGCGCACCATCCTGACAGCCATACACTGCTTCCCTTATGTGAAG
HC2C	GAAGAGCAGTGCAAACGGCGCACCATCCTGACAGCCATACACTGCTTCCCTTATGTGAAG
HC2D-KIAA1058	GAAGAGCAGTGCAAACGGCGCACCATCCTGACAGCCATACACTGCTTCCCTTATGTGAAG
HC2E	GAAGAGCAGTGCAAACGGCGCACCATCCTGACAGCCATACACTGCTTCCCTTATGTGAAG
HC2F	-----

FIG. 3A (cont.)

HC2A	AAGCGCATCCCTGTCATGTACCAGCACCACACTGACCTGAACCCCATCGAGGTGGCCATT
HC2-80	AAGCGCATCCCTGTCATGTACCAGCACCACACTGACCTGAACCCCATCGAGGTGGCCATT
HC2B	AAGCGCATCCCTGTCATGTACCAGCACCACACTGACCTGAACCCCATCGAGGTGGCCATT
HC2C	AAGCGCATCCCTTTTCATGTACCAGCACCACACTGACCTGAACCCCATCGAGGT--CCATT
HC2D-KIAA1058	AAGCGCATCCCTGTCATGTACCAGCACCACACTGACCTGAACCCCATCGAGGTGGCCATT
HC2E	AAGCGCATCCCTGTCATGTACCAGCACCACACTGACCTGAACCCCATCGAGGTGGCCATT
HC2F	-----
HC2A	GACGAGATGAGTAAGAAGGTGGCGGAGCTCCGGCAGCTGTGCTCCTCGGCCGAGGTGGAC
HC2-80	GACGAGATGAGTAAGAAGGTGGCGGAGCTCCGGCAGCTGTGCTCCTCGGCCGAGGTGGAC
HC2B	GACGAGATGAGTAAGAAGGTGGCGGAGCTCCGGCAGCTGTGCTCCTCGGCCGAGGTGGAC
HC2C	GACGAGATGAGTAAGAAGGTGGCGGAGCTCCGGCAGCTGTGCTCCTCGGCCGAGGTGGAC
HC2D-KIAA1058	GACGAGATGAGTAAGAAGGTGGCGGAGCTCCGGCAGCTGTGCTCCTCGGCCGAGGTGGAC
HC2E	GACGAGATGAGTAAGAAGGTGGCGGAGCTCCGGCAGCTGTGCTCCTCGGCCGAGGTGGAC
HC2F	-----
HC2A	ATGATCAAACCTGCAGCTCAAACCTCCAGGGCAGCGTGAGTGTTTCAGGTCAATGCTGGCCCA
HC2-80	ATGATCAAACCTGCAGCTCAAACCTCCAGGGCAGCGTGAGTGTTTCAGGTCAATGCTGGCCCA
HC2B	ATGATCAAACCTGCAGCTCAAACCTCCAGGGCAGCGTGAGTGTTTCAGGTCAATGCTGGCCCA
HC2C	ATGATCAAACCTGCAGCTCAAACCTCCAGGGCAGCGTGAGTGTTTCAGGTCAATGCTGGCCCA
HC2D-KIAA1058	ATGATCAAACCTGCAGCTCAAACCTCCAGGGCAGCGTGAGTGTTTCAGGTCAATGCTGGCCCA
HC2E	ATGATCAAACCTGCAGCTCAAACCTCCAGGGCAGCGTGAGTGTTTCAGGTCAATGCTGGCCCA
HC2F	-----
HC2A	CTAGCATATGCGCGAGCTTTCTTAGATGATACAAACACAAAGCGATATCCTGACAATAAA
HC2-80	CTAGCATATGCGCGAGCTTTCTTAGATGATACAAACACAAAGCGATATCCTGACAATAAA
HC2B	CTAGCATATGCGCGAGCTTTCTTAGATGATACAAACACAAAGCGATATCCTGACAATAAA
HC2C	CTAGCATATGCGCGAGCTTTCTTAGATGATACAAACACAAAGCGATATCCTGACAATAAA
HC2D-KIAA1058	CTAGCATATGCGCGAGCTTTCTTAGATGATACAAACACAAAGCGATATCCTGACAATAAA
HC2E	CTAGCATATGCGCGAGCTTTCTTAGATGATACAAACACAAAGCGATATCCTGACAATAAA
HC2F	-----
HC2A	GTGAAGCTGCTTAAGGAAGTTTTTCAGGCAATTTGTGGAAGCTTGCGGTCAAGCCTTAGCG
HC2-80	GTGAAGCTGCTTAAGGAAGTTTTTCAGGCAATTTGTGGAAGCTTGCGGTCAAGCCTTAGCG
HC2B	GTGAAGCTGCTTAAGGAAGTTTTTCAGGCAATTTGTGGAAGCTTGCGGTCAAGCCTTAGCG
HC2C	GTGAAGCTGCTTAAGGAAGTTTTTCAGGCAATTTGTGGAAGCTTGCGGTCAAGCCTTAGCG
HC2D-KIAA1058	GTGAAGCTGCTTAAGGAAGTTTTTCAGGCAATTTGTGGAAGCTTGCGGTCAAGCCTTAGCG
HC2E	GTGAAGCTGCTTAAGGAAGTTTTTCAGGCAATTTGTGGAAGCTTGCGGTCAAGCCTTAGCG
HC2F	-----
HC2A	GTAAACGAACGCTCTGATTAAAGAAGACCAGCTCGAGTATCAGGAAGAAATGAAAGCCAAC
HC2-80	GTAAACGAACGCTCTGATTAAAGAAGACCAGCTCGAGTATCAGGAAGAAATGAAAGCCAAC
HC2B	GTAAACGAACGCTCTGATTAAAGAAGACCAGCTCGAGTATCAGGAAGAAATGAAAGCCAAC
HC2C	GTAAACGAACGCTCTGATTAAAGAAGACCAGCTCGAGTATCAGGAAGAAATGAAAGCCAAC
HC2D-KIAA1058	GTAAACGAACGCTCTGATTAAAGAAGACCAGCTCGAGTATCAGGAAGAAATGAAAGCCAAC
HC2E	GTAAACGAACGCTCTGATTAAAGAAGACCAGCTCGAGTATCAGGAAGAAATGAAAGCCAAC
HC2F	-----

FIG. 3A (cont.)

HC2A	TACAGGGAAATGGCGAAGGAGCTTTCTGAAATCATGCATGAGCAG-----ATCTGCC
HC2-80	TACAGGGAAATGGCGAAGGAGCTTTCTGAAATCATGCATGAGCAG-----ATCTGCC
HC2B	TACAGGGAAATGGCGAAGGAGCTTTCTGAAATCATGCATGAGCAG-----ATCTGCC
HC2C	TACAGGGAAATGGCGAAGGAGCTTTCTGAAATCATGCATGAGCAG-----ATCTGCC
HC2D-KIAA1058	TACAGGGAAATGGCGAAGGAGCTTTCTGAAATCATGCATGAGCAGCTGGGATGATCTGCC
HC2E	TACAGGGAAATGGCGAAGGAGCTTTCTGAAATCATGCATGAGCAG-----ATCTGCC
HC2F	-----
HC2A	CCCTGGAGGAGAAGACGAGCGTCTTACCGAATTCCCTTCACATCTTCAACGCCATCAGTG
HC2-80	CCCTGGAGGAGAAGACGAGCGTCTTACCGAATTCCCTTCACATCTTCAACGCCATCAGTG
HC2B	CCCTGGAGGAGAAGACGAGCGTCTTACCGAATTCCCTTCACATCTTCAACGCCATCAGTG
HC2C	CCCTGGAGGAGAAGACGAGCGTCTTACCGAATTCCCTTCACATCTTCAACGCCATCAGTG
HC2D-KIAA1058	CCCTGGAGGAGAAGACGAGCGTCTTACCGAATTCCCTTCACATCTTCAACGCCATCAGTG
HC2E	CCCTGGAGGAGAAGACGAGCGTCTTACCGAATTCCCTTCACATCTTCAACGCCATCAGTG
HC2F	-----
HC2A	GGACTCCAACAAGCACAATGGTTACGGGATGACCAGCTCGTCTTCGGTCGTGTGATTAC
HC2-80	GGACTCCAACAAGCACAATGGTTACGGGATGACCAGCTCGTCTTCGGTCGTGTGATTAC
HC2B	GGACTCCAACAAGCACAATGGTTACGGGATGACCAGCTCGTCTTCGGTCGTGTGA----
HC2C	GGACTCCAACAAGCACAATGGTTACGGGATGACCAGCTCGTCTTCGGTCGTGTGA----
HC2D-KIAA1058	GGACTCCAACAAGCACAATGGTTACGGGATGACCAGCTCGTCTTCGGTCGTGTGATTAC
HC2E	GGACTCCAACAAGCACAATGGTTACGGGATGACCAGCTCGTCTTCGGTCGTGTGA----
HC2F	-----
HC2A	ATCTCATGGCCCGTGTGTGGGGACTTGCTTTGTCATTTGCAAACCTCAGGATGCTTTCCAA
HC2-80	ATCTCATGGCCCGTGTGTGGGGACTTGCTTTGTCATTTGCAAACCTCAGGATGCTTTCCAA
HC2B	-----
HC2C	-----
HC2D-KIAA1058	ATCTCATGGCCCGTGTGTGGGGACTTGCTTTGTCATTTGCAAACCTCAGGATGCTTTCCAA
HC2E	-----
HC2F	-----
HC2A	AGCCAATCACTGGGGAGACCGAGCACAGGGAGGACCAAGGGGAAGGGGAGAGAAAGGAAA
HC2-80	AGCCAATCACTGGGGAGACCGAGCACAGGGAGGACCAAGGGGAAGGGGAGAGAAAGGAAA
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AGCCAATCACTGGGGAGACCGAGCACAGGGAGGACCA--GGGGAAGGGGAGAGAAAGGAAA
HC2E	-----
HC2F	-----
HC2A	TAAAGAACAACGTTATTTCTTAACAGACTTTCTATAGGAGTTGTAAGAAGGTGCACATAT
HC2-80	TAAAGAACAACGTTATTTCTTAACAGACTTTCTATAGGAGTTGTAAGAAGGTGCACATAT
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TAAAGAACAACGTTATTTCTTAACAGACTTTCTATAGGAGTTGTAAGAAGGTGCACATAT
HC2E	-----
HC2F	-----

FIG. 3A (cont.)

HC2A	TTTTTTAAATCTCACTGGCAATATTCAAAGTTTTCATTGTGTCTTAACAAAGGTGTGGTA
HC2-80	TTTTTTAAATCTCACTGGCAATATTCAAAGTTTTCATTGTGTCTTAACAAAGGTGTGGTA
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TTTTTTAAATCTCACTGGCAATATTCAAAGTTTTCATTGTGTCTTAACAAAGGTGTGGTA
HC2E	-----
HC2F	-----
HC2A	GACACTCTTGAGCTGGACTTAGATTTTATTCTTCCTTGCAGAGTAGTGTTAGAATAGATG
HC2-80	GACACTCTTGAGCTGGACTTAGATTTTATTCTTCCTTGCAGAGTAGTGTTAGAATAGATG
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GACACTCTTGAGCTGGACTTAGATTTTATTCTTCCTTGCAGAGTAGTGTTAGAATAGATG
HC2E	-----
HC2F	-----
HC2A	GCCTACAGAAAAAAGGTTCTGGGATCTACATGGCAGGGAGGGCTGCACTGACATTGAT
HC2-80	GCCTACAGAAAAAAGGTTCTGGGATCTACATGGCAGGGAGGGCTGCACTGACATTGAT
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GCCTACAGAAAAAAGGTTCTGGGATCTACATGGCAGGGAGGGCTGCACTGACATTGAT
HC2E	-----
HC2F	-----
HC2A	GCCTGGGGGACCTTTTGCCTCGACTCGTGCCGAAATCTGATCGTAATCAGGGTACAGAA
HC2-80	GCCTGGGGGACCTTTTGCCTCGACTCGTGCCGAAATCTGATCGTAATCAGGGTACAGAA
HC2B	-----
HC2C	-----
HC2D-KIAA1058	GCCTGGGGGACCTTTTGCCTCGAGGCTGAGCTGGAAATCTTGAAAATATTTTTTT---T
HC2E	-----
HC2F	-----
HC2A	CTTACTAGTTTGTCTAGGAGTATGTTGTATGACTAGGATTTGTGCTATTATCTCATTCA
HC2-80	CTTACTAGTTTGTCTAGGAGTATGTTGTATGACTAGGATTTGTGCTATTATCTCATTCA
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TTTCCTGTGGCACATTCAGGTTGAATACAAGAACTATTTTGTGACTAGTTTTTGATGAC
HC2E	-----
HC2F	-----
HC2A	ACAACATAGAGCAAGAATAGTGAGCTAACTGAGCTAGACACTCAATTAATCCGCTACTGG
HC2-80	ACAACATAGAGCAAGAATAGTGAGCTAACTGAGCTAGACACTCAATTAATCCGCTACTGG
HC2B	-----
HC2C	-----
HC2D-KIAA1058	CTAAGGGAAGTGAACATTGTAATTTTGTACCAGTGAACCAGGAGATTTAGTGCTTTTAT
HC2E	-----
HC2F	-----

FIG. 3A (cont.)

HC2A	CTTCAAGTCAGAACTTTGTCATTAATCATCGACTCCGGGACGGTCATATATGTATTACAT
HC2-80	CTTCAAGTCAGAACTTTGTCATTAATCATCGACTCCGGGACGGTCATATATGTATTACAT
HC2B	-----
HC2C	-----
HC2D-KIAA1058	ATTCATTTTCCTTGCAATTTAAGAAAATATGAAAGCTTAAGGAATTATGTGAGCTTAAACT
HC2E	-----
HC2F	-----
HC2A	TTCTACATTTTTTAATACTCACATGGGCTTATGCATTAAGTTTAATTGTGATAAAATTTGTG
HC2-80	TTCTACATTTTTTAATACTCACATGGGCTTATGCATTAAGTTTAATTGTGATAAAATTTGTG
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AGTCAAGCAGTTTtagaaccAAAGGCCTATATTAATAACCGCAACTATGCTGAAAAGTACA
HC2E	-----
HC2F	-----
HC2A	CTGGTCCAGTATATGCAATACACTTTAATGGTTTATTCTTGTCATAAAAAATGTGCAATAT
HC2-80	CTGGTCCAGTATATGCAATACACTTTAATGGTTTATTCTTGTCATAAAAAATGTGCAATAT
HC2B	-----
HC2C	-----
HC2D-KIAA1058	AAGTAGTACAGTATATTGTTATGTACATATCATTGTTAATACAGTCCGGCATTCTGTAC
HC2E	-----
HC2F	-----
HC2A	GGAGATGTATACAAGTCTTTACT-----
HC2-80	GGAGATGTATACAAGTCTTTACT-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	ATATATGTATTACATTTCTACATTTTTTAATACTCACATGGGCTTATGCATTAAGTTTAAT
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TGTGATAAAATTTGTGCTGTTCCAGTATATGCAATACACTTTAATGTTTATTCTTGTACA
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TAAAAATGTGCAATATGGAGATGTATACAGTCTTTACTATATTAGGTTTATAAACAGTTT
HC2E	-----
HC2F	-----

FIG. 3A (cont.)

HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TAAGAATTCATCCTTTGCCAAAATGGTGGAGTATGTAATTGGTAAATCATAAATCCTG
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TGGTGAATGGTGGTGTACTTTAAAGCTGTCACCATGTTATATTTCTTTTAAGACATTAA
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	TTTAGTAATTTTATATTTGGGAAAAATAAAGGTTTTTAATTTTATTTAACTGGAATCACTG
HC2E	-----
HC2F	-----
HC2A	-----
HC2-80	-----
HC2B	-----
HC2C	-----
HC2D-KIAA1058	CCCTGCTGTAATTAAACATTCTGTACCACATCTGTATTAAAAAGACATTGCTGACC
HC2E	-----
HC2F	-----

FIG. 3A (cont.)

HC2A	-----
HC2A-80	-----
HC2B	-----
HC2C	-----
HC2D	ASGNLDKNARFS AIYRQDSNKL SNDDMLKLLADFRKPEKMAKLPVILGNLDITIDNVSSD
HC2E	-----
HC2F	-----
HC2A	-----
HC2A-80	-----
HC2B	-----
HC2C	-----
HC2D	FPNYVNSSYIPTKQFETCSKTPITFEVEEFVPCIPKHTQPYTIYTNHLYVYPKYLYDSQ
HC2E	-----
HC2F	-----
HC2A	-----VLHHHQNP E FYDEIK
HC2A-80	-----
HC2B	-----
HC2C	-----
HC2D	KSFAKARNIAICIEFKDSDEEDSQPLKCIYGRPGGPVFTRSAFAAVLHHHQNP E FYDEIK
HC2E	-----
HC2F	-----
HC2A	IELPTQLHEKHHL L L TFFHVSCDNSSKGSTKKRDV VETQVGYSWLP L L K DGRVVTSEQHI
HC2A-80	-----
HC2B	-----
HC2C	-----
HC2D	IELPTQLHEKHHL L L TFFHVSCDNSSKGSTKKRDV VETQVGYSWLP L L K DGRVVTSEQHI
HC2E	-----
HC2F	-----
HC2A	PVSANLPSGYLGYQELGMGRHYGPEIKWVDGGKPLLKISTHLVSTVYTQDQHLHNFFQYC
HC2A-80	-----
HC2B	-----
HC2C	-----
HC2D	PVSANLPSGYLGYQELGMGRHYGPEIKWVDGGKPLLKISTHLVSTVYTQDQHLHNFFQYC
HC2E	-----
HC2F	-----
HC2A	QKTESGAQALGNELVKYLSLHAMEGHVMI AFLPTILNQLFRVLTRATQEEVAVNVTRVI
HC2A-80	-----
HC2B	-----AMEGHVMI AFLPTILNQLFRVLTRATQEEVAVNVTRVI
HC2C	-----
HC2D	QKTESGAQALGNELVKYLSLHAMEGHVMI AFLPTILNQLFRVLTRATQEEVAVNVTRVI
HC2E	-----AMEGHVMI AFLPTILNQLFRVLTRATQEEVAVNVTRVI
HC2F	-----

Fig. 3B

HC2A	IHVVAQCHEEGLESHLRSYVKYAYKAEPYVASEYKTVHEELTKSMTTILKPSADFLTSNK
HC2A-80	-----
HC2B	IHVVAQCHEEGLESHLRSYVKYAYKAEPYVASEYKTVHEELTKSMTTILKPSADFLTSNK
HC2C	-----
HC2D	IHVVAQCHEEGLESHLRSYVKYAYKAEPYVASEYKTVHEELTKSMTTILKPSADFLTSNK
HC2E	IHVVAQCHEEGLESHLRSYVKYAYKAEPYVASEYKTVHEELTKSMTTILKPSADFLTSNK
HC2F	-----
HC2A	LLRYSWFFFDVLIKSMAQHLENSKVKLLRNQRFPA SYHHAETV VNMMLPHITQKFGDN
HC2A-80	-----
HC2B	LLRYSWFFFDVLIKSMAQHLENSKVKLLRNQRFPA SYHHAETV VNMMLPHITQKFGDN
HC2C	-----
HC2D	LLKYSWFFFDVLIKSMAQHLENSKVKLLRNQRFPA SYHHAETV VNMMLPHITQKFRDN
HC2E	LLRYSWFFFDVLIKSMAQHLENSKVKLLRNQRFPA SYHHAETV VNMMLPHITQKFGDN
HC2F	-----
HC2A	PEASKNANHSLAVFIKRCFTFMDRGFVFKQINNYISCFAPGDPKTLFEYKFEFLRVVCNH
HC2A-80	-----
HC2B	PEASKNANHSLAVFIKRCFTFMDRGFVFKQINNYISCFAPGDPKTLFEYKFEFLRVVCNH
HC2C	-----
HC2D	PEASKNANHSLAVFIKRCFTFMDRGFVFKQINNYISCFAPGDPKTLFEYKFEFLRVVCNH
HC2E	PEASKNANHSLAVFIKRCFTFMDRGFVFKQINNYISCFAPGDPKTLFEYKFEFLRVVCNH
HC2F	-----
HC2A	EHYIPLNLPMPFGKGRIQRYQDLQLDYSLTDEFCRNHFLVGLLLREVGTALQEFREVRLI
HC2A-80	-----QLDYSLTDEFCRNHFLVGLLLREVGTALQEFREVRLI
HC2B	EHYIPLNLPMPFGKGRIQRYQDLQLDYSLTDEFCRNHFLVGLLLREVGTALQEFREVRLI
HC2C	-----
HC2D	EHYIPLNLPMPFGKGRIQRYQDLQLDYSLTDEFCRNHFLVGLLLREVGTALQEFREVRLI
HC2E	EHYIPLNLPMPFGKGRIQRYQDLQLDYSLTDEFCRNHFLVGLLLREVGTALQEFREVRLI
HC2F	-----
HC2A	AISVLKNLLIKHSFDDRYASRSHQARIATLYLPLFGLLIENVQRINVRDVSPFPVNAGMT
HC2A-80	AISVLKNLLIKHSFDDRYASRSHQARIATLYLPLFGLLIENVQRINVRDVSPFPVNAGMT
HC2B	AISVLKNLLIKHSFDDRYASRSHQARIATLYLPLFGLLIENVQRINVRDVSPFPVNAGMT
HC2C	-----
HC2D	AISVLKNLLIKHSFDDRYASRSHQARIATLYLPLFGLLIENVQRINVRDVSPFPVNAGMT
HC2E	AISVLKNLLIKHSFDDRYASRSHQARIATLYLPLFGLLIENVQRINVRDVSPFPVNAGMT
HC2F	-----
HC2A	VKDESLALPAVNPLVTPQKGSTLDNSLHKDLLGAISGIASPYTTSTPNINSVRNADSRGS
HC2A-80	VKDESLALPAVNPLVTPQKGSTLDNSLHKDLLGAISGIASPYTTSTPNINSVRNADSRGS
HC2B	VKDESLALPAVNPLVTPQKGSTLDNSLHKDLLGAISGIASPYTTSTPNINSVRNADSRGS
HC2C	-----
HC2D	VKDESLALPAVNPLVTPQKGSTLDNSLHKDLLGAISGIASPYTTSTPNINSVRNADSRGS
HC2E	VKDESLALPAVNPLVTPQKGSTLDNSLHKDLLGAISGIASPYTTSTPNINSVRNADSRGS
HC2F	-----ADSRGS

FIG. 3B (cont.)

HC2A	LISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSEIKSLLMCFLYILKSMS
HC2A-80	LISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSEIKSLLMCFLYILKSMS
HC2B	LISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSEIKSLLMCFLYILKSMS
HC2C	-----
HC2D	LISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSEIKSLLMCFLYILKSMS
HC2E	LISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSEIKSLLMCFLYILKSMS
HC2F	LISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSEIKSLLMCFLYILKSMS
HC2A	DDALFTYWNKASTSELMDDFTTISEVCLHQFQYMGKRYIARNQEGLGPIVHDRKSQTLFVS
HC2A-80	DDALFTYWNKASTSELMDDFTTISEVCLHQFQYMGKRYIARNQEGLGPIVHDRKSQTLFVS
HC2B	DDALFTYWNKASTSELMDDFTTISEVCLHQFQYMGKRYIARNQEGLGPIVHDRKSQTLFVS
HC2C	-----
HC2D	DDALFTYWNKASTSELMDDFTTISEVCLHQFQYMGKRYIAR-----
HC2E	DDALFTYWNKASTSELMDDFTTISEVCLHQFQYMGKRYIARNQEGLGPIVHDRKSQTLFVS
HC2F	DDALFTYWNKASTSELMDDFTTISEVCLHQFQYMGKRYIAS-----VR--KISSVLGIS
HC2A	RNRTGMMHARLQQLGSLDNSLTFNHSYGHSDADVLHQSLLEANIATEVCLTALDTLSLFT
HC2A-80	RNRTGMMHARLQQLGSLDNSLTFNHSYGHSDADVLHQSLLEANIATEVCLTALDTLSLFT
HC2B	RNRTGMMHARLQQLGSLDNSLTFNHSYGHSDADVLHQSLLEANIATEVCLTALDTLSLFT
HC2C	-----
HC2D	---TGMHARLQQLGSLDNSLTFNHSYGHSDADVLHQSLLEANIATEVCLTALDTLSLFT
HC2E	RNRTGMMHARLQQLGSLDNSLTFNHSYGHSDADVLHQSLLEANIATEVCLTALDTLSLFT
HC2F	V-----D-NG-----YGHSDADVLHQSLLEANIATEVCLTALDTLSLFT
HC2A	LAFKNQLLADHGHNPMLKKVFDVYLCFLQKHQSETALKNVFTALRSLIYKFPSTFYEGRA
HC2A-80	LAFKNQLLADHGHNPMLKKVFDVYLCFLQKHQSETALKNVFTALRSLIYKFPSTFYEGRA
HC2B	LAFK--LLADHGHNPMLKKVFDVYLCFLQKHQSETALKNVFTALRSLIYKFPSTFYEGRA
HC2C	-----
HC2D	LAFKNQLLADHGHNPMLKKVFDVYLCFLQKHQSETALKNVFTALRSLIYKFPSTFYEGRA
HC2E	LAFKNQLLADHGHNPMLKKVFDVYLCFLQKHQSETALKNVFTALRSLIYKFPSTFYEGRA
HC2F	LAFKNQLLADHGHNPMLKKK-----
HC2A	DMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKSFVRTHLQVII SVSQLI
HC2A-80	DMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKSFVRTHLQVII SVSQLI
HC2B	DMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKSFVRTHLQVII SVSQLI
HC2C	-----
HC2D	DMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKSFVRTHLQVII SVSQLI
HC2E	DMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKSFVRTHLQVII SVSQLI
HC2F	-----
HC2A	ADVVGIGETRFOQSLSIINNCANSRDLIKHTSFSSDVKDLTKRIRTVLMATAQMKEHEND
HC2A-80	ADVVGIGETRFOQSLSIINNCANSRDLIKHTSFSSDVKDLTKRIRTVLMATAQMKEHEND
HC2B	ADVVGIGETRFOQSLSIINNCANSRDLIKHTSFSSDVKDLTKRIRTVLMATAQMKEHEND
HC2C	-----
HC2D	ADVVGIGETRFOQSLSIINNCANSRDLIKHTSFSSDVKDLTKRIRTVLMATAQMKEHEND
HC2E	ADVVGIGETRFOQSLSIINNCANSRDLIKHTSFSSDVKDLTKRIRTVLMATAQMKEHEND
HC2F	-----

FIG. 3B (cont.)

HC2A	PEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGDLSEAAMCYVHVTALVAEYLTRK
HC2A-80	PEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGDLSEAAMCYVHVTALVAEYLTRK
HC2B	PEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGDLSEAAMCYVHVTALVAEYLTRK
HC2C	-----
HC2D	PEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGDLSEAAMCYVHVTALVAEYLTRK
HC2E	PEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGDLSEAAMCYVHVTALVAEYLTRK
HC2F	-----
HC2A	-----GVFRQGCTAFRVITPNIDEEASMMEDVGMQDVHFNE
HC2A-80	-----GVFRQGCTAFRVITPNIDEEASMMEDVGMQDVHFNE
HC2B	-----GVFRQGCTAFRVITPNIDEEASMMEDVGMQDVHFNE
HC2C	-----FRQGCTAFRVITPNIDEEASMMEDVGMQDVHFNE
HC2D	EAVQWEPPLLPHSHSACLRRSRGGVFRQGCTAFRVITPNIDEEASMMEDVGMQDVHFNE
HC2E	-----GVFRQGCTAFRVITPNIDEEASMMEDVG-----
HC2F	-----
HC2A	DVLMELLEQCADGLWKAERYELIADIYKLIPIYEKRR-----
HC2A-80	DVLMELLEQCADGLWKAERYELIADIYKLIPIYEKRR-----
HC2B	DVLMELLEQCADGLWKAERYELIADIYKLIPIYEKRRDFERLAHLYDTLHRAYSK
HC2C	DVLMELLEQCADGLWKAERYELIADIYKLIPIYEKRRDFERLAHLYDTLHRAYSK
HC2D	DVLMELLEQCADGLWKAERYELIADIYKLIPIYEKRRDFERLAHLYDTLHRAYSK
HC2E	-----KAERYELIADIYKLIPIYEKRRDFERLAHLYDTLHRAYSK
HC2F	-----
HC2A	-----DFFEDEDGKEYIYKEPKLTPLSE
HC2A-80	-----DFFEDEDGKEYIYKEPKLTPLSE
HC2B	VTEVMHSGRRLLGTYFRVAFFG-----QGFFEDEDGKEYIYKEPKLTPLSE
HC2C	VTEVMHSGRRLLGTYFRVAFFG-----QGFFEDEDGKEYIYKEPKLTPLSE
HC2D	VTEVMHSGRRLLGTYFRVAFFGQAAQYQFTDSETDVEGFFEDEDGKEYIYKEPKLTPLSE
HC2E	VTEVMHSGRRLLGTYFRVAFFG-----QGFFEDEDGKEYIYKEPKLTPLSE
HC2F	-----
HC2A	ISQRLLKLYSDKFGSENVKMIQDSGKVNPKDLDSKYAYIQVTHVIPFFDEKELQERKTEF
HC2A-80	ISQRLLKLYSDKFGSENVKMIQDSGKVNPKDLDSKYAYIQVTHVIPFFDEKELQERKTEF
HC2B	ISQRLLKLYSDKFGSENVKMIQDSGKVNPKDLDSKYAYIQVTHVIPFFDEKELQERKTEF
HC2C	ISQRLLKLYSDKFGSENVKMTQDSGKVNPKDLDSKYAYIQVTHVIPFFDEKELQERKTEF
HC2D	ISQRLLKLYSDKFGSENVKMIQDSGKVNPKDLDSKYAYIQVTHVIPFFDEKELQERKTEF
HC2E	ISQRLLKLYSDKFGSENVKMIQDSGKVNPKDLDSKYAYIQVTHVIPFFDEKELQERKTEF
HC2F	-----
HC2A	ERSHNIRRFMFEMPFTQTGKRQGGVEEQCKRRTILTAIHCFPYVKKRIPVMYQHHTDLNP
HC2A-80	ERSHNIRRFMFEMPFTQTGKRQGGVEEQCKRRTILTAIHCFPYVKKRIPVMYQHHTDLNP
HC2B	ERSHNIRRFMFEMPFTQTGKRQGGVEEQCKRRTILTAIHCFPYVKKRIPVMYQHHTDLNP
HC2C	ERSHNIRRFMFEMPFTQTGKRQGGVEEQCKRRTILTAIHCFPYVKKRIPVMYQHHTDLNP
HC2D	ERSHNIRRFMFEMPFTQTGKRQGGVEEQCKRRTILTAIHCFPYVKKRIPVMYQHHTDLNP
HC2E	ERSHNIRRFMFEMPFTQTGKRQGGVEEQCKRRTILTAIHCFPYVKKRIPVMYQHHTDLNP
HC2F	-----

FIG. 3B (cont.)

HC2A	IEVAIDEMSKKVAELRQLCSSAEVDMIKLQLKLQGSVSVQVNAGPLAYARAFLLDDTNTKR
HC2A-80	IEVAIDEMSKKVAELRQLCSSAEVDMIKLQLKLQGSVSVQVNAGPLAYARAFLLDDTNTKR
HC2B	IEVAIDEMSKKVAELRQLCSSAEVDMIKLQLKLQGSVSVQVNAGPLAYARAFLLDDTNTKR
HC2C	IEVHZ-----
HC2D	IEVAIDEMSKKVAELRQLCSSAEVDMIKLQLKLQGSVSVQVNAGPLAYARAFLLDDTNTKR
HC2E	IEVAIDEMSKKVAELRQLCSSAEVDMIKLQLKLQGSVSVQVNAGPLAYARAFLLDDTNTKR
HC2F	-----
HC2A	YPDKNVKLLKEVFRQFVEACGQALAVNERLIKEDQLEYQEEMKANYREMAKELSEIMHEQ
HC2A-80	YPDKNVKLLKEVFRQFVEACGQALAVNERLIKEDQLEYQEEMKANYREMAKELSEIMHEQ
HC2B	YPDKNVKLLKEVFRQFVEACGQALAVNERLIKEDQLEYQEEMKANYREMAKELSEIMHEQ
HC2C	-----
HC2D	YPDKNVKLLKEVFRQFVEACGQALAVNERLIKEDQLEYQEEMKANYREMAKELSEIMHEQ
HC2E	YPDKNVKLLKEVFRQFVEACGQALAVNERLIKEDQLEYQEEMKANYREMAKELSEIMHEQ
HC2F	-----
HC2A	ICPLEEKTSVLPNSLHIFNAISGTPTSTMVHGMTSSSSSVVZ----
HC2A-80	ICPLEEKTSVLPNSLHIFNAISGTPTSTMVHGMTSSSSSVVZ----
HC2B	ICPLEEKTSVLPNSLHIFNAISGTPTSTMVHGMTSSSSSVVZ----
HC2C	-----
HC2D	LG-----
HC2E	ICPLEEKTSVLPNSLHIFNAISGTPTSTMVHGMTSSSSSVVZ----
HC2F	-----

FIG. 3B (cont.)




FIG. 4A




FIG. 4B

HC2A	-----
KIAA	ASGNLDKNARFSATYRQDSNKLSDNDMLKLLADFRKPEKMAKLPVILGNLDITIDNVSSD
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	-----
KIAA	FPNYVNSSYIPTKQFETCSKTPITFEVEEFVPCIPKHTQPYTIYTNHLYVYPKYLKYDSQ
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	-----VLHHHQNPFEYDEIK
KIAA	KSFAKARNIAICIEFKDSDEEDSQPLKCIYGRPGGPVFTRSAFAAVLHHHQNPFEYDEIK
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	IELPTQLHEKHHLTFFHVSCDNSSKSGSTKKRDVVEQVGYSWLPLLKDGRVVTSEQHI
KIAA	IELPTQLHEKHHLTFFHVSCDNSSKSGSTKKRDVVEQVGYSWLPLLKDGRVVTSEQHI
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	PVSANLPSGYLGYQELGMGRHYGPEIKWVDGGKPLLKISTHLVSTVYTQDQHLNFFQYC
KIAA	PVSANLPSGYLGYQELGMGRHYGPEIKWVDGGKPLLKISTHLVSTVYTQDQHLNFFQYC
rat	-----
HC4	-----
HC1	-----
HC3	-----GPGPARSTVSIISLISNSARV
HC5	-----
HC2A	OKTESGAQALGNEIVKYLKSLHAMEGHVMIAFLPTILNQLFRVLT-RATQEEVAVNVTRV
KIAA	OKTESGAQALGNEIVKYLKSLHAMEGHVMIAFLPTILNQLFRVLT-RATQEEVAVNVTRV
rat	-----
HC4	-----MEIQVLIRFLSVILMQLFWVLPNMIHEDDVPISCPMV
HC1	-----MSFLPIILNQLFKVLV-QNEEDEITTTVTRV
HC3	NRSRSLSNSNPDISGTPTSPDDEVRSIIGSKGLDRSNSWNTGGPKAAPWGSNPSPSAES
HC5	-----

FIG.5A

HC2A	I IHVVAQCHEEGLESHLRSYVKYAYKAEPYVASEYKTVHEELTKSMTTILKPSADFLTSN
KIAA	I IHVVAQCHEEGLESHLRSYVKYAYKAEPYVASEYKTVHEELTKSMTTILKPSADFLTSN
rat	-----
HC4	LFHIVSKCHEEGLD SYLSSFIKYSFRPGKPSAPQAPLIHETLATMMIALLLKQSADFLAIN
HC1	LPDIVAKCHEEQLDHSVQSYIKFVKTR---ACKERPVEDHLAKNVTGLLK-SNDSPTVK
HC3	TQAMDRSCNRMSSHTETSSFLQTLTGRLP---TKKLFHEELALQWVCSG--SVR---E
HC5	-----
	Cadherin Cleavage
HC2A	KLLRYSWFFFDVLIKSMAQHLENSKVKLIRNQRFPASYHHAAETVVNMLMPHITQKFGD
KIAA	KLLKYSWFFFDVLIKSMAQHLENSKVKLIRNQRFPASYHHAAETVVNMLMPHITQKFRD
rat	-----
HC4	KLLKYSWFFFEIIAKSMATYLLLEENKIKLTHGQRFPKAYHHALHSFLAIT-IVESQYAE
HC1	HVLKHSWFFFAIILKSMAQHLEIDTNKIQLERPQRFPESYQNELDNLMVLSHDHVIWKYKD
HC3	SALQQAWFFFEILMVKSMVHHLYFNDKLEAERKSREPERFMDIDIAALVSTIASDIVSRFQK
HC5	-----
HC2A	NPEASKNANHSLAVFIKRCFTFMDRGFVFKQIN---NYIS--CFAPGDPKTLFEYKFEFL
KIAA	NPEASKNANHSLAVFIKRCFTFMDRGFVFKQIN---NYIS--CFAPGDPKTLFEYKFEFL
rat	-----
HC4	IPKESRNVNYSLASFLKCCLTLMDRGFVFNLIN---DYIS--GFSPKDPKVLAEYKFEFL
HC1	ALEETRRATHSVAREFLKRCFTFMDRGCVFKMVN---NYIS--MFSSGDLKTLQYKFDL
HC3	DTEMVERLNTSLAFFLNDLLSVMDRGFVFSLIKSCYQVSSKLYSLPNPSVLVSLRLDFL
HC5	-----
HC2A	RVVCNHEHYIPLNLPM-----PFGKGRIQR-----YQDLQL---DYSLTDEF
KIAA	RVVCNHEHYIPLNLPM-----PFGKGRIQR-----YQDLQL---DYSLTDEF
rat	-----
HC4	QTICNHEHYIPLNLPM-----AFAKPKLQR-----VQDSNL---EYSLSDEY
HC1	QEVQCHEHFIPCLPIRSANIPDPLTPSES-----TQELHASDMPEYSVTNEF
HC3	RIICSHHYVTNLNPLPCSLTPPASPSVSSATSQSSGFSSTNVQDQKIANMFELS--VPF
HC5	-----MNADTAPTSPCPSIS---SQNSSSCSSFQDQKIASMFDRTSRVPA
HC2A	CRNHFLVGLLLREVGTALQEFRE---VRLIAISVLKNLLIKHSFDDRYASRSHQARIAT
KIAA	CRNHFLVGLLLREVGTALQEFRE---VRLIAISVLKNLLIKHSFDDRYASRSHQARIAT
rat	-----
HC4	CKHHFLVGLLLRETSIALQDNYE---IRYTAISVIKNLLIKHAFDTRYQHKNQQAQIAQ
HC1	CRKHFLIGILLREVGFALQEDQD---VRHLALAVLKNLMAKHSFDDRYREPRKQAQIAS
HC3	RQHYLAGLVLTALVLDPAEGLFGLHKKVINMVHNLSSHDSDPRYSDPQIKARVAM
HC5	SSTS-SPGLLFTLAAALDAEGEGISEVQRKAVSAIHSLLSSHDLDPRCVKPEVKVKIAA
HC2A	LYLPLFGLLIENVQRINVRDVSPFPVNAG-MTVKDESIALPAVNPLVTPQKGSTLDNSLH
KIAA	LYLPLFGLLIENVQRINVRDVSPFPVNAG-MTVKDESIALPAVNPLVTPQKGSTLDNSLH
rat	-----
HC4	LYLPPVGLLLENIQRLAGRTLYSCAAMPNSASRDEFPCG-----FTSP--AN--RGSLS
HC1	LYMPLYGMLLDNMPRIYLDLYPFTVNTSNQGSRDDLSNNGGFQSQTAKHANSVDTSFS
HC3	LYLPLIGIIMETVPQLYDFTETHNQRGRPICIAATDDYSE-----SG---SMIS
HC5	LYLPLVGIIILDALPQLCDFTVADTRRYR---TSGSDEEQE-----GA---GAIT
HC2A	KDLLGAISGIASPYTTSTPNINSVRNADSRGSLISTDSGNLSPERNSEKSNSLDKHQSS
KIAA	KDLLGAISGIASPYTTSTPNINSVRNADSRGSLISTDSGNLSPERNSEKSNSLDKHQSS
rat	-----
HC4	TDKDTAYGSGFQNG-----HGKREDSRGSLIP-EGATGFPDQGNTEG-----TRQS
HC1	KDVLNSIAAFSS-----IAISTVNHADSRASLASLDSNPSTNEKSSEKTDNCEKIPRL
HC3	QTVAMAIAGTSVPQ-----LTRPGSFLLTSTSGRQHT-----
HC5	QNVALAIAAGNFN-----LKTSG-IVLSSSLPYKQYN-----

FIG. 5A (cont.)

HC2A	TLGNSVVRCDKLDQSEIKSLLMCFLYILKSMSDDALFTYWN-KASTSEIMDFFTISEVCL
KIAA	TLGNSVVRCDKLDQSEIKSLLMCFLYILKSMSDDALFTYWN-KASTSEIMDFFTISEVCL
rat	-----
HC4	STRSSVSQYNRLDQYEIRSLLMCYLYIVKMI SEDTLLTYWN-KVSPQELINILILLEVCL
HC1	ALIGSTLRFDRLDQAETRSLIMCFLHIMKTI SYETLIAYWQ-RAPSPVEVSDFFSILDVCL
HC3	-----TFSAESSRSLLICLLWVLKN-ADETVLQKWFTDLSVLQNLRLDLLLYLCV
HC5	-----MLNADTTRNIMICFLWIMKN-ADQSLIRKWIADLPSTQLNRLDLLFICV
HC2A	HQFQYMGKRYIARNQEGLG--PIVHDRKS-----QTLPVSRNRTGMM
KIAA	HQFQYMGKRYIAR-----TGMM
rat	-----
HC4	FHFRYMGKRNIARVHDAWLSKHFGIDRKS-----QTMPALNRNSGVM
HC1	QNFYRLGKRNIIRKIAAAF--KFVQSTQNNGTILKGSNPSCQTSGLLAQWMHSTSRHEGHK
HC3	SCFEYKGGKVFERMNSLTFK--KSKDMRAK-----LEEAILGSIGARQEMV
HC5	LCFEYKGGKQSSDKVSTQVLQ--KSRDVKAR-----LEEALLRGEARGEMV
HC2A	HARLQQL-----GSLDNS-----LTFNHSYGHSDADVLHQSLLEANIATEVC
KIAA	HARLQQL-----GSLDNS-----LTFNHSYGHSDADVLHQSLLEANIATEVC
rat	-----
HC4	QARLQHL-----SSLESS-----FTLNHSSTTTEADIFHQALLEGNTATEVS
HC1	QHRSQTLPIIRGK---NALSNPKL----LQMLDNTMTSNSNEIDIVHHVDTEANIATEGC
HC3	RRSRGQLERSPSGSAFGSQENLRWRKDMTHWRQNTKLDKSRAEIEHEALIDGNLATEAN
HC5	RRRAPGNDRFP----GLNENLRWKKEQTHWRQANEKLDKTKAELDQEALISGNLATEAH
HC2A	LTALDTLSLFTLAFKNQLLADHGHNPIMKKVFDVYLCFLQKHQSE TALKNVFTALRSLIY
KIAA	LTALDTLSLFTLAFKNQLLADHGHNPIMKKVFDVYLCFLQKHQSE TALKNVFTALRSLIY
rat	-----KLSRGHSPIMKKVFDVYLCFLQKHQSE MALKNVFTALRSLIY
HC4	LTVLDTISFFTQCFKTHFLNNDGHNPIMKKVFDIHLAFLKNGQSEVSLKHVFASLRAFIS
HC1	LTILDVLSLFTQTHQRQLQCCDCQNSIMKRGFDTYMLFFQVQNQSATALKHVFASLRLFVC
HC3	LIILDTLEIVVQTVS--VTES--KESILGGVLKVLHSMACNQSAVYLQHC FATQRALVS
HC5	LIILDMQENIIQASS--ALDC--KDSILGGVLRVLVNSLNC DQSTYLTLCFATLRLAIA
HC2A	KFPSTFYEGRADMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNEFDYTGKKSFVRTH
KIAA	KFPSTFYEGRADMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNEFDYTGKKSFVRTH
rat	KFPSTFYEGRADMCASLCYEVLKCCNSKLSSIRTEASQLLYFLMRNNEFDYTGKKSFVRTH
HC4	KFPSAFFKGRVNMCAAFCEVLKCCTSKISSTRNEASALLYLLMRNNEFYTKRKTFLRTH
HC1	KFPSAFFQGPADLCGSFCYEVLKCCNHRSRSTQTEASALLYLFMRKNFEFNKQKSIVRSH
HC3	KFPELLFEEETEQCADLCRLRLRHCCSSIGTIRSHPSASLYLLMRQNF EIGN--NFARVK
HC5	KFGDLLFEEVEQCDFDLCHQVLHHCSSMDVTRSQCACATLYLLMRFSFGATS--NFARVK
HC2A	LQVIIISVSQLIADVVVGIGETRFQQSLSIINN CANSDRLIKHTSFSSDVKDLTKRIRTVLM
KIAA	LQVIIISVSQLIADVVVGIGETRFQQSLSIINN CANSDRLIKHTSFSSDVKDLTKRIRTVLM
rat	LQVIIISLSQLIADVVVGIGETRFQQSLSIINN CANSDRLIKHTSFSSDVKDLTKRIRTVLM
HC4	LQIIIAVSQLIADVALSGSRFQESLFIINN FANSRPLARAFPAEVKDLTKRIRTVLM
HC1	LQLIKAVSQLIAD-AGIGGSRFQHSLAITN NFANGDKQMN SNFPAEVKDLTKRIRTVLM
HC3	MQVPMSSLSSLVGTSQNFEEFLRRSLK TILTYAEEDLELRETTFFDQVQDLVFNLMILS
HC5	MQVTMSLASLVGRAPDFNEEHLRRSLRTILAYSEEDTAMQMPFPFTQVEELL CNLNSILY

FIG. 5A (cont.)

FIG. 5A (cont.)

		Coiled-Coil 1
HC2A	IHC	FPYVKKRIPVMYQHHTDINPIEVAIDEMSKKVAELRQLCSSAEVDMIKLQKLQGSV
KIAA	IHC	FPYVKKRIPVMYQHHTDINPIEVAIDEMSKKVAELRQLCSSAEVDMIKLQKLQGSV
rat	IHC	FPYVKKRIPVMYQHHTDINPIEVAIDEMSKKVAELHQLCSSAEVDMIKLQKLQGSV
HC4	SNS	FPYVKKRIPINCEQQINLKPIDGATDEIKDKTAELOKLCSSSTDVDMIQLQKLQGSV
HC1	SHL	FPYVKKRIQVISQSSTEINPIEVAIDEMSRKVSELNQLCTMEVDMISLQKLQGSV
HC3	SHAF	PYIKTRVNVTHKEEIIITPIEVAIEDMQKKTQELAFATHQDPADPKMLQKLQGSV
HC5	MHAF	PYIKTRISVIOKEEFVITPIEVAIEDMKKKTLQLAVAINQEPDAKMLQKLQGSV
		Coiled-Coil 2
HC2A	SVQ	VNAGPLAYARAFLDDTNTKRYPDNKVKLLKEVFRQFVEACGQALAVNERLIKEDQLE
KIAA	SVQ	VNAGPLAYARAFLDDTNTKRYPDNKVKLLKEVFRQFVEACGQALAVNERLIKEDQLE
rat	SVQ	VNAGPLAYARAFLDDTNTKRYPDNKVKLLKEVFRQFVEACGQALAVNERLIKEDQLE
HC4	SVQ	VNAGPLAYARAFLDNDQASKYPPKKVSELKDMFRKFIQACSIALLNERLIKEDQVE
HC1	SVK	VNAGPMAYARAFLEETNAKKYPDNQVKLLKEIFRQFADACGQALAVNERLIKEDQLE
HC3	GTT	VNQGPLEVAQVFLSEIPSDPKLFRHHNKLRLCFKDFTKRCEDALAKNKSLIGPVQKE
HC5	GAT	VNQGPLEVAQVFLAEIPADPKLYRHHNKLRLCFKEFIMRCGEAVEKNKRLITADQRE
		Coiled-Coil 2
HC2A	YQE	EMKANYREMAKELSEIMHEQICPLEEKTS-VLPNSLHIFNAISGTPSTMVHGMTSS
KIAA	YQE	EMKANYREMAKELSEIMHEQLG-----
rat	YQE	EMKANYREIRKELSDIIVPRICPGEDKRATKFFPAHLQRHQRTNKHSGSRVDQFILS
HC4	YHE	GLKSNFRDMVKELSDIIEHQILQEDTMHSPWMSNTLHVFCASISGSSDRGYGSPRYA
HC1	YQE	ELRSHYKDMSELSTVMNEQITGRDDLK---RGVDQCTTRVISKATPALPTVISISS
HC3	YQ	ELG----KLSS-----PZ-----
HC5	YQ	ELKKNYNKLENLRPMIERKIPELYKPIFRVESQKRDSFHRSSFRKCETQLSQGSZ-
		PBM
HC2A	SSV	V-----
KIAA	SSV	V-----
rat	CVT	LPHPHVGTCTCFVMCKLRTTFRANHWFCQAQEEAMGNGREKEPWTVIENSRFYRSWGK
HC4	EV	-----
HC1	SAE	V-----
HC3	SAE	V-----
HC5	SAE	V-----
HC2A	----	
KIAA	----	
rat	VHIF	F-----
HC4	----	
HC1	----	
HC3	----	
HC5	----	

FIG. 5A (cont.)




FIG. 5B

DOCK2=KIAA0209
DOCK3=KIAA0299
CLASP2variant=KIAA1058

1
A


2
 GTT TTA CAC CAT CAC CAA AAC CCA GAA TTT TAT GAT GAG ATT AAA ATA GAG TTG CCC ACT
 val leu his his his gln asn pro glu phe tyr asp glu ile lys ile glu leu pro thr
 32
 62
 CAG CTG CAT GAA AAG CAC CAC CTG TTG CTC ACA TTC TTC CAT GTC AGC TGT GAC AAC TCA
 gln leu his glu lys his his leu leu leu thr phe phe his val ser cys asp asn ser
 92
 122
 AGT AAA GGA AGC ACG AAG AAG AGG GAT GTC GTT GAA ACC CAA GTT GGC TAC TCC TGG CTT
 ser lys gly ser thr lys lys arg asp val val glu thr gln val gly tyr ser trp leu
 152
 182
 CCC CTC CTG AAA GAC GGA AGG GTG GTG ACA AGC GAG CAG CAC ATC CCG GTC TCG GCG AAC
 pro leu leu lys asp gly arg val val thr ser glu gln his ile pro val ser ala asn
 212
 242
 CTT CCT TCG GGC TAT CTT GGC TAC CAA GAG CTT GGG ATG GGC AGG CAT TAT GGT CCG GAA
 leu pro ser gly tyr leu gly tyr gln glu leu gly met gly arg his tyr gly pro glu
 272
 302
 ATT AAA TGG GTA GAT GGA GGC AAG CCA CTG CTG AAA ATT TCC ACT CAT CTG GTT TCT ACA
 ile lys trp val asp gly gly lys pro leu leu lys ile ser thr his leu val ser thr
 332
 362
 GTG TAT ACT CAG  GAT CAG CAT TTA CAT AAT TTT TTC CAG TAC TGT CAG AAA ACC GAA TCT
 val tyr thr gln asp gln his leu his asn phe phe gln tyr cys gln lys thr glu ser
 392
 422
 GGA GCC CAA GCC TTA GGA AAC GAA CTT GTA AAG TAC CTT AAG AGT CTG CAT GCG ATG GAA
 gly ala gln ala leu gly asn glu leu val lys tyr leu lys ser leu his ala met glu
 452
 482
 GGC CAC GTG ATG ATC GCC TTC TTG CCC ACT ATC CTA AAC CAG CTG TTC CGA GTC CTC ACC
 gly his val met ile ala phe leu pro thr ile leu asn gln leu phe arg val leu thr
 512
 542
 AGA GCC ACA CAG GAA GAA GTC GCG GTT AAC GTG ACT CGG GTC ATT ATT CAT GTG GTT GCC
 arg ala thr gln glu glu val ala val asn val thr arg val ile ile his val val ala
 572
 602
 CAG TGC CAT GAG GAA GGA TTG GAG AGC CAC TTG AGG TCA TAT GTT AAG TAC GCG TAT AAG
 gln cys his glu glu gly leu glu ser his leu arg ser tyr val lys tyr ala tyr lys
 632
 662
 GCT GAG CCA TAT GTT GCC TCT GAA TAC AAG ACA GTG CAT GAA GAA CTG ACC AAA TCC ATG
 ala glu pro tyr val ala ser glu tyr lys thr val his glu glu leu thr lys ser met
 692

FIG. 6A

722 752
ACC ACG ATT CTC AAG CCT TCT GCC GAT TTC CTC ACC AGC AAC AAA CTA CTG AGG TAC TCA
thr thr ile leu lys pro ser ala asp phe leu thr ser asn lys leu leu arg tyr ser

782 812
TGG TTT TTC TTT GAT GTA CTG ATC AAA TCT ATG GCT CAG CAT TTG ATA GAG AAC TCC AAA
trp phe phe phe asp val leu ile lys ser met ala gln his leu ile glu asn ser lys

842 |Cadherin Cleavage| 872
GTT AAG TTG CTG CGA AAC CAG AGA TTT CCT GCA TCC TAT CAT CAT GCA GCG GAA ACC GTT
val lys leu leu arg asn gln arg phe pro ala ser tyr his his ala ala glu thr val

902 932
GTA AAT ATG CTG ATG CCA CAC ATC ACT CAG AAG TTT GGA GAT AAT CCA GAG GCA TCT AAG
val asn met leu met pro his ile thr gln lys phe gly asp asn pro glu ala ser lys

962 992
AAC GCG AAT CAT AGC CTT GCT GTC TTC ATC AAG AGA TGT TTC ACC TTC ATG GAC AGG GGC
asn ala asn his ser leu ala val phe ile lys arg cys phe thr phe met asp arg gly
ref 2.1

1022 1052
TTT GTC TTC AAG CAG ATC AAC AAC TAC ATT AGC TGT TTT GCT CCT GGA GAC CCA AAG ACC
phe val phe lys gln ile asn asn tyr ile ser cys phe ala pro gly asp pro lys thr

1082 1112
CTC TTT GAA TAC AAG TTT GAA TTT CTC CGT GTA GTG TGC AAC CAT GAA CAT TAT ATT CCG
leu phe glu tyr lys phe glu phe leu arg val val cys asn his glu his tyr ile pro

1142 1172
TTG AAC TTA CCA ATG CCA TTT GGA AAA GGC AGG ATT CAA AGA TAC CAA GAC CTC CAG CTT
leu asn leu pro met pro phe gly lys gly arg ile gln arg tyr gln asp leu gln leu

1202 1232
GAC TAC TCA TTA ACA GAT GAG TTC TGC AGA AAC CAC TTC TTG GTG GGA CTG TTA CTG AGG
asp tyr ser leu thr asp glu phe cys arg asn his phe leu val gly leu leu leu arg

1262 1292
GAG GTG GGG ACA GCC CTC CAG GAG TTC CGG GAG GTC CGT CTG ATC GCC ATC AGT GTG CTC
glu val gly thr ala leu gln glu phe arg glu val arg leu ile ala ile ser val leu
ref 3.1

1322 1352
AAG AAC CTG CTG ATA AAG CAT TCT TTT GAT GAC AGA TAT GCT TCA AGG AGC CAT CAG GCA
lys asn leu leu ile lys his ser phe asp asp arg tyr ala ser arg ser his gln ala

1382 1412/471
AGG ATA GCC ACC CTC TAC CTG CCT CTG TTT GGT CTG CTG ATT GAA AAC GTC CAG CGG ATC
arg ile ala thr leu tyr leu pro leu phe gly leu leu ile glu asn val gln arg ile

1442 1472
AAT GTG AGG GAT GTG TCA CCC TTC CCT GTG AAC GCG GGC ATG ACC GTG AAG GAT GAA TCC
asn val arg asp val ser pro phe pro val asn ala gly met thr val lys asp glu ser

FIG. 6A (cont.)

1502 1532
CTG GCT CTA CCA GCT GTG AAT CCG CTG GTG ACG CCG CAG AAG GGA AGC ACC CTG GAC AAC
leu ala leu pro ala val asn pro leu val thr pro gln lys gly ser thr leu asp asn
ref 4.1 and 4.2

1562 1592
AGC CTG CAC AAG GAC CTG CTG GGC GCC ATC TCC GGC ATT GCT TCT CCA TAT ACA ACC TCA
ser leu his lys asp leu leu gly ala ile ser gly ile ala ser pro tyr thr thr ser

1622 1652
ACT CCA AAC ATC AAC AGT GTG AGA AAT GCT GAT TCG AGA GGA TCT CTC ATA AGC ACA GAT
thr pro asn ile asn ser val arg asn ala asp ser arg gly ser leu ile ser thr asp
ref 5.1 and 5.2

1682 1712
TCG GGT AAC AGC CTT CCA GAA AGG AAT AGT GAG AAG AGC AAT TCC CTG GAT AAG CAC CAA
ser gly asn ser leu pro glu arg asn ser glu lys ser asn ser leu asp lys his gln

1742 1772
CAA AGT AGC ACA TTG GGA AAT TCC GTG GTT CGC TGT GAT AAA CTT GAC CAG TCT GAG ATT
gln ser ser thr leu gly asn ser val val arg cys asp lys leu asp gln ser glu ile

1802 1832
AAG AGC CTA CTG ATG TGT TTC CTC TAC ATC TTA AAG AGC ATG TCT GAT GAT GCT TTG TTT
lys ser leu leu met cys phe leu tyr ile leu lys ser met ser asp asp ala leu phe

1862 1892
ACA TAT TGG AAC AAG GCT TCA ACA TCT GAA CTT ATG GAT TTT TTT ACA ATA TCT GAA GTC
thr tyr trp asn lys ala ser thr ser glu leu met asp phe phe thr ile ser glu val
ref 6.1

1922 1952
TGC CTG CAC CAG TTC CAG TAC ATG GGG AAG CGA TAC ATA GCC AGG AAC CAG GAG GGG TTG
cys leu his gln phe gln tyr met gly lys arg tyr ile ala arg asn gln glu gly leu

1982 2012
GGA CCC ATA GTT CAT GAT CGA AAG TCT CAG ACA TTG CCT GTT TCC CGT AAC AGA ACA GGA
gly pro ile val his asp arg lys ser gln thr leu pro val ser arg asn arg thr gly

2042 2072
ATG ATG CAT GCC AGA TTG CAG CAG CTG GGC AGC CTG GAT AAC TCT CTC ACT TTT AAC CAC
met met his ala arg leu gln gln leu gly ser leu asp asn ser leu thr phe asn his

2102 2132
AGC TAT GGC CAC TCG GAC GCA GAT GTT CTG CAC CAG TCA TTA CTT GAA GCC AAC ATT GCT
ser tyr gly his ser asp ala asp val leu his gln ser leu leu glu ala asn ile ala

2162 2192
ACT GAG GTT TGC CTG ACA GCT CTG GAC ACG CTT TCT CTA TTT ACA TTG GCG TTT AAG AAC
thr glu val cys leu thr ala leu asp thr leu ser leu phe thr leu ala phe lys asn
ref 7.1

2222 2252
CAG CTC CTG GCC GAC CAT GGA CAT AAT CCT CTC ATG AAA AAA GTT TTT GAT GTC TAC CTG
gln leu leu ala asp his gly his asn pro leu met lys lys val phe asp val tyr leu

2282 2312

FIG. 6A (cont.)

FIG. 6A (cont.)

ref 9.1

3122 ATC ATC CCC ATT TAT GAG AAG CGG AGG GAT TTC TTT GAA GAT GAA GAT GGA AAG GAG TAT
ile ile pro ile tyr glu lys arg arg asp phe phe glu asp glu asp gly lys glu tyr

3182 ATT TAC AAG GAA CCC AAA CTC ACA CCG CTG TCG GAA ATT TCT CAG AGA CTC CTT AAA CTG
ile tyr lys glu pro lys leu thr pro leu ser glu ile ser gln arg leu leu lys leu
ref 10.1

3242 TAC TCG CAT AAA TTT GGT TCT GAA AAT GTC AAA ATG ATA CAG GAT TCT GGC AAG GTC AAC
tyr ser asp lys phe gly ser glu asn val lys met ile gln asp ser gly lys val asn

3302 CCT AAG GAT CTG GAT TCT AAG TAT GCA TAC ATC CAG GTG ACT CAC GTC ATC CCC TTC TTT
pro lys asp leu asp ser lys tyr ala tyr ile gln val thr his val ile pro phe phe

3362 GAC GAA AAA GAG TTG CAA GAA AGG AAA ACA GAG TTT GAG AGA TCC CAC AAC ATC CGC CGC
asp glu lys glu leu gln glu arg lys thr glu phe glu arg ser his asn ile arg arg

3422 TTC ATG TTT GAG ATG CCA TTT ACG CAG ACC GGG AAG AGG CAG GGC GGG GTG GAA GAG CAG
phe met phe glu met pro phe thr gln thr gly lys arg gln gly gly val glu glu gln
ref 11.1

3482 TGC AAA CCG CGC ACC ATC CTG ACA GCC ATA CAC TGC TTC CCT TAT GTG AAG AAG CGC ATC
cys lys arg arg thr ile leu thr ala ile his cys phe pro tyr val lys lys arg ile

3542 CCT GTC ATG TAC CAG CAC CAC ACT GAC CTG AAC CCC ATC GAG GTG GCC ATT GAC GAG ATG
pro val met tyr gln his his thr asp leu asn pro ile glu val ala ile asp glu met

3602 xxxxxxxxxxxx Coiled coil 1 cont'd xxxxx 3632 xx
AGT AAG AAG GTG GCG GAG CTC CGG CAG CTG TGC TCC TCG GCC GAG GTG GAC ATG ATC AAA
ser lys lys val ala glu leu arg gln leu cys ser ser ala glu val asp met ile lys
ref 12.1

3662 xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx | 3692 GTC AAT GCT GGC CCA CTA GCA TAT
CTG CAG CTC AAA CTC CAG GGC AGC GTG AGT GTT CAG val gln val asn ala gly pro leu ala tyr
leu gln leu lys leu gln gly ser val ser val gln val

3722 GCG CGA GCT TTC TTA GAT GAT ACA AAC ACA AAG CGA TAT CCT GAC AAT AAA GTG AAG CTG
ala arg ala phe leu asp asp thr asn thr lys arg tyr pro asp asn lys val lys leu

3782 CTT AAG GAA GTT TTC AGG CAA TTT GTG GAA GCT TGC GGT CAA GCC TTA GCG GTA AAC GAA
leu lys glu val phe arg gln phe val glu ala cys gly gln ala leu ala val asn glu
| xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx

3842 xxxxxxxx Coiled coil 2 xxxxxxxxxxxxxx 3872 xx
CGT CTG ATT AAA GAA GAC CAG CTC GAG TAT CAG GAA GAA ATG AAA GCC AAC TAC AGG GAA
arg leu ile lys glu asp gln leu glu tyr gln glu glu met lys ala asn tyr arg glu

3902 xxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxxx . 3932 xxx|

FIG. 6A (cont.)

ATG GCG AAG GAG CTT TCT GAA ATC ATG CAT GAG CAG ATC TGC CCC CTG GAG GAG AAG ACG
 met ala lys glu leu ser glu ile met his glu gln ile cys pro leu glu glu lys thr

 3962 3992
 AGC GTC TTA CCG AAT TCC CTT CAC ATC TTC AAC GCC ATC AGT GGG ACT CCA ACA AGC ACA
 ser val leu pro asn ser leu his ile phe asn ala ile ser gly thr pro thr ser thr

 4022 |XXXX PBM XXXXX|
 ATG GTT CAC GGG ATG ACC AGC TCG TCT TCG GTC GTG TGA TTA CAT CTC ATG GCC CGT GTG
 met val his gly met thr ser ser ser ser val val STP

 4082 4112
 TGG GGA CTT GCT TTG TCA TTT GCA AAC TCA GGA TGC TTT CCA AAG CCA ATC ACT GGG GAG

 4142 4172
 ACC GAG CAC AGG GAG GAC CAA GGG GAA GGG GAG AGA AAG GAA ATA AAG AAC AAC GTT ATT

 4202 4232
 TCT TAA CAG ACT TTC TAT AGG AGT TGT AAG AAG GTG CAC ATA TTT TTT TAA ATC TCA CTG

 4262 4292
 GCA ATA TTC AAA GTT TTC ATT GTG TCT TAA CAA AGG TGT GGT AGA CAC TCT TGA GCT GGA

 4322 4352
 CTT AGA TTT TAT TCT TCC TTG CAG AGT AGT GTT AGA ATA GAT GGC CTA CAG AAA AAA AAG

 4382 4412
 GTT CTG GGA TCT ACA TGG CAG GGA GGG CTG CAC TGA CAT TGA TGC CTG GGG GAC CTT TTG

 4442 ↓ ref 13.1 4472
 CCT CGA CTC GTG CCG GAA ATC TGA TCG TAA TCA GGG TAC AGA ACT TAC TAG TTT TGT CTA

 4502 4532
 GGA GTA TGT TGT ATG ACT AGG ATT TGT GCT ATT ATC TCA TTC AAC AAC ATA GAG CAA GAA

 4562 4592
 TAG TGA GCT AAC TGA GCT AGA CAC TCA ATT AAT CCG CTA CTG GCT TCA AGT CAG AAC TTT
 ↓ ref 14.1
 4622 4652
 GTC ATT AAT CAT CGA CTC CGG GAC GGT CAT ATA TGT ATT ACA TTT CTA CAT TTT TAA TAC

 4682 4712
 TCA CAT GGG CTT ATG CAT TAA GTT TAA TTG TGA TAA ATT TGT GCT GGT CCA GTA TAT GCA

 4742 4772
 ATA CAC TTT AAT GGT TTA TTC TTG TCA TAA AAA TGT GCA ATA TGG AGA TGT ATA CAA GTC

 4802
 TTT ACT

FIG. 6A (cont.)

BAC sequences of Human CLASP 2

Ref 1.1

Sequence of BAC4 using primer HC2AS2, which spans nucleotides 327-346 of the cDNA. Exon sequence is underlined and represents nucleotides 356-375.

TTTCTACAGNGTNTACTCAGGTATGTGCTCCTTCAACAAAATTAGCAGTTGCTGCTCT
GTGACAAAGTTTGCACCATTTTGCAAGAAGAAAAAATCCTAATGTGTTATATTACTA
TATTTTACTCTATAGATCTTTTCTAAAGAAAGAAAGTACAACCTGAAGTGCTTATAT
GTATTCATATAAATGACTAGTACAAGCATCATTTTGCAACAGATTTCCTCTTTCATTG
GAGGATCTTCTTGATGTTATTTGTACACGATCAATTTTGTAGTCTTAATAAGATGAGGC
TGGGTGTGGTGGCTCACACCTGTAATCCTAGCATTTTGGAGGCCAAGGTGGGCAGAT
CACTTTAGCCCAGGGGTTTGAGACCAGCCTGGCCAAACATGGCAAAACCTTGTCTCTA
CAAAAATACNAAAATTATCCAGGCATGGTGATGTGTGCCTGTAGTCCCAACTNCCCTAG
GAGGCTAGGGGTAGGGGGATTGCAAGAGGCTGGGAGGGTCAAAGCCCNAANTGAG
CCATTGGTNCATGTCACTTGGACCCCAAGCNGGGGNGANCAAGAGCAAAGGACTNN
TGTNNTTTANAAAAAAAACCGGGCTACCATAACNNACCAACCCNCNNACCTACCCNACC
TTCCANNTTAAANAAGGCTTTGNCTTGCAAGGAAAANCAAAATNNCC

Ref 1.2

Sequence of BAC26 using primer HC2AS2, which spans nucleotides 327-346 of the cDNA. Exon sequence is underlined and represents nucleotides 351-375.

TCTGGTTTCTACAGTGTATACTNAGGTATGTGCTCCTTNAACAAAATTAGCAGTTGCT
GCTCTGTGACAAAGTTTGCACCATTTTGCAAGAAGAAAAAATCCTAATGTGTTATAT
TACTATATTTTACTCTATAGATCTTTTCTAAAGAAAGAAAGTACAACCTGAAGTGCTT
ATATGTATTCATATAAATGACTAGTACAAGCATCATTTTGCAACAGATTTCCTCTTTC
ATTGGAGGATCTTCTTGATGTTATTTGTACACGATCAATTTTGTAGTCTTAATAAGATG
AGGCTGGGTGTGGTGGCTCACACCTGTAATCCTAGCATTTTGGAGGCCAAGGTGGGC
AGATCACTTTAGCCCAGGGGTTTGAGACCAGCCTGGCCAAACATGGCAAAACCTTGTCT
TCTACAAAAATACAAAAATTATCCAGGCATGGTGATGTGTGCCTGTAGTCCCAGCTAC
CTAGGAGGCTAGGGTAGGGGGATTGCAAGAGGCTNGGAGGTCAAGGCCCGCAGTGA
GCCATGGTCATGTCACTGCACCCCAAGCCAGGGCCGACAGGAGCAAGACTNNTGTNT
CAAAAAAAAACAGNAACCAACANCCAACAACAACNACCTTTCNGCAAAAANAAGC
TTGCTNCAANGAAACCAAAATGNCTTCTTNTTTTCCCCCN

Ref 1.3

Sequence of BAC26 using primer HC2AS2, which spans nucleotides 327-346 of the cDNA. Exon sequence is not found within this sequence. This sequence most likely represent intron sequence since this sequence matches the intron sequence found in the previous two BAC sequences.

AGNNNNNCCNCTACNCCACTTTTAACCTTTTGAAAACACAGTGTTNCTCAANTATG
 CGTCTCTTACATATTAGCAGTTGCTGCTCTGTGACATAGTTGCACCATTNTGCAAGA
 AGAAAAAATCCTAAGTGTNATATCACTATATNNNTACTCTATAGATCTTNTCTAAAGA
 AAGAAAGTCAACTGATGTGCTTATATGTATNCATATAAATGACTAGTACATGCATCAT
 TTTGCAACAGATNTCTCTCACATTGGAGGATCTTCTNGANGNATTCGACACGATNAN
 TATTAGTCTNAATAAGATGANGCTGGTGTGGNGGTACACTGNATCTAGCATNTGGAN
 GCATGTGGCAGACACTTANCCNCGGTNGAGACAGCTGTCACTGNCNAACTGTCTCTN
 TAAANCAAANNCTCCGCNGGNGATGGGCTGAGCCAGTCCTAGNNGCTAGNTAGNGAT

FIG. 6A (cont.)

GNNGAGNTGTNGCACGNCGAGNGAGCATGNTCTGTACTGACTCATCAGGCGNCNACA
CGNTCTGTTCNAAAAACATACCACACACACTCNCACCTNCGCAAATGCTCTNNAAAN
ATGCTTNTTTCACACNGNTNCAATCNCCTATATNNTCTTCTATTCTNACGTNTNATTA
NNATCTTNCNCTGCANAAACNATNCGNCCACCTNNANNACCTTANGCTTNGTTTCACGC
TTATAGCTCCCCCTACACNTNNCAGCNNTTNCNNGTGAAGGGCCNCCCGAATCTACGA
NCATACTCTCTCCGTATATNGCCTCGGTCAACGCCATCTGCTGTNTNCTCNCNCTNG
CNNTTNANCNGTNGCCTATCTCTNNNCCGGATCCNCCCATATNNTNNTCTACTTAN
AGCGTAANNTNTNCCNCACTANTCACAACCTTNTNCTNNAACTCTATCTNCTCCTCT
CTACCACCTCACTTACTACCTNTTACNCANTCTCCTTNCNTNTCCACTGATCTCCACA
TAGCTGCTNTACTCGCCANTTTATCATATNCAACNCTCTACGCTNNNTNT

Ref 2.1

Sequence of BAC4 using primer HC2S1, which spans nucleotides 1107-1126 of the cDNA. Exon sequence is underlined and represents nucleotides 1079-1097.

CTTGTATTNAAAGAGGGTCTGCAGGAAGAAGTGTGTAGTCATAAATACCTCACTGGA
TATTTTATACAGGATTCTAAAAAACCTATTAGCAATAGTATGCTAGAAATAGTCATTA
GCTTCTTGACCTTCTTAGAACTGCACACTCTATTGCACTGTACAGATTTCAAGGATGGC
TGCAGGGGATTGATTTGAAAACTAAGGACACATTTCAATAAACAATGTCTTCAATTGAT
TTTTAGGGCTCCTCCTACTTCAATGAAGGACTTCAGGTAGCTTATAATTACAGACACA
GGCTCAATAACAATAAAAAAATTAGTAAGGCAGAGCTTTAAAAAAGGAAAAA
GATAATTCTACCAGAGAAAGGCTACATGGTGACTTCTGTTACCAGTAACAACCCCG
CACTACCTTTGGGTCTCCAGGAGCAAAACAGCTAATGTAGTTGTTGATCTGCTTGAAG
ACAAAGCCCCTGTCCATGAAGGTGAAACATCTCTGTGGAGGAAAAACAAGCAAAAAAG
TTATTTCAAGGTCCAAACATTTTCGGAAATTTGGATTCAAAGCAGGCATTATTGCTAAT
AAGTTTATCCACTGACATAAAAAACATGCCTTCAACATTGCCAGAGCACCTACTCTAT
TNTAGTCNCN

Ref 3.1

Sequence of BAC4 using primer C96AS, which spans nucleotides 1443-1452 of the cDNA. Exon sequence is underlined and represents nucleotides 1370-1422.

AATCAGCAGACCAAACAGAGGCAGGTAGAGGGTGGCTATCCTTGCCTGATGGCTCTG
AAAAGAAGACACACATGGTAAGTTTGACCCAGGATTCTGAGAACCGAACTAAGTTGG
TGCTGACCATCTCCTTTATTTGGATCCTTCTATAAAGACAGATATTTGATTTTAGTCC
CAAAATAGAGCAAAATCTTAGTGCTGTTACCATGAATTTTCTAACTGATTACTTTCTTT
ACACCACTTAAATAAAGGACATTATCAATGCACATTCCTTCCATTGGGGACCACTCA
CCCTTGAAGCATATCTGTCATCAAAAGAATGCTTTATCAGCAGGTTCTTGAGCACACT
GATGGCGATCAGACGGACCTCCCGGAACCTCCTGGAGGGCTGTCCCCACCTCCCTNAG
TAACAGTCCCACCAAGAGTGGTTTCTGCAGAACTCATCTGTTAATGAGTAGTCAAGC
TGGGAGGTCTGAAATGAGGATAGAACTACTTTGNGTTAGGAAAGATGCAATGCTCT
TTTGAATAAAACAAACAAACAAACAAACAAACAAACAAACAAACAAACAAACAAACAAAC
ATTTCAAGCCCACCTGGGGTNGGTCAAAGAGATGATCAGNANTTTGGCNTTNAAT
GAAGAAAGAAATNAATTNTCCAGGGGNTGTTCTNCTTTTGTAGCACANGGAGGGATNT
TAANTGAAAACCAATTTAAATCCAATTNAGNG

FIG. 6A (cont.)

Ref 4.1

Sequence of BAC4 using primer C2AS5, which spans nucleotides 1716-1735 of the cDNA. Exon sequence is underlined and represents nucleotides 1602-1703.

TTCTTTCTGCAAGGCTGTTCCCGAATCTGTGCTTATGAGAGATCCTCTCGAATCAGC
ATTTCTCACACTGTTGATGTTTGGAGTTGAGGTTGTATATGGAGAAGCTAAATGGAAA
TCAAGCCAACAATAAAGTTTTATTAAGACAGAACAAAATAAAGATGAGTACTGAACTT
TAAGGGAAATTGCTTTTATTGCACTTATTTTTCTGTTAGGAAGTTGGCTCAAGAGTT
GCATTCCATTACTTCACCTTTAAAGAACCAGGTCATATACAATGAGATAAAAAGAAAC
TAGTCTGAAACATTGAGATGTAAACATCAATTCACCTTGTTAGAAACCACCTTTGATCG
CTAAAGACTAAATGCATACCTGTTTCAGAATGTGATAGAATGAAGACTTAAAAAATT
AAAAGATAAATCCACCTACAACCTATCAAAATCACAAAATTAACCACACAACAACTTG
TAGCATTCAAACCTGGTAATAAAACACTGAGGAGCCTACCCAACTCTGAGGGGTGTCAT
GGGGTATTTTAAATTTTCGAGGAGAACACAGTGATATGTGACCTCAGCCAGAAGCTG
CTGTTTINAGCAGCAGGTTGGTGCTATGCTCCTTTTTGAAGACATATTTGTGAAGCTGG
GTATTTTGGGGGGCCTGCTTATGATAAAAANGGCAAGGTNTTCAATGNAGGGGN

Ref 4.2

Sequence of BAC26 using primer C2AS5, which spans nucleotides 1716-1735 of the cDNA. Exon sequence is underlined and represents nucleotides 1602-1703.

TTCTTTCTGGAAGGCTGTTACCCGAATCTGTGCTTATGAGAGATCCTCTCGAATCAG
CATTCTCACACTGTTGATGTTTGGAGTTGAGGTTGTATATGGAGAAGCTAAATGGAA
ATCAAGCCAACAATAAAGTTTTATTAAGACAGAACAAAATAAAGATGAGTACTGAACT
TTAAGGGAAATTGCTTTTATTGCACTTATTTTTCTGTTAGGAAGTTGGCTCAAGAGT
TGCATTCCATTACTTCACCTTTAAAGAACCAGGTCATATACAATGAGATAAAAAGAAA
CTAGTCTGAAACATTGAGATGTAAACATCAATTCACCTTGTTAGAAACCACCTTTGATC
GCTAAAGACTAAATGCATACCTGTTTCAGAATGTGATAGAATGAAGACTTAAAAAAT
TAAAAGATAAATCCACCTACAACCTATCAAAATCACAAAATTAACCNCACAACAACTT
GTAGCATTCAAACCTGGTAATAAAACACTGAGGAGCCTACCCAACTTTGAGGGGTGTC
AATGGGGTNTTTTAAATTTTCGNNGGANANCCAGTGNTATGGTGACCTTCACCCA
AGAAGCTTGTTTGTTTACCAAGCNAGGTTGNNTNTGCTCCTTTTAGAANACNNTA
TTTTNNNAAATNCTGGNTTTTTTNNNGGCCCCCTTNCNTTNTT

Ref 5.1

Sequence of BAC4 using primer C2S6, which spans nucleotides 1686-1705 of the cDNA. Exon sequence is underlined and represents nucleotides 1724-1736.

TTCTGGATAAGGTAATTGCTTTTACCCAACACAAATGTTTCTTATAATCAATGGATT
TAGCCCAAAGTAAACGTACTTCATGTTCTAGTGCCTTTTAAGTGTGACCTTTTGTTTT
TTTCTAAACCACCCGGCTGACCTGGAGTAGGTGATGAGAGCTTTAAGGTTGGGGCCC
ATTCTTGAAGTGCTCTGATTCTGTTTCCAGTACCTCAGATCCTGGGCAGGGTTTG
AGTGGAGCGTCTTGAGTGAATGGCTCTGGTGGGTTGAACGGGGAGGGACTCAAAAT
GCTGCCCATCTCAATTTCTGTAGTCTTTTTATTTATTTATTTTATTTTGGAGACAGAG
TCTCGCTCTGTGCGCCAGGCTGGAGTACAGCGGCACGATCTCAATTNACTGCAACCT
CCGCCTCC:TGGGTTCAAACGACTCCTCTGCCTCAGCCTCCCCAGCAGC:TGGGACCA
CAGGCACAAGCCACCACCGCCGGCTAATTTTTGTNTTTTTAGTA:GAGAT:GGGGTT
TCACCATAATTTGGCCAGGCTGGGCTCAAACCTCCTGACC:TCGTATCCGCNCCCTCGG
NCTNCCAAAGTGCTTGGGATTNCAGGCNGTGAGCCCACTTACACCTNNGGCAATTCC
CTGTNAGTCTTTTTTACCAGAGACACCATCATTCAACACAGCTTTTCCACCCACAA

FIG. 6A (cont.)

Ref 5.2

Sequence of BAC26 using primer C2S6, which spans nucleotides 1686-1705 of the cDNA. Exon sequence is underlined and represents nucleotides 1712-1736.

TGAGAAGAGCAATTTCTGGATAAGGTAATTGCTTTTACCCAACACAAATGTTTCTTA
TAATCAATGGATTTAGCCCAAAGTAAACGTACTTCATGTTCTAGTGCCTTTTAAAGTGT
GACCTTTTGTGTTTTTCTAAACCACCCGGCTGACCTGGAGTAGGTGATGAGAGCTTTA
AGGTTGGGGCCCATTCCTTGAAGTGCTCTGATTCTGTTTCCAGTACCTCAGATCCTG
GGCAGGGTTTGCAGTGGAGCGTCTTGAGTGAATGGCTCTGGTGGGTTGAACGGGGA
GGGACTCAAAATGCTGCCATCTCAATTTCTGTAGTCTTTTTATTTATTTATTTATTT
TTTGAGACAGAGTCTCGCTCTGTGCGCCAGGCTGGAGTACAGCGGCACGATCTCAAT
TCACTGCAACCTCCGNCCTCCCTGGGTTCAAACGACTCCTCTGNCTNAGNCTCCC:AGC
AGCCTGGGAACCACAGGCTCANGCCACCACGCCCGGCTAATTNTTGTAAATTTTNAAGT
AANAAATTGGGGGTTCTCACCATTNTGGCCCAAGNCTTGGGCCTAAAAACCTTNCNTNA
CCNTCGNCATTNCNCCCNACCNTGGGCNCTNCTCAAANGNGCTTGGGGATTTANC
ANNGGCNTTAACCCCCCNTATCACCGTGGNCCTTAATTT

Ref 6.1

Sequence of BAC4 using primer C2S7, which spans nucleotides 1918-1937 of the cDNA. Exon sequence is not found within this sequence. Since the primer is directed against exon sequence we presume that sequence derived from C2S7 is intron sequence.

NAGNGNGGGTTTNAGNCGTTTGAAGCCTGNNACGNGGTGNGTGCTNGAACTCTGTGG
GCTTTCAGGTACTGGGGTATCTGGGAGCCTGCTGTTTGCATTGCTAGTGCATCAGAC
CAGGGCTTTTCTCCCTGTAGCTGCTACTTATACACATAGCTCTAACTGAGATGATT
CTCCAGACAACTGATGCAGAGCAGCAAAAGCTTCTGCCGTTCTCCCTTCTAGGAGT
GTCTCCTTTCTTTGGAAAGAGATCATGAGGGGCTAGATTGTAATGAAGTGAGGCTCA
GTGCTTGAGCACATCCGGTAAAAAGTTCCAATATATTGGTCATAAAGTTTCTCATTCTT
TATAGCAGTTAATTTCTCTGGCTCATGAGTTTTCTTAGTTTTAATCTGACTTTTAAATT
AATGTCTCCAGCACCAGTCATATCCCCAGGGCAAACTCAAAGGCATGAGAGGCCAGA
CTCGGGTCTGGTCATAGCAACCCCTGTCTAGGGCCTTGGTCCCTGCCCTCCGCTTGT
GTGCTGTGGCGCAGGTCCTATGGGCCCTTAGGAAACAGGACCACCCTGTCGCACCCC
CTACAGAGACCAGCCAAGTTTGACATTAGATCACCGTAGCAATGTNTGCAAATTCCA
GTTTCTTGCTAAAAACAGGTTAAGCCTTGACGCCACTTTATCTGTAAGTGGCNGAGGTT
TTGACATAAAA

Ref 7.1

Sequence of BAC4 using primer C2S8, which spans nucleotides 2143-2162 of the cDNA. Exon sequence is underlined and represents nucleotides 2182-2219.

CTCTCGACACGCTGTTTCTATTAACATTGGCGTTTAAGGTTTGTATCAATTTGCTGTT
CGNGGTTCTAGTTTTACCTTTCACATTCATTCTGCTTGGTAAGCTCAGTGAGCACAAA
CTTACTATGTTGCATTTTACTTCAGCAATTATTTTGTCCCTGTAAGGAAACCATTA
TCTTTAAATTCCTTTAATGAAATCATTCCACAGTGAATGGCTTGAATGCCCTGAAATA
AAATTTAACTGGTCAGTGTGTGCTGCGCGCTTGGGTATGGTGGAACACGGTCTCTG
GAGGCAGTTAACTCTTGGCTCGAACCTTGAGGATGGTGAATATAGGCACCTAATCAG

FIG. 6A (cont.)

GCATTTCTGCCTTGAATATCTTTAAATATATCCAAATGTTATAGCGTTTAATTAGATT
TTATGTAGAAAGGAGCAATAAACACAAGACACATGTTTTTCAGTTTTTATCTGTTACT
GCATTAATGATAAAAAACGTTTTGGAGATAGAAAATGAAAGGGGTTTTTTTTTTGTCT
TGTTTTAAAGTTTTAGCAAATAATATTCAAGTAGGTGGAGATGGACTCTTCACCACTC
TCCTGTTTTTAGGAACCAATACTTTTTTCATTCTTGCTAAATGATTACTTCCATTTCTA
GCATAGAAAAGGAGAAAATTGGAATGAGTGTATATAT

Ref 8.1

Sequence of BAC4 using primer C2S9, which spans nucleotides 2992-3011 of the cDNA. Exon sequence is not found within this sequence. Since the primer is directed against exon sequence we presume that sequence derived from C2S9 is intron sequence

CGCTTTNAAATNCCAGCCGCTACTGCGGGGCGNTNAATTGAAACGTGTTGTTNTCT
GTGATGCCTGGCTCTGATTGTGTGGGATTGGTCATCAGTGGCGGTTGGCAGNTGGGG
TTCATGGAAGCGGCCATGGGGACTGATGGCAGGCCCTTGGATTGCCACCGCAGAGCC
TGGCAGTGTCTTTGGTCTGCATTCTACCGGCGAAAGTCTCATTTCACCTCACGTGTTA
TCTCTTGAAAGCATTCCTTTAGCGGGCTGTGTCTACCCCTCCATCCTCTCGTCCAAA
CTCCCCCTCCTTCTCTGTCTCTCTCCCTCCCATCCTCTCTCCCCAGTCTTCTTCTCT
ATGTTCTCTTCCCTCAGTGGTTTTCTCTTCTCTGTTTGACTTTCCAAGGTCATTTTGACTG
TTCCTGCTCCCAACTACAAAGATACTAAATCTCACCTAACCACTCTTCTTCTTTCTTA
ATGAAAGAATGTTTTTCAGTCCATCCCAAATTTGTGTGGACTTCACAAACCTTCTCTAA
AATGGAGCCTTTTCTCTTCTACTCTTGACTAGNTGGTAAACGCTCCATGTTCTTGGC
CAGAACTCCCTGGTGAGTAGCGTCACTCCCACTTTCTGTGCAGAACCAAGCCTCCT
AGAAAACCTCTTTGCANCTGAGTGGGTTGGGACACGCCCTTTNTTTGGG

Ref 9.1

Sequence of BAC4 using primer C2AS10, which spans nucleotides 3276-3295 of the cDNA. Exon sequence is underlined and represents nucleotides 3147-3234.

TTTANACCNATNTATCCGNGTCAGTTANAGGAGTCTCTGAGAAATTTCCGACAGCGGT
GTGAGTTTGGGTTCTTGTAAATATACTCCTTTCCATCTTCATCTTCAAAGAATCCCT
GTGACATAAAGCACAAATTAGAGCTATCCCTGAACGTAAGCCCAGGGCTTACCACCTA
GGAAGCGTTCTTTTATTACAAGGGGGGAAAAAAGGAATGGGTCTAAAAATCCAGCTG
AAATGGGCTTTCTGAATGAGAAAAGAAAATGCTAATAACATGAAGTCTAGGTGCAAAG
GTAAAGGAAAAACACAACATTGCAAACTTATTCAAGAATGCAGTCATTAAGTGTGAG
TGAAATGAAAGATTTTGGATACAAGACTAAGCTGTCCCAGGGAAGTCTAATGGGAGT
CAAGCCTGTTTCACTTTCCCAAGAAGCAGAACTCACTANAAAATGATGAGCAGCCCA
CGACAGGCAGGCTCAGAAGTGGACATGCCTCCCTTCTCCTGATGGCTNCCATGCACA
CAGGATTTTATGGCATGAACTGAAGCGTTTGGGGGTCTGGAGTAAGTTTAGTAAAAG
TTAGGTAAAGCTTGTATAAATTGTATTTTGTCTTACCCGATGAGAAAAAATATTN
AAGACCTGGTAGCTTCAATATTCAAGAAAAATATTTTTCATNTCACCCG

FIG. 6A (cont.)

Ref 10.1

Sequence of BAC4 using primer C2S11, which spans nucleotides 3167-3186 of the cDNA. Exon sequence is underlined and represents nucleotides 3231-3296.

NGNANGTGGAGCCNCGANCCAGGGACAATCTNAACCTNCTTAAACTGTACTCGGATN
AATTGGTTCTGAAAATGTCAAAATGATACAGGATTCTGGCAAGGTATTGACCATGTT
TGGANAAGTTTCATAGCAATGTAATGTTGTGATNCGATTACATATNATATATTTTTAA
ATGTNTATAGAAAAAACACANGAAAAATATTAAGGATTGTTGGCCCCGTGAGTGGCA
GGTGTATNTTCTNCTGATCCTTTAGNGCTTCCATTACATGCNTGACATTAAAAAAA
NCTTTATCGCCTAATTTTTGAAACATCTAATTTTACAAAATAATTAACCGTNTGGCCAN
GNATATNTCATTTTTAGGNCCAGCTATTTAGAAACTCTGACANAAATGAGGGGCTGT
GGCTTNCCTNCCCTNNACTTGNCCTCTTTTNNGNATGTACCACATGAACTTGNCNCCT
CTTTCNNCTNACCGGGTGGCATGTTANAGGACAGGTTGAAACCNCANTNGGGCNGGA
NTTNGGTNNAATTGGGACACAATGGTACNANGCTCTATNGGAATNGAAACTCTCCCN
ACNNCNGTGNNCCNTGGGGAAAATGNGNCNNATTCATTTTN

Ref 11.1

Sequence of BAC4 using primer C2S12, which spans nucleotides 3474-3493 of the cDNA. Exon sequence is not found within this sequence. Since the primer is directed against exon sequence we presume that sequence derived from C2S9 is intron sequence

AGNANNGTTNNGCAGCTGCANNTCTGGACCCANAGGCCGCANGGGCACGAGCCNNG
ACACGCTCGGCAAAGAGCTGTCCAGAGGGATTGAGAAGCTTCAGGACTGGAAGGGTC
TTTCGAGCTCAGTTAGCCACCCACACCCATTTCAGTTTCACATTTATCTAGTGCTT
CCTTTTGAATCATTGGGATGTTTTTCTGTTGATCTGTTGGCACTTCCTTCTTCCACAA
GACCAGAAGCTCATATCCAATCTAAGGTCACTTACCCTTCTGAGAATCTGATGAAAAT
GGCGTGCTTATGTGCCTAGATGCTTTTGCACACAGTCTAAGGTGACTTATGGACTCC
AGGTCCAGCAGCCACACCCAGTCCTGGGTCTCCGCACAGGGAGGGACCCGTCTTCAC
ACACCTGTCTCAGGTTCTAGCATTGGGCTGCTTCAGCGGTCTCAGGCTGTGAGTAAA
TGGGATGTGAGCTTGGATCGCCCCACGCTGTTGNCCCCCGGGGGGCTTGGCCAGCTG
GCCACTTNGAAATGCCTCCTTTTGCCCAAGGAAAGCTCACTGCATTTCAATGGGGNTTN
TCCACGAAGTTCANCTTTANGGG

Ref 12.1

Sequence of BAC4 using primer C2S13, which spans nucleotides 3645-3664 of the cDNA. Exon sequence is underlined and represents nucleotides 3683-3699.

AGNAAGGTNNCTCANTNAANNCAGCGTGAGNGTTCAGGTGAGCCAGGCACAGCAGGC
CGGAGGGGCAGCAGGGGACGTCCTTGCCCCCTGGGTGACTTGAGAGTCGTTTCCACTAA
CAAGGTCTACTTGAGAGCCTCGGTTTACCAAGTGATCCCTGCTCCCTTCCCCCAACGT
NTGTGACATTTCTCCTGATATCAGAGGGGGAGGAAACCTCATGATCCCTGCCCCCG
CCCCATGAGGACTGACTGTGGGGACAAAGAGCCAGATCTCATAGACTACCTGATTT
GTCAGTATTTGGGGAATTCTGGGTGCTGATTAGAAGCATCAAGACTCTTCTAAATNC
AAAGAAGTGTGGAGAGCAGTAGATTTTCTATAAACTGGTGTTGCTGGTTTCTATGA
AAATTGGATCCAAAAAAGTCCTTAAGTTTACCTCTTAATGGNATCTTTTGATTAAAT
GGAATTCATTATTTAATATAGCCCAATCAATCCAATTTTCTTTATTGGTAGCATTTT
TATGTTCTCTTTAAAAAATCTTGGNCTACCTCCAAAATTCACAGATGTTCTCCTAG
GGTTTCTCCTTTTGGTTCAAGCATCCCATTCANGTCTTGCAGTCCATTCTGGGG

FIG. 6A (cont.)

Ref 13.1

Sequence of BAC4 using primer C2S14, which spans nucleotides 4289-4308 of the cDNA. Exon sequence is underlined and represents nucleotides 4321-4448.

GACTTANATTTATTCTTCCTTGCAGAGTAGTGTTAGAATAGATGGCCTACAGAAAAAA
AAGGTTCTGGGATCTACATGGCAGGGAGGGCTGCACTGACATTGATGCCTGGGGGAC
CTTTGCCTCGAGGCTGAGCTGGAAAATCTTGAAAATATTTTTTTTTTCCTGTGGCAC
ATTCAGGTTGAATACAAGAACTATTTTTGTGACTATGTTTTTGATGACCTAAGGGAAC
TGACCATTGTAATTTTTGTACCANTGAACCANGAGATTTAAGTGCTTTTATATTCATTT
CCTTGCATTTAAGAAAATATGAAAGCTTAAGGAATTATGTGAGCTTAAACTAGTCAA
GCANTTTAGAACCAAAGGCCTATNTTNATAACCGCAACTATGCTNAAAAGNACAAAGT
AGTACAGNATATTGNTATGTACATATCATTTGGTAATACACNCCNGGCNTTCTGTACA
TATATGTATTACATTTCTACNTTTTAATACTCCCNTGGGCTTATGCCNTTAAGGTTAA
NTTNGGATAAATTTNGGCTGTTCCNGTNTATNCNATACNCTTTT

Ref 14.1

Sequence of BAC4 using primer C2AS15, which spans nucleotides 4680-4700 of the cDNA. Exon sequence is underlined and represents nucleotides 4660-4683.

ATGAGAATGTAATACATATATGTACAGAATGCCAGGACTGTATTAACAATGATATGTA
CATAACAATATACTGTACTACTTTGTACTTTTCAGCATAGTTGCGGTTATTAATATAG
GCCTTTGGTTCTAACTGCTTGACTAGTTTTAAGCTCACATAATTCCTTAAGCTTTTCAT
ATTTTCTTAAATGCAAGGAAATGAATATAAAAGCACTAAATCTCCTGGTTCAGTGGTA
CAAAAATTACAATGGTCAGTTCCTTAGGTCATCAAAAAGTCAAAAAATAGTTC
TTGTATTCAACCTGAATGTGCCACAGGAAAAAAAAAATATTTTCAAGATTTTCCAGCT
CAGCCTCGAGGCAAAAGGCCCCAGGCATCAATGTCAGNGCAGCCCTCCTGCCATGT
AGATCCCAGAACCCTTTTTTTTCTGTAGGCCATCTATTCTAACACTACTCTGCAGGGAG
AATAAAATCTAAAGNCCAGCTCAAGAGTGCTACCACACCTTTGTTAAGACACAATGAA
AACTTTGGATATTGGCAGGNGAGATTTAAAAAAATGTGCCCTTTCTTACCACTCCT
ATAGNAAAGTCTGGTTAAGAAATAACCGTTGGTCTTTATTTTCCTTTNTTTCCCTTC
CCTGGGNCTTCCTGGGGCTCGG

FIG. 6A (cont.)

HC2A	-----
KIAA	ASGNLDKNARFSATYRQDSNKLSDNDMLKLLADFRKPEKMAKLPVILGNLDITIDNVSSD
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	-----
KIAA	FPNYVNSSYIPTKQFETCSKTPITFEVEEFVPCIPKHTQPYTIYTNHLYVYPKYLYKYSQ
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	-----VLHHHQNPFEFYDEIK
KIAA	KSFAKARNIAICIEFKDSDEEDSQPLKCIYGRPGGPVFTRSAFAAVLHHHQNPFEFYDEIK
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	IELPTQLHEKHHLILLTFFHVSCDNSSKSGSTKKRDVVEQVGYSWLPLLKGRVVTSEQHI
KIAA	IELPTQLHEKHHLILLTFFHVSCDNSSKSGSTKKRDVVEQVGYSWLPLLKGRVVTSEQHI
rat	-----
HC4	-----
HC1	-----
HC3	-----
HC5	-----
HC2A	PVSANLPSGYLGYQELGMGRHYGPEIKWVDGGKPLLKISTHLVSTVYTQDQHLHNFQYC
KIAA	PVSANLPSGYLGYQELGMGRHYGPEIKWVDGGKPLLKISTHLVSTVYTQDQHLHNFQYC
rat	-----
HC4	-----
HC1	-----
HC3	-----GPGPARSTVSISLISNSARV
HC5	-----
HC2A	QKTESGAQALGNELVKYLKSLHAMEGHVMI AFLPTILNQLFRVLT-RATQEEVAVNVTRV
KIAA	QKTESGAQALGNELVKYLKSLHAMEGHVMI AFLPTILNQLFRVLT-RATQEEVAVNVTRV
rat	-----
HC4	-----MEIQVLIRFLSVILMQLFWVLPNMIHEDDVPI SCPMV
HC1	-----MSFLPIILNQLFKVLV-QNEEDEITTTVTRV
HC3	NRSRSLSNSNPDISGTPTSPDDEVRSIIIGSKGLDRSNSWVNTGGPKAAPWGSNPSPSAES
HC5	-----

FIG. 6B

		Ref.
HC2A	IIHVVAQCHEEGLESHLSYVKYAYKAEFYVASEYKTVHEELTKSMTTILKPSADFLTSN	
KIAA	IIHVVAQCHEEGLESHLSYVKYAYKAEFYVASEYKTVHEELTKSMTTILKPSADFLTSN	
rat		
HC4	LFHIVSKCHEEGLDLSYSSFIKYSEFPGKPSAPQAPLIHETLATMIALLKQSAFLAIN	
HC1	LPDIVAKCHEEQLDHSVQSYIKFVKTR---ACKERPVEDLAKNVTGLLK-SNDSPTVK	
HC3	TQAMDRSCNRMSSHTETSSFLQTLTGRLP---TKKLFHEELALQWVVCSG--SVR---E	
HC5		
	Cadherin Cleavage	
HC2A	KLLRYSWFFFDVLIKSMAQHLEIENSKVKLLIRNQRFPASYHHAETVVMMLMPHITQKFGD	
KIAA	KLLKYSWFFFDVLIKSMAQHLEIENSKVKLLIRNQRFPASYHHAETVVMMLMPHITQKFRD	
rat		
HC4	KLLKYSWFFFEIIAKSMATYILLEENKIKLTHCQRFPPKAYHHAHLSLFLAIT-IVESQYAE	6.1
HC1	HVLKHSWFFFAIILKSMQHLEIDTNKIQLERPQRFPESYQNELDNLMVLSHVWIKYKD	1.2/1.2/2.1/
HC3	SALQQAFFFEELMVKSMVHHLFYNDKLEAPRKSRFPFERFMDIAALVSTIASDIVSRFQK	
HC5		
HC2A	NPEASKNANHSLAVFIKRCFTFMDRGEVFKQIN---NYIS--CFAPGDKTLFEYKFEFL	2.1
KIAA	NPEASKNANHSLAVFIKRCFTFMDRGEVFKQIN---NYIS--CFAPGDKTLFEYKFEFL	
rat		
HC4	IPKESRNVNYSLASFLKCCILMDRGEVFNLIN---DYIS--GFSPKDPKVLAEYKFEFL	
HC1	ALEETRATHSVARFLKRCFTFMDRGCVEFKMVN---NYIS--MFSSGDLKTLQYKFDL	7.1
HC3	DTEVERLNTSLAFLNDLLSVMDRGEVFSLIKSCYKQVSSKLYSLPNPSVLVSLRDLFL	3.1/3.2
HC5		
HC2A	RVVCNHEHYIPLNLFM-----PFGKGRIQR-----YQDLQL---DYSLTDEF	
KIAA	RVVCNHEHYIPLNLFM-----PFGKGRIQR-----YQDLQL---DYSLTDEF	
rat		
HC4	QTICNHEHYIPLNLFM-----AFAPKPLQR-----VQDSNL---EYSLSDEY	
HC1	QEVCOHEHEFIPLCLPIRSANIPDPLTPSES-----TQELHASDMPEYSVTNEF	4.1/4.2
HC3	RIICSEHYVTNLNLFCSLLTPPASPSFSVSSATISQSSGFSTNVQDQKIANMFELS--VPE	
HC5	-----MNADTAPTSPCPISIS---SQNSSSCSSEQDQKIASMFDRTSRVPA	
HC2A	CRNHFLVGLLLREVGTALQEFRE----VRLIAISVLKNLLIKHSFDDRYASRSHQARIAT	3.1
KIAA	CRNHFLVGLLLREVGTALQEFRE----VRLIAISVLKNLLIKHSFDDRYASRSHQARIAT	
rat		
HC4	CKHHFLVGLLLRETSIALQDNYE----IRYTAISVIKNNLLIKHAFDTRYQHKNQQAQIAQ	
HC1	CRKHFLIGILLREVGFALQEDQD----VRHLALAVLKNIMAKHSFDDRYEPRKQAQIAS	8.1
HC3	RQOHYLAGIVLTELAVILDPAEGLFGLHKKVINMVINLLSSHSDPRYSDDPQIKARVAM	
HC5	SSTS--SPGLLFTELAAALDAEGEGISEVQRKAVSAIHSLLSSHLDLPPRCVKPEVKVIAA	
HC2A	LYLPLFGLLIENVQRINVRDVSPFFVNAG-MTVKDESLALPAVNPLVTPQKGSTLDNSLH	
KIAA	LYLPLFGLLIENVQRINVRDVSPFFVNAG-MTVKDESLALPAVNPLVTPQKGSTLDNSLH	
rat		
HC4	LYLPPFVGLLLENIQRLAGRDITLYSCAMPNSASRDEFFCG-----FTSP--AN--RGSLS	9.1
HC1	LYMPLYGMLLDNMPRIYLKOLYPTVNTSNQCSRDDLTNGGFQSQTAIKHANSVDTSPFS	
HC3	LYLELIGIIMETVPQLYDFTEHNRQRCRPICATDDYESE-----SG---SMIS	
HC5	LYLPLVGIIIDALPQLCDFTVADTRRYR---TSGSDEEQE-----GA---GAIT	
	4.1/4.2	
HC2A	KDLLGAISGLASPYTTSTPNINSVRNADSRGSLISTDSGNLPERNSEKSNSLDKHQSS	5.1/5.2
KIAA	KDLLGAISGLASPYTTSTPNINSVRNADSRGSLISTDSGNLPERNSEKSNSLDKHQSS	
rat		
HC4	TDKDTAYGSFQNG-----HGKIKREDSRGSLIP-EGATGFPDQNGTGEN-----TRQS	
HC1	KDVINSIAAFSS-----IAISTVNHADSRASLSDSNPSTNEKSSEKTDNCEKIPRPL	10.1
HC3	QTVAMAIAGTSVPQ-----LTRPGSFLLTSTSGRQHT-----	3.1
HC5	QNVALAIACNNFN-----LKTSG-IVLSSLEFYKQYN-----	2.1

FIG. 6B (cont.)

		Ref.
HC2A	TLGNSVVRCDKLDQSEIKSLLMCFYILKMSDDALFTYWN-KASTSELMDFFTISEVCL	
KIAA	TLGNSVVRCDKLDQSEIKSLLMCFYILKMSDDALFTYWN-KASTSELMDFFTISEVCL	
rat	-----	
HC4	STRSSVSQYNRLDQYEIRSLIMCYLYIVKMI SEDTLLTYWN-KVSPQELINILILEVCL	
HC1	ALIGSTLRFDRLDQAE TRSLLMCFLHIMKTISYETLIAYWQ-RAPSPEVSDFFSIIIDVCL	11.1/11.2
HC3	-----TFSAESRSLLICLLWVLKN-ADETVLQKWFTDLSVLQINRLIDLLYL CV	
HC5	-----MLNADTTRNIMICFLWIMKN-ADQSLIRKWIADLPSTQLNRILDLLFICV	
HC2A	HQFQYMGKRYIARNQEGLG--PIVHDRKS-----QTLFVSRNRTGMM	6.1
KIAA	HQFQYMGKRYIAR-----TGMM	
rat	-----	
HC4	FHFRYMGRNIARVHDAWLSKHFGIDRKS-----QTMPALNRNSGVM	
HC1	QNFYRLGKRNIIRKIAAAF--KFVQSTQNNGT LKGSNPSCQTSGLLAQWMHSTSRHEG HK	
HC3	SCFEYKGKKVFERMNSLTFK--KSKDMRAK-----LEEAILGSIGARQEMV	
HC5	LCFEYKGKQSSDKVSTQVLQ--KSRDVKAR-----LEEALLRGE GARGEMM	
HC2A	HARLQQL-----GSLDNS-----LTFNHSYGHSDADVLHQSLLEANIATEVC	
KIAA	HARLQQL-----GSLDNS-----LTFNHSYGHSDADVLHQSLLEANIATEVC	
rat	-----	
HC4	QARLQHL-----SSLESS-----FTLNHSSTTEADIFHQALLEGNTATEVS	
HC1	QHRSQTLPIIRGK--NALS NPKL----LQMLDNTMTS NSNEIDIVHHVDTEANIATEGC	12.1/12.2
HC3	RRSRGQIERSPSGSAFGSQENLRWRKDMTHWRQNT EKLDKSRAEIEHEALIDGNLATEAN	6.1/6.2
HC5	RRRAPGND RFP-----CLNENLRWKKEQTHWRQANEKLDKTKAELDQEALISGNLATEAH	
HC2A	LTALD TSLSFTLAFKNQLLADHGHNPLMKKVFDVYLCFLQKHQSE TALKNVFTALRS LIY	7.1
KIAA	LTALD TSLSFTLAFKNQLLADHGHNPLMKKVFDVYLCFLQKHQSE TALKNVFTALRS LIY	
rat	-----KLSRGHSP LMKKVFDVYLCFLQKHQSE MALKNVFTALRS LIY	
HC4	LTVIDTISFFTQC FKT HFLNNDGHNPLMKKVFDIHLAFLKNGQSEVSLKHVFASLRAFIS	
HC1	LTILD LVSLFTQTHQRQLQCCDQNSLMKRGFDTYMLFFQV NQSATALKHVFASLR L FVC	13.1
HC3	LIILD TLEIVVQTVS--VTES--KESILGGVLKVLLHSMACNQSAVYLOHCFATORALVS	
HC5	LIILDMQENIIQASS--ALDC--KDSLLGGVLRVLVNSLNC DQSTTYLTHCFATLRALIA	3.1
HC2A	KFPSTFFYEGRADMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKS FVRTH	
KIAA	KFPSTFFYEGRADMCAALCYEILKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKS FVRTH	
rat	KFPSTFFYEGRADMCAALCYEVLKCCNSKLSSIRTEASQLLYFLMRNNFDYTGKKS FVRTH	
HC4	KFPSAFFKGRVNMCAAF CYEVLKCC TSKISSRNEASALLYL LMRNNFEYTKRKTFLRTH	
HC1	KFPSAFFQGPADLCGSFCYEVLKCCNHRSRSTQTEASALLYL FMRKNFEF NKQKSIVRSH	
HC3	KFPELLFEEETEQCADLCRLRLRHCCSSSIGTIRSHPSASLYLLMRQNF EIGN--NFARVK	7.1/7.2
HC5	KFGDILLFEEVEQE QCDLCHQVLHHCSSSMDVTRSQACATLYLLMRFSFGATS--NFARVK	
HC2A	LQV IISVSQLIADVVGIGGTRFQQSLSIINN CANS DRLIKHTSFSSDVKDLTKRIRTVLM	
KIAA	LQV IISVSQLIADVVGIGGTRFQQSLSIINN CANS DRLIKHTSFSSDVKDLTKRIRTVLM	
rat	LQV IISLSQLIADVVGIGGTRFQQSLSIINN CANS DRLIKHTSFSSDVKDLTKRIRTVLM	
HC4	LQIIIAVSQLIADVALSGGSRFQESLFTIINN FANS DRMLARAFPAEVKDLTKRIRTVLM	
HC1	LQLIKAVSQLIAD-AGIGGSRFQHS LAITNNFANGDKQMKQNSNFP AEVKDLTKRIRTVLM	14.1/14.2/15
HC3	MQVPM SLSL VGTSQNFNEEFLRRSLKITLTYAEEDLELRETTFPDQVDLVFNLMILS	
HC5	MQVTMSLASLVGRAPDFNEEHLRRSLRILAYSEEDTAMQMTPFPTQVEELL CNLNSILY	

FIG. 6B (cont.)

		Transmembrane	Ref.
HC2A	ATAQMKHEHNDPEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGL	LSEAAMCYVHV	
KIAA	ATAQMKHEHNDPEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGL	LSEAAMCYVHV	
rat	ATAQMKHEHNDPEMLVDLQYSLAKSYASTPELRKRWLDSMARIHVKNGL	LSEAAMCYVHV	
HC4	ATAQMKHEHNDPEMLVDLQYSLAKSYASTPELRKRWLDSMAKIHVKNGL	FSEAAMCYVHV	
HC1	ATAQMKHEHNDPEMLVDLQYSLANSYASTPELRRTWLESMAKIHARNGDL	LSEAAMCYIHI	16.1/16.2
HC3	ITVQCKEHDPEMLIDLMYRIAKGYQTS PDLRLTWLQNMAGKHSERSN	HAEAAQCLVHS	
HC5	ITVQCKEHDPEMLIDLMYRIAKSYQAS PDLRLTWLQNMAEKHTKKKCY	TEAAMCLVHA	
SH3			
HC2A	TALVAEYLTIRKGV-----	FRQGCTAFRVITPN	
KIAA	TALVAEYLTIRKTA-----	VQWEPPLPHSHSACLRRSRGGVFRQGCTAFRVITPN	
rat	TALVAEYLTIRHEAD-----	LALQREPPVFPPYSHTSCQRKSRGGMFRQGCTAFRVITPN	
HC4	AALVAEFLIRFKL-----	FPNGCSAFKKITPN	
HC1	AALTAEYIIRFGYWKVEKIC-----	TASLISEDTHPCDSNSLLTTPSGGSMFSMGWPAFLSITPN	
HC3	AALVAEYLTIRLED-----	RYLFPVGCVTFCNISSN	8.1/8.2
HC5	AALVAEYLTIRLED-----	HSYLPVGSVSFONISSN	
HC2A	IDEEASMTEDVGMQD-----	VHENEDVIMELLEQCADGLWKAERYELIADIYKLIPI	8.1
KIAA	IDEEASMTEDVGMQD-----	VHENEDVIMELLEQCADGLWKAERYELIADIYKLIPI	
rat	IDEEASMTEDVGMQD-----	VHENEDVIMELLEQCADGLWKAERYELIADIYKLIPI	
HC4	IDEEGAKGELAGMD-----	VHYSEEVILLELLEQCVNGLWKAERYEIISEISKLIPI	
HC1	IKKEGAAKEDSGMHD-----	TPYNEIILVEQLYMCGEFLWKSERYELIADVNPPIIAV	17.1/17.2
HC3	VLEKSAVSDVVSPEEGICSGKYFTE SGLVGLLEQAAASF	SMAGMYEAVNEVYKVLPI	
HC5	VLEESVSEDTLSPDEDCVCACQYFTE SCLVGLLEQAAELF	STGGLYETVNEVYKVLPI	
ITAM ITAM ITAM ITAM			
HC2A	YTKRFD-----	TYFRVAFYFGQ	9.1
KIAA	YTKRRDFFELAHYDTLHRAYSKVTEVMHSGRRLLGTYFRVAFYFGQ	AGSWDLLPGGLEFGQ	
rat	SMKSGGTLETHYDTLHREYSKVTEVITR-----	A-----	
HC4	YENRREFENLTQYRTHGAYTKILEVMHTKKRLG-----	TYFRVAFYFGQ	
HC1	FEKQDFKKLSDIYYDIHRSYLKVAEVVNSEKRLFG-----	TYFRVAFYFGQ	
HC3	KEANRDAKELSTHCKLQEAFSKIVHQTGWERMFG-----	TYFRVAFYFGQ	9.1
HC5	LEANREFRKLTLTHSKLQRAFDIVNKH--KRMFC-----	TYFRVAFYFGQ	
HC2A	-FFEDEDGKEYIYKEPKLTPLSEISQRLKLYSDKFGSENVKMIQDSGK	NPKDLDSKYA	10.1
KIAA	GFFEDEDGKEYIYKEPKLTPLSEISQRLKLYSDKFGSENVKMIQDSGK	VNPKDLDSKYA	
rat	GFFEDEDGKEYIYKEPKLTPLSEISQRLKLYSDKFGSENVKMIQDSGK	VNPKDLDSKFA	
HC4	SFFEDEDGKEYIYKEPKLTGLSEISLRLVKLYGEKFGTENVKI	IQSDKVNAKELDPKYA	
HC1	GFFFEEDGKEYIYKEPKLTGLSEISQRLKLYADKFGADNVKI	IQSDKNVNPKDLDPKYA	
HC3	TKPCDLDEQEFVYKEPAITKLAEISHRIEGFYGERFGEDVVE	VIKDSNPNVDKCKLDPNKA	10.1/10.2
HC5	SKFGDLDEQEFVYKEPAITKLPEISHRLEAFYGCQCGAEFVE	VIKDSNPNVDKCKLDPNKA	4.1
HC2A	YIQVTHVIPFFDEKELQERKTEFERSHNIRRFMFEMPFTQTGKRQGGV	EEQCKRRITILTA	11.1/11.2
KIAA	YIQVTHVIPFFDEKELQERKTEFERSHNIRRFMFEMPFTQTGKRQGGV	EEQCKRRITILTA	
rat	YIQVTHVIPFFDEKELQERKTEFERSHNIRRFMFEMPFTQTGKRQGGV	EEQCKRRITILTA	
HC4	HIQVTVKPYFDDKELTERKTEFERNHNSRFVFEAPYTLSGKKQGCIEE	QCKRRITILT	
HC1	YIQVTVVTPFFEEKEIEDRKTDFFEMHNNIRFVFETPFTL	SGKKHGGVAAEQCKRRITILT	18.1
HC3	YIQITYVEPYFDTYEMKDRITYFDKNVLRFRMYCTPFTLD	GRAHGEHQYRRNTVLT	
HC5	YIQITFVEPYFDEYEMKDRVITYFEKNFNLRFRMYTTPFTLE	GRPRGELHEQYRRNTVLT	

FIG. 6B (cont.)

	Coiled-Coil 1	Ref
HC2A	IHCFFPYVKKRIPVMYQHHTDLNPIEVAIDEMSKKVAELRQLCSSAEVDMIKLQKLQGSV	
KIAA	IHCFFPYVKKRIPVMYQHHTDLNPIEVAIDEMSKKVAELRQLCSSAEVDMIKLQKLQGSV	
rat	IHCFFPYVKKRIPVMYQHHTDLNPIEVAIDEMSKKVAELRQLCSSAEVDMIKLQKLQGSV	
HC4	SNSFPYVKKRIPINCEQQINLKPIDGATDEIKDKTAELOKLCSSSTDVDMIQLQLKLGWV	
HC1	SHLFPYVKKRIQVISQSSTELNPIEVAIDEMSRKVSELNQLCTMEEVDMISLQKLQGSV	
HC3	SHAFPIKTRVNVTHKEIIILTPIEVAIDEMQKKTQELAFATHQDPADPKMLQMVLOGSV	11.1
HC5	MHAFPIKTRISVIOKEEFVLTPIEVAIDEMKKKTLQLAVAINQEPDAKMLQMVLOGSV	
	Coiled-Coil 2	
HC2A	SVQVNAGPLAYARAFLLDDTNTKRYPDNKVKLLKEVFRQFVEACGQALAVNERLIKEDQLE	11.1/12.1
KIAA	SVQVNAGPLAYARAFLLDDTNTKRYPDNKVKLLKEVFRQFVEACGQALAVNERLIKEDQLE	
rat	SVQVNAGPLAYARAFLLDDTNTKRYPDNKVKLLKEVFRQFVEACGQALAVNERLIKEDQLE	
HC4	SVQVNAGPLAYARAFLLDSQASKYPPKKVSELKDMFRKFIQACSI ALELNERLIKEDQVE	
HC1	SVKVNAGPMAYARAFLEETNAKKYPDNQVKLLKEIFRQFADACGQALDVNERLIKEDQLE	
HC3	GTTVNQGPLEVAQVFLSEIPSDPKLFRHHNKLRLCFKDFTKRCEDALRKKNKSLIGPVQKE	
HC5	GATVNQGPLEVAQVFLAEIPADPKLYRHHNKLRLCFKEFIMRCGEAVEKNKRLITADQRE	
	Coiled-Coil 2	
HC2A	YQEEMKANYREMAKELSEIMHEQICPLEEKTS-VLPNSLHIFNAISGTPTSTMVHGMTSS	
KIAA	YQEEMKANYREMAKELSEIMHEQIG-----	
rat	YQEEMKANYREIRKELSDIIVPRICPGEDKRA TKFPAHLQRHQD TNKHSGSRVDQFILS	
HC4	YHEGLKSNERDMVKELSDIIEHQIQEDTMHSPWMSNTLHVFCASGTSSDRQYGS PRYA	
HC1	YQEELRSHYKDMSELSTVMNEQITGRDDL SK---RGVDQTC TRVISKATPALPTVSISS	19.1
HC3	YQRELG---KLSS---PZ-----	
HC5	YQELKKYNKLNKLNLRPMIERKIPELYKPIFRVESQKRDSFHRSSFRKCETQLSQGSZ-	
	PBM	
HC2A	SSVVZ-----	
KIAA	-----	
rat	CVTLPHPEPHVGTCTCFVMCKLRTTFRANHWFCQAQEEAMNGGREKEPWTVI FNSRFYRSWGK	
HC4	EVZ-----	
HC1	SAEVZ-----	
HC3	-----	
HC5	-----	
HC2A	-----	
KIAA	-----	
rat	VHIFF-----	
HC4	-----	
HC1	-----	
HC3	-----	
HC5	-----	

FIG. 6B (cont.)




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 9D

	10	20	30	40	50	60	70	80	
1	AATTGTAATA	CGACTCACTA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCTC	GAGGTGACG	GTATCGATAA	GCTTGATATC	80
81	GAATTCGGCA	CGAGTTTAC	ACCATCACC	AAACCCAGAA	TTTTATGATG	AGATTAAAT	AGAGTTGCC	ACTCAGCTGC	160
161	ATGAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTCCG	CTGTGACAA	TCAAGTAAAG	GAAGCCAGAA	GAAGAGGGAT	240
241	GTCGTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCCTCC	TGAAGAGCGG	AAGGGTGGTG	ACAAGCCGAG	AGCACATCCC	320
321	GGTCTCGGG	AACCTTCCTT	CGGGCTATCT	TGGCTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAATTAATAT	400
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTACATA	480
481	ATTTTTTCCA	GTACTGTCAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACCTG	TAAAGTACCT	TAAAGTCTCG	560
561	CATGCGATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTC	CGAGTCCTCA	CCAGAGCCAC	640
641	ACAGGAAGAA	GTCCGCGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TTGGAGAGCC	720
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAACTG	800
801	ACCAAATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCCTCACCAG	CAACAACTA	CTGAGGTACT	CATGGTTTTT	880
881	CTTTGATGTA	CTGATCAAA	CTATGGCTCA	GCAATTTGATA	GAGAACTCCA	AAGTTAAGTT	GCTCCGAAAC	CAGAGATTTC	960
961	CTGCATCCTA	TCATCATGCA	GCGGAAACCG	TTGTAAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTGG	AGATAATCCA	1040
1041	GAGGCACTCA	AGAACCGGAA	TCATAGCCTT	GCTGCTTTCA	TCAAGAGATG	TTTCACTTTC	ATGGACAGGG	GCTTTGCTCT	1120
1121	CAAGCAGATC	AACAACCTACA	TTAGCTGTTT	TGCTCCTGGA	GACCCAAAGA	CCCTCTTTGA	ATACAAGTTT	GAATTTCTCC	1200
1201	GTGTAGTGTG	CAACCATGAA	CATTATATTC	CCTTGAACCT	ACCAATGCCA	TTTGGAAAG	GCAGGATTCA	AAATATACCA	1280
1281	GACCTCCAGC	TTGACTACTC	ATTAACAGAT	GAGTTCTGCA	GAACCACTT	CTTGGTGGGA	CTGTACTGTA	GGGAGGTGGG	1360
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATTAAG	CATTCTTTTG	1440
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACCTCTAC	CTGCTCTGT	TTGGTCTGCT	GATTGAAAC	1520
1521	GTCCAGCGGA	TCAATGTGAG	GGATGTGTCA	CCCTTCCCTG	TGAACCGCGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT	1600
1601	ACCAGCTCTG	AATCCGCTCG	TCACGCCGCA	GAAGCGAAGC	ACCTTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGGCCCCA	1680
1681	TCTCCGCGAT	TGCTTCTCCA	TATACAACCT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCGAG	AGGATCTCTC	1760
1761	ATAAGCACAG	ATTCCGGTAA	CAGCCTTCCA	GAAGGAAATA	GTGAGAAAG	CAATTCCCTG	GATAAGCAC	AACAAGTAG	1840
1841	CACATTGGGA	AATTCGCTGG	TTCGCTGTGA	TAACTTTGAC	CAGTCTGAGA	TAAAGAGCCT	ACTGATGTGT	TTCTCTTACA	1920
1921	TCTTAAAGAG	CATGCTGTAT	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACTTATTGA	TTTTTTTACA	2000
2001	ATATCTGAAG	TCTGCTTGCA	CCAGTTCCAG	TACATGGGGA	AGCGATACAT	AGCCAGGAAC	CAGGAGGGGT	TGGGACCCAT	2080
2081	AGTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTCCCGT	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG	2160
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAAAC	ACAGCTATGG	CCACTCCGAC	GCAGATGTTT	TGCACCACTC	ATTACTTGAA	2240
2241	GCCAAATATG	CTACTGAGGT	TTGCCCTGCA	GCTCTGGACA	CGCTTTCTCT	ATTTACATTT	CGGTTTAAAG	ACCAGCTCCT	2320
2321	GCCCGACCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGTCTACC	TCTGTTTTCT	TCAAAAACAT	CAGTCTGAAA	2400
2401	CGCCTTTAAA	AAATGTCTTC	ACTGCTTTAA	GGTCCCTAAT	TTATAAGTTT	CCCTCAACAT	TCTATGAAGG	GAGAGCGGAC	2480
2481	ATGTGTGCGG	CTCTGTGTTA	CGAGATTCTC	AAGTGTGTGA	ACTCCAAGCT	GAGCTCCATC	AGGACGGAGG	GCTCCAGAT	2560
2561	GCTCTACTTC	CTGATGAGGA	ACAACCTTTG	TTACATGGGA	AAGAAGTCTT	TTGTCCGGAC	ACATTTGCAA	CTCATATAT	2640
2641	CTGTGAGCCA	GCTGATAGCA	GACGTTGTTG	GCATTGGGGA	AACCAGATTG	CAGCAGTCCC	TGTCATCAT	CAACAACCTG	2720
2721	GCCAAACAGT	ACCGGCTTAT	TAAAGCACACC	AGCTTCTCTT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTGCT	2800
2801	AATGGCCACC	GCCAGATGGA	AGGAGCATGA	GAACGACCCA	GAGATGCTGG	TGGACCTCCA	GATACGCCCT	CCCAATCTCT	2880
2881	ATGCCAGCAC	GCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG	2960
2961	GCAGCAATGT	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	AATATCTCAC	ACGGAAGGCG	GTGTTTAGAC	AAGGATGCAC	3040
3041	CGCCTTCAGG	GTCAATTACC	CAAACTCCGA	CGAGGAGCCC	TOCATGTAGG	AAGACGTGGG	GATGCAAGGT	GTCCATTCCA	3120
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	AGTCCGCAGA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCGCCGAC	3200
3201	ATCTACAAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTACAA	3280
3281	GGAAACCCAA	CTCACACCGC	TGTCGGAAAT	TTCTCAGAGA	CTCCTTAAC	TGTACTCGGA	TAAATTTGGT	TCTGAAAATG	3360
3361	TCAAAATGAT	ACAGGATCTT	GGCAAGGTCA	ACCCTAAGGA	TCTGGATTCT	AAGTATGCA	ACATCCAGGT	GACTCAGCTC	3440
3441	ATCCCTTCT	TTGACGAAA	AGAGTTGCAA	GAAGGAAA	CAGACTTTGA	GAGATCCCA	AACATCCGCC	GCTTCATGTT	3520
3521	TGAGATGCCA	TTTACGCAGA	CCGGGAAGAG	GCAGGGCGGG	GTGGAAGAGC	AGTGCACACG	GCCCAACATC	CTGACAGCCA	3600
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCA	GTACCAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC	3680
3681	ATTGACGAGA	TGAGTAAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGTCTCTC	GGCCGAGGTG	GACATGATCA	AACTGCAGCT	3760
3761	CAAACTCCAG	GGCAGCGTCA	GTGTTCAAGT	CAATGCTGGC	CCACTAGCAT	ATGCGCGAGC	TTTCTTAGAT	GATACAAACA	3840
3841	CAAGCGGATA	TCCTGACAA	AAAGTGAAGC	TGCTTAAGGA	AGTTTTTCAGG	CAATTTGTGG	AAGCTTGCGG	TCAAGCCTTA	3920
3921	GCGGTAAACG	AACGCTGTAT	TAAAGAGAC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACATCAGGG	AAATGGCGAA	4000
4001	GGAGCTTTCT	GAATCATGCT	ATGAGCAGAT	CTGCCCCCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTCC	CTTCACATCT	4080
4081	TCACAGCCAT	CAGTGGGACT	CCAACAAGCA	CAATGGTTCA	CGGGATGACC	AGCTCGTCTT	CGGTCTGTCT	ATTACATCTC	4160
4161	ATGCCCCCTG	TGTGGGACT	TGCTTTGTCA	TTTCCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA	4240
4241	CAGGGAGGAC	CAAGGGGAAG	GGGAGAGAAA	GGAAATAAAG	AACAACGTTA	TTTCTTAACA	GACTTTCTAT	AGGAGTTGTA	4320
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGSCAATATT	CAAAAGTTTC	ATTGTGTCTT	AACAAAGGTG	TGGTAGACAC	4400
4401	TCTTGAGCTG	GACTTAGATT	TTATCTCTCC	TTGCAGAGTA	GTGTTAGAAT	AGATGGCCTA	CAGMAAAAA	AGGTTCTGGG	4480
4481	ATCTACATGG	CAGGGAGGGC	TGCACTGACA	TTGATGCTCT	GGGGACCTTT	TGCTCTGACT	CGTGCCGGAA	ATCTGATCGT	4560
4561	AATCAGGGTA	CAGAAGTTAC	TAGTTTGTCT	TAGGAGTATG	TTGTATGACT	AGGATTGTGT	CTATTCTCTC	ATTCAACAAC	4640
4641	ATAGAGCAAG	AATAGTGAGC	TAACTGAGCT	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGAACT	TTGTCAATTA	4720
4721	TCATCGACTC	CGGGACGGTC	ATATATGTAT	TACATTTCTA	CATTTTAAAT	ACTCACATGG	GCTTATGCA	TAAAGTTAAT	4800
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGCTTTA	TTCTTGTCTAT	AAAAATGTGC	AATATCGAGA	4880
4881	TCTATACAA	TCTTTACT							4898

FIG.10A

	10	20	30	40	50	60	70	80
1	MEGHVMI	PTILNQLFRV	LTRATQEEVA	VNVTRVTHV	VAQCHEEGLE	SHLSYVVKYA	YKAEFYVASE	YKTVHEELTK 80
81	SMTTILKPSA	DFLTSTNKLRL	YSWEFFDVL	KSMAQHLTEN	SKVKLLRNQR	FPASYHHAAE	TVVNMMLMPHI	TQKFGDNPEA 160
161	SKNANHSLAV	FIKRCFTFMD	RGFVFKQINN	YISCFAPCDP	KTLEFYKFEF	LKVVCNHEHY	IPLNLMPPFG	KGRIQRYQDL 240
241	QLDYSILTDEF	CRNHFLVGLL	LREVGTAQOE	FREVRLIAIS	VLKNLLIKHS	FDDRYASRSH	QARIATLYLP	LFGLLIENVQ 320
321	RINVRDVSPF	PVNAGMTVKD	ESLALPAVNP	LVTPOKGSTL	DNSLHKDLLG	AISGIASPYT	TSTPNINSVR	NADSRGSLIS 400
401	TDSGNSLPER	NSEKSNSLDK	HQQSSTLGNS	VVRCDILQDS	EIKSLLMCFL	YILKMSDDA	LFTYWNKAST	SEIMDFFTIS 480
481	EVCLHQFYQM	GKRYIARNOE	GLGPIVHDK	SQTLFVSRNR	TGAGHARLQQ	LGSLDNLTF	NHSYCHSDAD	VLHQSLLEAN 560
561	IATEVCLTAL	DTLSLFTLAF	KNQLLADHGH	NPLMKKVFVDV	YLCFLQKHQS	ETALQNVFTA	LRSLIYKFPS	TFYEGRADMC 640
641	AALCYEILKC	CNSKLSSIRT	EASQLLYFLM	RNNFDYTGKK	SEVRTHLQVI	ISVSQLIADV	VGIGETRFQQ	SLSIINNCAN 720
721	SDRLIKHTSF	SSDVKDLTKR	IRTVLMATAQ	MGEHENDPEM	LVDLQYSLAK	SYASTPELRK	TWLDSEARIE	VKNGDLSEAA 800
801	MCYVHVTAIV	AEYLTRKGVF	RQGCFAFRVI	TPNIDEEASH	MEDVGMQDVH	FNEDVLMELL	EQCADGLWKA	ERYELIADIY 880
881	KLIPIIYEKR	RDFEDEDGK	EYIYKEPKLT	PLSEISQRL	KLYSDKFGSE	NVKMIQDSCK	VNPKDLDSKY	AYIQVTHVIP 960
961	FFDEKELQER	KTEFERSHNI	RRFMFEMFFT	QTGKROGGVE	EQCKRRTILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID 1040
1041	EMSKKVAELR	QLCSSAEVDM	IKLQKLQGS	VSVQVNAGPL	AYARAFDDT	NTKRYFDNKV	KLLKEVFRQF	VEACQALAV 1120
1121	NERLIKEDQL	EYQEEMKANY	REMAKELSEI	MHEQICPLEE	KTSLVENSLE	IFNAISGTP	STAVHGMTSS	SSVV 1195

FIG. 10A (cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCAGTA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCCCC	GAGGTGACG	GTATCGATAA	GCTTGATATC 80
81	GAATTCGGCA	CGAGTTTAC	ACCATCACCA	AAACCCAGAA	TTTTATGATG	AGATTAATAA	AGAGTTGCCC	ACTCAGCTGC 160
161	ATGAAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTGAG	CTGTGACAA	TCAAGTAAAG	GAAGCACGAA	GAAGAGGGAT 240
241	GTCGTTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCCCTC	TGAAAGACGG	AAGGGTGGTG	ACAAGCGAGC	AGCACATCCC 320
321	GGTCTCGGGC	AACCTTCCTT	CGGGCTATCT	TGGGTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAATTTAAAT 400
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTTACATA 480
481	ATTTTTCCTA	GTACTGTCAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAGAGTCTG 560
561	CATGCGATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTC	CGAGTCTCTA	CCAGAGCCAC 640
641	ACAGGAAGAA	GTGCGGGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCACTGCCA	TGACGAAGGA	TTGGAGAGCC 720
721	ACTTGAGGTC	ATATGTAAAG	TACGCGTATA	AGGCTGAGCC	ATATGTGTCC	TCTGAATACA	AGACAGTGCA	TGAAGAAGTC 800
801	ACCAAATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCCTCACCAG	CAACAAACTA	CTGAGGTACT	CATGGTTTCT 880
881	CTTTGATGTA	CTGATCAAA	CTATGGCTCA	GCATTGATA	GAGAATCCCA	AAGTTAAGTT	GCTGCGAAAC	CAGAGATTTC 960
961	CTGCAATCCTA	TCATCATGCA	CGGGAACCGG	TTGTAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTGG	AGATAATCCA 1040
1041	ATAAGCATCTA	AGAACGCGAA	TCATAGCCTT	GCTGTCTTCA	TCAAGAGATG	TTTCACTTTC	ATGGACAGGG	GCCTTTGTCT 1120
1121	CAAGCAGATC	AACAACATCA	TTAGCTGTCT	TGCTCCTGGA	GACCCAAAGA	CCCTCTTTGA	ATACAGTTT	GAATTTCTCC 1200
1201	GTGTAGTGTG	CAACCATGAA	CATTATATTC	CGTTGAACTT	ACCAATGCCA	TTTGGAAAAG	GCAGGATTCA	AAGATACCAA 1280
1281	GACCTTCAGC	TTGACTACTC	ATTAACAGAT	GAGTTCTGCA	GAAACCACTT	CTTGGTGGGA	CTGTTACTGA	GGGAGGTGGG 1360
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATAAAG	CATTCTTTTG 1440
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACTCTCTAC	CTGCCTCTGT	TTGGTCTGCT	GATTGAAAAC 1520
1521	GTCCAGCGGA	TCAATGTGAG	GGATGTGTCA	CCCTTCCCTG	TGAACGCGGG	CATGACCGTG	AAGGATGAAT	CCCTGGTCTT 1600
1601	ACCAGCTGTG	AATCCGCTCG	TGACGCCGCA	GAAGGGAAGC	ACCCTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGGCGCCA 1680
1681	TCTCCGCGAT	TGCTTCTCCA	TATACAACTT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTTCAG	AGGATCTCTC 1760
1761	ATAAGCACAG	ATTCGGGTAA	CAGCCTTCCA	GAAAGGAATA	GTGAGAAAGG	CAATTCCCTG	GATAAGCACC	AACAAAGTAG 1840
1841	CACATTGGGA	AATTCGCTGG	TTGCTGTGTA	TAAACTTGAC	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCTCTTACA 1920
1921	TCCTTAAAGG	CATGTCTGAT	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACTTATGGA	TTTTTTTACA 2000
2001	TATCTGGAAG	TTCGCTGTGA	CCAGTCCGAG	TACATGGGGA	AGCCATACAT	AGCCAGGAAC	CAGGAGGGGT	TGGGACCCAT 2080
2081	AGTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTCCCGT	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG 2160
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAACC	ACAGCTATGG	CCACTCGGAC	GCAGATGTTT	TGCACCACTG	ATTACTTGAA 2240
2241	GCCAACTAGT	CTACTGAGGT	TTGCTGTGCA	GCTCTGGACA	GCCTTTCTCT	ATTTACATTC	CGGTTTAAAG	ACCGCTCTCT 2320
2321	GGCCGACCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGCTTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA 2400
2401	CGGCTTTAAA	AAATGTCTTC	ACTGCCTTAA	GGTCCCTTAA	TTATAAGTTT	CCCTCAACAT	TCTATGAAGG	GAGAGCGGAC 2480
2481	ATGTGTGCGG	CTCTGTGTTA	CGAGATTCTC	AAGTGTCTGA	ACTCCAAAGC	GAGCTCCATC	AGGACGGAGG	CCTCCAGACT 2560
2561	GCTCTACTTC	CTGATGAGGA	ACAACTTTGA	TTCACTGGA	AAGAAGTCTT	TTGTCCGGAC	ACATTTGCAA	GTCTCATCAT 2640
2641	CTGTAGAGCA	GCTGATAGCA	GACGTTGTGG	GCATTTGGGA	AACCAGATTG	CAGCAGTCCC	TGTCCATCAT	CAACAACCTG 2720
2721	GCACCAAGTG	ACCGGCTTAT	TAAGCACACC	AGCTTCTCCT	CTGATGTGAA	GGACTTAAAC	AAAAGGATAC	CCACGGCTCT 2800
2801	AATGGCCACC	GCCAGATGTA	AGGACATGTA	GAACGACCCA	GAGATGCTGG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT 2880
2881	ATGCCAGCAC	CGCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG 2960
2961	GCGCAATCTG	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	AATATCTCAC	ACGGAAAGGC	GTGTTTAGAC	GAAGATGCAC 3040
3041	CGCCTTCAGG	GTCAATTACC	CAAAACATCGA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCAGGAT	GTCCATTTCA 3120
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	AGTGCCGAGA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCCGCCGAC 3200
3201	ATCTTCAAAA	TTATCATCCC	CATTATGAG	AAGCCGAGGG	ATTCTTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTCATA 3280
3281	GGAAACCCAA	CTCACACCCG	TGTCGGAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGGT	TCTGAAAATG 3360
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCTTAGGGA	TCTGGATTCT	AAGTATGCA	ACATCCAGGT	GACTCACGCT 3440
3441	ATCCCTTCTT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCAC	AACATCCGCC	GCTTCATGTT 3520
3521	TGAGATGCCA	TTTACGCGAG	CCGGGAAGAG	CGAGGGCGGG	GTGGAAGAGC	AGTGCAAAAC	CGCACCACAT	CTGACAGCCA 3600
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCTGTCTAT	GTACCAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC 3680
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGCTCCTC	GGCCGAGGTG	GACATGATCA	AACATGAGCT 3760
3761	CAAACCTCCG	GGCAGCGTGA	GTGTTGAGGT	CAATGCTGGC	CCACTAGCAT	ATGCGCGAGC	TTTCTTAGAT	GATACAAACA 3840
3841	CRAAGCGATA	TCCTGACAA	AAAGTGAAGC	TGCTTAGGGA	AGTTTTCAGG	CAATTTGTGG	AAGCTTGGCG	TCAGGCTTTA 3920
3921	GCGGTAACAG	AACGTCTGAT	TAAAGAAGAC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACTACAGGG	AAATGGCGAA 4000
4001	GGAGCTTTCT	GAATCATGTC	ATGAGCAGAT	CTGCCCCCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTCC	CTTCACATCT 4080
4081	TCACGCCCAT	CAGTGGGACT	CCAACAAGCA	CAATGGTTCA	CGGGATGACC	AGCTCGTCTT	CGGTGCTGTG	ATTACATCTC 4160
4161	ATGGCCCGTG	TGTGGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA 4240
4241	CAGGGAGGAC	CAAGGGGAAG	GGGAGAGAAA	GGAATAAAG	AACAACGTTA	TTTCTTAAAC	GACTTTCTAT	AGGAGTTGTA 4320
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAGTTTTT	ATTGTGTCTT	AACAAAGGTG	TGGTAGACAC 4400
4401	TCCTTGAGCTG	GACTTAGATT	TTATTCTTCC	TTGCAAACT	GTGTTAGAA	AGATGGCCCTA	CAGAAAAAAA	AGGTTCTGGG 4480
4481	ATCTACATGG	CAGGAGGGGC	TGCACTGACA	TTGATGCTGT	GGGGACCTTT	TGCTCGGACT	CGTGCCGGA	ATCTGATCGT 4560
4561	AATCAGGGTA	CAGAACTTAC	TAGTTTTGTC	TAGGAGTATG	TTGTATGACT	AGGATTTGTC	CTATTATCTC	ATTCACCAAC 4640
4641	ATAGAGCAAG	AAATGTGAGC	TAACGTGAGC	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGAACT	TGTGATTTAA 4720
4721	TCATCGACTC	CGGGACCGTC	ATATATGTAT	TACATTTCTA	CATTTTAAAT	ACTCACATGG	GCTTATGCAT	TAAGTTTAAAT 4800
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGGTTTA	TTCTTGTGAT	AAAAATGTGC	AATATGGAGA 4880
4881	TGTATACAAG	TCTTTACT						4898

FIG.10B

	10	20	30	40	50	60	70	80
1	MEGHVMIAFL	PTILNQLFRV	LTRATQEEVA	VNVTRVIIHV	VAQCHEEGLE	SHLSYVKYA	YKAEFYVASE	YKTVHEELTK 80
81	SMTTILKPSS	DFLTSNKLLR	YSWFFFDVLI	KSMAQH LIEN	SKVKLLRNQR	FPASYHHAAE	TVVNMMLPHI	TQKFGDNPEA 160
161	SKNANHSLAV	FIKRCFTFMD	RGEFVEKQINN	YISCFAPGDP	KTLFEYKFEF	LRVVCNHEHY	IPLNLPMPFG	KGRIQRYQDL 240
241	QLDYSLTDEF	CRNHFLVGLL	LREVG TALQE	FREVRLIAIS	VLKNLLIKHS	FDDRYASRSH	QARIATLYLE	LFGLLIENVQ 320
321	RINVRDVSPF	PVNAGMTVKD	ESLALPAVNP	LVTPQKGSTL	DNSLHKDLLG	AISGIASPYT	TSTPNINSVR	NADSRGSLIS 400
401	TDSGNSLPER	NSEKSNSLDK	HQQSSTLGNS	VVRCDKLDQS	EIKSLIMCFL	YILKSMSDDA	LFTYWNKAST	SELMDDFTIS 480
481	EVCLHQFYM	GKRYIARNQE	GLGPIVHDRK	SQTLFVSRNR	TGMHARLQQ	LGSLDNSLTF	NHSYGHSDAD	VLHQSLLEAN 560
561	IATEVCLTAL	DTLSLFTLAF	KNQLLADHGH	NPLMKKVFDV	YLCFLQKHQS	ETALKNVFTA	LRS LIYKFPS	TFYEGRADMC 640
641	AALCYEILKC	CNSKLSSIRT	EASQILLYFLM	RNNFDYTGKK	SEVRTHLQVI	ISVSQLIADV	VGIGETRFOQ	SLSIINN CAN 720
721	SDRLIKHTSF	SSDVKOLTKR	IRTVLMATAQ	MKEHENDPEM	LVDLQYSLAK	SYASTPELRK	TWLD SMARIH	VKNGDLSEAA 800
801	MCYVHV TALV	AEYLTRKGVF	RQGC TAFRVI	TPNIDEEASM	MEDVGMQDVH	FNEDVLMELL	EQCADGLWKA	ERYELIADIY 880
881	KLIIP IYEKR	RDFFEDEDGK	EYIYKEPKLT	PLSEISQRLL	KLYSDKFGSE	NVKMIQDSGK	VNPKDLDSKY	AYIQVTHVIP 960
961	FEDEKELQER	KTEFERSHNI	RRFMFEMPFT	QTGKRQGGVE	EQCKRR TILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID 1040
1041	EMSKKVAELR	QLCSSAEVDM	IKLQKLQGS	VSVQVNAGPL	AYARAFLDDT	NTKRYPDNKV	KLLKEVERQF	VEACGQALAV 1120
1121	NERLIKEDQL	EYQEEMKANY	REMAKELSEI	MHEQICPLEE	KTSVLPNSLH	IFNAISGTPT	STMVHGMTSS	SSVV 1195

FIG. 10B (cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCACTA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCTC	GAGGTCGACG	GTATCGATAA	GCTTGATATC 80
81	GAATTGCGCA	CGAGTTTAC	ACCATCACCA	AAACCCAGAA	TTTTATGATG	AGATTAAAT	AGAGTTGCC	ACTCAGCTGC 160
161	ATGAAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTCAG	CTGTGACAAC	TCAAGTAAAG	GAAGCACGAA	GAAGAGGGAT 240
241	GTCGTTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCCTCC	TGAAAGACGG	AAGGGTGGTG	ACAAGCGGCG	AGCACATCCC 320
321	GGTCTCGGCG	AACCTTCCTT	CGGGCTATCT	TGGCTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAATTTAAAT 400
401	GGGTAGATCG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTATACATA 480
481	ATTTTTTCCA	GTACTGTCAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAGAGTCTG 560
561	CATGCGATCG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCTTAA	CCAGCTGTTT	CGAGTCTCTA	CCAGAGCCAC 640
641	ACAGGAAGAA	GTCCCGCTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TTGGAGAGCC 720
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAACTG 800
801	ACCAAATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCTCACCAG	CAACAACTA	CTGAGGTACT	CATGGTTTTT 880
881	CTTTGATCTA	CTGATCAAA	CTATCGCTCA	GCATTGATA	GAGAACTCCA	AAGTTAAGTT	GCTGCCAAAC	CAGAGATTTT 960
961	CTGCATCCTA	TCATCATCCA	CCGGAACCG	TTGTAAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTGG	AGATAATCCA 1040
1041	GAGGCATCTA	AGAACGGGAA	TCATAGCCTT	GCTGTCTTCA	TCAAGAGATG	TTTCACCTTC	ATGGACAGGG	GCTTTGTCTT 1120
1121	CAAGCAGATC	AACAACCTACA	TTAGCTGTTT	TGCTCCTGGA	GACCCAAAGA	CCCTCTTTGA	ATACAAGTTT	GAATTTCTCC 1200
1201	GTGTAGTCTG	CAACCATGAA	CATTATATTC	CGTTGAACCT	ACCAATAGCA	TTTGGAAAAG	GCAGGATTTA	AAGATACCAA 1280
1281	GACCTCCAGC	TTGACTACTC	ATTACAGAT	GAGTTCTGCA	GAAACCACTT	CTTGGTGGGA	CTGTTACTGA	GGGAGGTGCG 1360
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATAAAG	CATTCTTTTG 1440
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACCTCTTAC	CTGCCTCTGT	TTGGTCTGCT	GATTGAAAAC 1520
1521	GTCCAGCCGA	TCAATGTCAG	GGATGTGTCA	CCCTTCCCTG	TGAACCGCGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT 1600
1601	ACCAGCTGTG	AATCCGCTCG	TGACGCCGCA	GAAGGGAAGC	ACCTTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGGCGCCA 1680
1681	TCTCCGGCAT	TGCTTCTCCA	TATACAACCT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCGAG	AGGATCTCTC 1760
1761	ATAAGCACAG	ATTCGGGTAA	CAGCCTTCCA	GAAGGGAATA	GTGAGAAGAG	CAATTCCCTG	GATAAGCACC	AACAAAGTAG 1840
1841	CACATTGGGA	AATTCGCTGG	TTGCGTGTGA	TAAACTTGAC	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCCTCTACA 1920
1921	TCTTAAAGAG	CATGCTGAT	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACTTATGGA	TTTTTTTACA 2000
2001	ATATCTGAAG	TCTCCCTCCA	CCAGTTCCAG	TACATGCGGA	AGCGATACAT	AGCCAGGAAC	CAGGAGGAGT	TGGGACCCAT 2080
2081	AGTTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTCCCGT	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG 2160
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAAAC	ACAGTATGG	CCACTCGGAG	GCAGATGTTT	TGCACCAGTC	ATTACTTGAA 2240
2241	GCCAAACATT	TGCTGAGGT	TTGCCGTGACA	GCTCTGAGCA	CGCTTTCTCT	ATTACATTG	CGGTTTAAAG	ACCAGCTCCT 2320
2321	GGCCGACCAT	GCACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGTCTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA 2400
2401	CGGCTTTAAA	AAATGCTTTC	ACTGCCTTAA	GGTCTTAAAT	TTATAGATTT	CCCTCAACAT	TCTATGAAG	GAGAGCCGAC 2480
2481	ATGTGTGCGG	CTCTCTGTTA	CGAGATTCTC	AAGTCTGTGA	ACTCCAAGCT	GAGCTCCATC	AGGACGGAGG	CTGCCAGGCT 2560
2561	GCTCTACTTC	CTGATGAGGA	ACAACCTTTGA	TTACATGGA	AAGAAGTCTT	TTGTCCGGAC	ACATTTGCAA	GTCTCATAT 2640
2641	CTGTACGCCA	GCTGATAGCA	GACGTTGTTG	GCATTGGGGA	AACCAGATTG	CAGCAGTCCC	TGTCCATCAT	CAACACATGT 2720
2721	GCCAACAGTG	ACCGGCTTAT	TAAGCACACC	AGCTTCTCCT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTGCT 2800
2801	AATGGCCACC	CGCCAGATGA	AGGAGCATGA	GAACGACCCA	ATCGATCTGG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT 2880
2881	ATGCCAGCAC	CGCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCCTCAGAG 2960
2961	GCAGCAATGT	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	AATATCTCAC	ACGGAAAGGC	GTGTTTAGAC	AAGGATGCAC 3040
3041	CGCCTTCAGT	GTCATTACCC	CAACATCGCA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCAGGAT	GTCCATTCTA 3120
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	AGTGCACAGA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCGCCGAC 3200
3201	ATCTACAAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTACAA 3280
3281	GGAACCCAAA	CTCACACCGC	TGTCGGAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGTT	TCTGAAAATG 3360
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCCTAAGGA	TCTGGATTCT	AAGTATGCAT	ACATCCAGGT	GACTCACGTC 3440
3441	ATCCCCCTCT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCAC	AACATCCGCC	GCTTCATGTT 3520
3521	TGAGATGCCA	TTTACGCAGA	CCGGGAAGAG	GCAGGGCGGG	GTGGAAGAGC	AGTGCAAACG	GCGCACCATC	CTGACAGCCA 3600
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCTAT	TACACAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC 3680
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGCTCCTC	GGCCGAGGTG	GACATGATCA	AACTGCAGCT 3760
3761	CAAACCTCAG	GGCAGCGTGA	GTGTTTCAGGT	CAATGCTGGC	CCACTAGCAT	ATGCCGCGAG	TTTCTTAGAT	GATACAAACA 3840
3841	CAAGGCCGATA	TCCGTGACAAT	AAAGTGAAGC	TGCTTAAAGGA	AGTTTTCAGG	CAATTTGTGG	AAGCTTGGCG	TCAGGCCCTTA 3920
3921	GCGGTAAACG	AACGTCTGAT	TAAAGAAGAC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACTACAGGG	AAATGGCGAA 4000
4001	GGAGCTTTCT	GAATCATGCG	ATGAGCAGAT	CTGCCCCCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTCC	CTTCACATCT 4080
4081	TCAACGCCAT	CAGTGGGACT	CCAACAAGCA	CAATGGTTCA	CGGGATGACC	AGCTCGTCTT	CGGTCTGTGT	ATTACATCTC 4160
4161	ATGGCCCGTG	TGTGGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA 4240
4241	CAGGGAGGAC	CAAGGGGAAG	GGGAGAGAAA	GGAAATAAAG	AACAACGTTA	TTTCTTAACT	GACTTTCTAT	AGGAGTTGTA 4320
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAGTTTTT	ATTGTGTCTT	AACAAAGGTG	TGGTAGACAC 4400
4401	TCTTGAGCTG	GACTTAGATT	TTATTCTTCC	TTCGACAGTA	GCTTTAGAAT	AGATGCGCTA	CAGAAAAAAA	AGGTTCTGGG 4480
4481	ATCTACATGG	CAGGGAGGGC	TGCACTGACA	TTGATGCCTG	GGGGACCTTT	TGCCTCGACT	CGTGCCGGAA	ATCTGATCGT 4560
4561	AATCAGGGTA	CAGAACTTAC	TAGTTTTGTC	TAGGAGTATG	TTGTATGACT	AGGATTTGTG	CTATTATCTC	ATTCAACAAC 4640
4641	ATAGAGCAAG	AATAGTGAGC	TAACTGAGCT	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGAACT	TTGTCAATTAA 4720
4721	TCATCGACTC	CGGGACGGTC	ATATATGTAT	TACATTTCTA	CATTTTAAAT	ACTCACATGG	GCTTATGCAT	TAAGTTTAAAT 4800
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATCACTTT	TAATGGTTTA	TTCTTGTCAT	AAAAATGTGC	AAATATGGAGA 4880
4881	TGTATACAAG	TCTTTACT						

FIG.10C

		10		20		30		40			50		60		70		80	
1		MEGHVMI	AFL	PTILNQL	FRV	LTRATQEE	VA	VNVTRV	IIHV		VAQCHEE	GLE	SHLSYVK	YA	YKAE	PYVASE	YKTVHEELTK 80	
81		SMTTILK	PSA	DELT	SNKLLR	YSWFFFD	VLI	KSMAQH	LIE		SKVKLLR	NQR	FPASYH	HAAE	TVVNML	MPHI	TQKFGDNPEA 160	
161		SKNANH	SLAV	FIKRCF	TFMD	RGFVFK	QINN	YISCFAP	GDP		KTLFEYK	FEF	LRVVCN	HEHY	IPLNLP	MPFG	KGRIQRYQDL 240	
241		QLDYSL	TDEF	CRNHFL	VGLL	LREVGTA	LQE	FREVRL	IAIS		VLKNNLL	IKHS	FDDRYAS	RSH	QARIAT	LYLP	LFGLLIENVQ 320	
321		RINVRD	VSPF	PVNAGM	TVKD	ESLALPA	VNP	LVTPOK	GSTL		DNSLHKD	LLG	AISGIA	SPYT	TSTPNIN	SVR	NADSRGSLIS 400	
401		TDSGNS	LPER	NSEKSN	SLDK	HQQSSTL	GNS	VVRCDK	LDO		EIKSLLM	CFL	YILKSM	SDDA	LFTYWN	KAST	SELMDEFTTIS 480	
481		EVCLHQ	FQYM	GKRYIA	RNOE	GLGPIV	HDRK	SQTLEP	SRNR		TGMMHAR	LQO	LGSLDN	SLTF	NHSYGH	SDAD	VLHQSLLEAN 560	
561		IATEVCL	TAL	DTLSLF	TALF	KNQLLAD	HGH	NPLMKK	VFDV		YLCFLQ	KHQS	ETALKN	VFTA	LRSLIY	KFPS	TFYEGRADMC 640	
641		AALCYE	ILKC	CNSKLS	SIRT	EASQLLY	FLM	RNNFDY	TGKK		SFVRTHL	QVI	ISVSQLI	ADV	VGIGETR	FOO	SLSIINNCAN 720	
721		SDRLIK	HTSF	SSDVKDL	TKR	IRTVLM	ATAQ	MKEHEN	DPEM		LVDLQYS	LAK	SYASTPE	LK	TWLD	SMARIH	VKNGDLSEAA 800	
801		MCYVHV	TALV	AEYLTR	KGVF	RQGCTAF	RVI	TPNIDE	EASM		MEDVGM	QDVH	FNEDVLM	ELL	EQCADGL	LWKA	ERYELIADIY 880	
881		KLIPIY	EKR	RDFEDED	GK	EYIYKE	PKLT	PLSEIS	QRLL		KLYSDK	FGSE	NVKMIQ	DSGK	VNPKDL	DSKY	AYIQVTHVIP 960	
961		FFDEKE	LQER	KTEFERS	HNI	RRFMFEM	PFT	QTGKRQ	GGVE		EQCKRRT	ILT	AIHCFPY	VKK	RIPVMY	QHHT	DLNPIEVAID 1040	
1041		EMSKKV	AEELR	QLCSSAE	VDM	IKLQLK	LQGS	VSVQVN	AGPL		AYARAF	LDDT	NTKRYPD	NKV	KLLKEV	FRQF	VEACGQALAV 1120	
1121		NERLIK	EDQL	EYQEEM	KANY	REMAKEL	SEI	MHEQIC	PLEE		KTSVLP	NSLH	IFNAIS	GTPT	STMVHG	MTSS	SSV 1195	
			10		20		30		40			50		60		70		80

FIG. 10C (cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCACTA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCTC	GAGGTCGACG	GTATCGATAA	GCTTGATATC 80
81	GAATTCGGCA	CGAGTTTAC	ACCATCACCA	AAACCCAGAA	TTTATGATG	AGATTAAAAT	AGAGTTGCC	ACTCAGCTGC 160
161	ATGAAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTGAC	CTGTGACAAC	TCAAAGTAAG	GAAGCAGGAA	GAAGAGGGAT 240
241	GTCGTTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCCTCC	TGAAAGACGG	AAGGGTGGTG	ACAAGCGAGC	AGCACATCCC 320
321	GGTCTCGCGG	AACCTTCTCT	CGGGCTATCT	TGGGTACCAA	GAGCTTGGA	TGGGCAGGCA	TTATGGTCCG	GAATTTAAAT 400
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTTCATA 480
481	ATTTTTTCCA	GTACTGTCAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAGAGTCTG 560
561	CATGCGATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTC	CGAGTCCTCA	CCAGAGCCAC 640
641	ACAGGAAGAA	GTCGCGGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TGGAGAGGCC 720
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAAGTC 800
801	ACCAAATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCCTCACCAG	CAACAAACTA	CTGAGGTACT	CATGGTTTTT 880
881	CTTTGATGTA	CTGATCAAAAT	CTATGGCTCA	GCATTTGATA	GAGAAGTCCA	AAGTTAAGTT	GCTGCCGAAC	CAGAGATTTT 960
961	CTGCATCCTA	TCATCATGCA	GCGGAAACCG	TTGTAAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTGG	AGATAATCCA 1040
1041	GAGGCATCTA	AGAACGCGAA	TCATAGCCTT	GCTGCTTTCA	TCAAGAGATG	TTTCACCTTC	ATGGACAGGG	GCTTTGTCTT 1120
1121	CAAGCAGATC	AACAACATCA	TTAGCTGTTT	TGCTCCTGGA	GACCCAAAGA	CCCTCTTTGA	ATACAAGTTT	GAATTTCTCC 1200
1201	GTGTAGTGTG	CAACATGAA	CATTATATTC	CGTTGAACCT	ACCAATGCCA	TTTGGAAAAG	GCAGGATTTA	AAGATACCAA 1280
1281	GACCTCCAGC	TTGACTACTC	ATTAACAGAT	GAGTTCTGCA	GAAACCACTT	CTTGGTGGGA	CTGTTACTGA	GGGAGGTGGG 1360
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATAAAG	CATTCTTTTG 1440
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATGAC	CACCTCTCAC	CTGCCCTGCT	TTGGTCTGCT	GATTGAAAAC 1520
1521	TCCAGCCGGA	TCAATGTGAG	GGATGTGTCA	CCCTTCCCTG	TGAACGCGGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT 1600
1601	ACCAGCTGTG	AATCCGCTGG	TGACGCCGCA	GAAGGGAAGC	ACCTTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGGCGCCA 1680
1681	TCTCCGGCAT	TGCTTCTCCA	TATACAACCT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCGAG	AGGATCTCTC 1760
1761	ATAAGCACAG	ATTCGGGTAA	CAGCCTTCCA	GAAAGGAATA	GTGAGAAGAG	CAATTCCCTG	GATAAGCACC	AACAAAGTAG 1840
1841	CACATTGGGA	AATTCCGTGG	TTGCTGTGTA	TAACTTGAC	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCTCTTACA 1920
1921	TCTTAAAGAG	CATGCTGTAG	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACCTATGGA	TTTTTTTACA 2000
2001	ATATCTGAAG	TCTGCCTGCA	CCAGTTCCAG	TACATGGGGA	AGCGATACAT	AGCCAGGAAC	CAGGAGGGGT	TGGGACCCAT 2080
2081	AGTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTCCCGT	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG 2160
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAAAC	ACAGCTTATG	CCACTCGGAG	GCAGATGTTT	TGCACCAGTC	ATTACTTGAA 2240
2241	GCCACATTG	CTACTGAGGT	TTGCCCTGAC	GCTCTGGACA	CGCTTCTCT	ATTTACATTG	GCGTTTAAAG	ACCAGCTCCT 2320
2321	GGCCGACCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGCTTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA 2400
2401	CGGCTTTAAA	AAATGTCTTC	ACTGCCTTAA	GGTCCTTAAT	TTATAAGTTT	CCCTCAACAT	TCTATGAAGG	GAGAGCGGAC 2480
2481	ATGTGTGCGG	CTCTGTGTTA	CGAGATTCTC	AAGTGTCTGA	ACTCCAAGCT	GAGTCCATC	AGGACGGAGG	CTCCACGCT 2560
2561	GCTCTAGTTC	CTGATGAGGA	ACAACCTTGA	TTACACTGGA	AAGAAGTCC	TTGTCCGGAC	ACATTTGCAA	GTCTCATAT 2640
2641	CTGTGAGCCA	GCTGATAGCA	GACGTTGTTG	GCATTGGGGA	AACCAGATTG	CAGCAGTCCC	TGTCATCAT	CAACAACTGT 2720
2721	GCCAAACAGT	ACCGGCTTAT	TAAGCACACC	AGCTTCTCCT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTGCT 2800
2801	AATGGCCACC	GCCGAGATGA	AGGAGCATGA	GAACGACCCA	GAGATCTGCG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT 2880
2881	ATGCCAGCAC	GCCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG 2960
2961	GCAGCAATGT	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	AATATCTCAC	ACGGAAAGGC	GTGTTTAGAC	AAGGATGCAC 3040
3041	CGCCTTCAGG	GTCATTACCC	CAAACATCGA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCTTACG	GTCCATTCCA 3120
3121	ACGAGGATGT	CGTGTAGGAG	CTCCTTGAGC	AGTGGCGAGA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCGCCGAC 3200
3201	ATCTACAAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTCATA 3280
3281	GGAAACCAAA	CTCACACCGC	TGTCGGAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGGT	TCTGAAATG 3360
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCCTAAGGA	TCTGGATTCT	AAGTATGCA	ACATCCAGGT	GACTCACGTC 3440
3441	ATCCCTTTCT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCAC	AACATCCGCC	GCTTCATGTT 3520
3521	TGAGATGCCA	TTTACGCAGA	CCGGGAAGAG	GCAGGGCGGG	GTGGAAGAGC	AGTGCAAACG	GCGCACCATC	CTGACAGCCA 3600
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCTAT	GTACAGCAGC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC 3680
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGCTCCTC	GGCCGAGGTG	GACATGATCA	AACTGCAGCT 3760
3761	CAAACCTCCG	GGCAGCGTGA	GTGTTCAAGT	CAATGCTGGC	CCACTAGCAT	ATGCCGAGC	TTTCTTAGAT	GATACAAACA 3840
3841	CAAAGCGATA	TCTTGACAAT	AAAGTGAAGC	TGCTTAAGGA	AGTTTTTCAGG	CAATTGTGG	AAGCTTGGCG	TCAAGCCCTTA 3920
3921	GCGGTAAACG	AACGTCTGAT	TAAAGAAGAC	GAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACTACAGGG	AAATGGCGAA 4000
4001	GGAGCTTTCT	GAAATCATGC	ATGAGCAGAT	CTGCCCTCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTC	CTTCACATCT 4080
4081	TCAACGCCAT	CAGTGGGACT	CCAACAAGCA	CAATGGTTCA	CGGGATGACC	AGCTCGTCTT	CGGTGCTGTG	ATTACATCTC 4160
4161	ATGGCCCGTG	TGTTGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA 4240
4241	CAGGAGGAGC	CAAGGGGAAG	GGGAGAGAAA	GGAATAAAG	AACAACGTTA	TTTCTTAACA	GACTTTCTAT	AGGAGTTGTA 4320
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAGTTTTT	ATTGTGTCTT	AACAAGGTG	TGGTAGACAC 4400
4401	TCTTGAGCTG	CAGCTAGATT	TTATCTTTCC	TTGCAGAGTA	GTGTTAGAAT	AGATGGCCTA	CAGAAAAAAA	AGGTTCTGGG 4480
4481	ATCTACATGG	CAGGGAGGGC	TGCACTGACA	TTGATGCCTG	GGGACCTTTT	TGCCTCGACT	CGTGCCGGA	ATCTGATCGT 4560
4561	AATCAGGGTA	CAGAACTTAC	TAGTTTTGTC	TAGGAGTATG	TTGTATGACT	AGGATTGTGT	CTATTATCTC	ATTCAACAAC 4640
4641	ATAGAGCAAG	AATAGTGAGC	TAACTGAGCT	AGACACTCAA	TAAATCGGCT	ACTGGCTTCA	AGTCAGAACT	TGTGATTTAA 4720
4721	TCATCGACTC	CGGGACGGTC	ATATATGTAT	TACATTTCTA	CATTTTTTAAT	ACTCACATGG	GCTTATGCAT	TAAGTTTAAAT 4800
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGGTTTA	TTCTTGTCAT	AAAAATGTGC	AATATGGAGA 4880
4881	TGTATACAAG	TCTTTACT						4898

FIG.10D

	10	20	30	40	50	60	70	80
1	MEGHVMIAPL	PTILNQLFRV	LTRATQEEVA	VNVTRVIIHV	VAQCHEEGLE	SHLSYVKYA	YKAEPYVASE	YKTVHEELTK 80
81	SMTTILKPSA	DFLTSNKLLR	YSWFFFDVLI	KSMAQHLIEN	SKVKLLRNQR	FPASYHHAAE	TVVNMLMPHI	TQKFGDNPEA 160
161	SKNANHSLAV	FIKRCFTFMD	RGEVEKQINN	YISCFAPGDE	KTLFEYIGFEF	LRVVCNHEHY	IPLNLPMPFG	KGRIQRYQDL 240
241	QLDYSLTDEF	CRNHFLVGLL	LREVGTAQOE	FREVRLIAIS	VLKILLIKHS	FDDRYASRSH	QARIATLYLP	LFGLLIENVQ 320
321	RINVROVSPF	PVNAQMTVMD	ESLALPAVNP	LVTPOKGSTL	DNSLHKDILG	AISGIASPYT	TSTPNINSVR	NADSRGSLIS 400
401	TDSGNSLPER	MSEKSNSLDK	HQQSSTLGNS	VVRCDKLDQS	EIKSLIMCFL	YILKSMSDDA	LFTYWNKAST	SELMDEFTIS 480
481	EVCLHQFQTM	GCRYIARNOE	GLGPIVHDRK	SQTLFVSRNR	TGMMHARIQQ	LGSLDNLSTF	NHSYGHSDAD	VLHQSLLEAN 560
561	IATEVCLTAL	DTLSLFTLAF	KNQLLADHGH	NPLMKKVFEDV	YLCFLQKHQS	ETALKNVFTA	LRSLIYKFPS	TFYEGRADMC 640
641	AALCYEILKC	CNSKLSSTRT	EASQLLYFLM	RNNFDYTGGK	SFVRTHLQVI	ISVSQLIADV	VGIGETRFQQ	SLSIINNCAN 720
721	SDRLIKHTSF	SSDVKDLTKR	IRTVLMATAQ	MKEHENDPEM	LVDLQYSLAK	SYASTPELRK	TWLDSMARIH	VKNGDLSEAA 800
801	MCYVHVITALV	AEYLTRKGVF	RQCCTAFRVI	TENIDEEASM	MEDVGMQDVH	FNEDVIMELL	EQCADGLWKA	ERYELIADIY 880
881	KLIIPYIEKA	ADFFEDIECK	EYIYKEPKLT	PLSEISQRLI	KLYSDKFGSE	NVKMIQDSGK	VNPKDLDSKY	AYIQVTHVIP 960
961	FEDEKELQER	KTEPERSHN:	RPFMFEMPFT	QTGKRQGGVE	EQCKRRTILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID 1040
1041	EMSKKVAELA	QLC55AEVDM	IKLQLKLQGS	VSVQVNAGPL	AYARAFLDDT	NTKRYPDNKV	KLLKEVFRQF	VEACGQALAV 1120
1121	NERLIKEDQL	EYQEDMKANT	REMAKELSEI	MHEQICPLEE	KTSVLPNSLH	IFNAISGTPT	STMVHGMTSS	SSVV 1195

FIG. 10D(cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCACTA	TAGGGCGAAT	TGGGTACC	GGGCCCCCTC	GAGGTGACG	GTATCGATAA	GCTTGATATC
81	GAATTCGGCA	CGAGTTTTAC	ACCATCACCA	AAACCCAGAA	TTTATATGATG	AGATTAAAT	AGAGTTGCC	ACTCAGCTGC
161	ATGAAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTCAG	CTGTGACAA	TCAAGTAAAG	GAAGCAGAA	GAAGAGGGAT
241	GTCTTGA	CCCAAGTTGG	CTACTCCTGG	CTTCCCCTCC	TGAAAGACGG	AAGGGTGGTG	ACAAGCGAGC	AGCACATCCC
321	GGTCTCGCG	AACCTTCCTT	CGGGCTATCT	TGGCTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAATTTAAAT
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTTACATA
481	ATTTTTTCCA	GTACTGTGAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAAGAGTCTG
561	CATCCGATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTC	CGAGTCTCTCA	CCAGAGCCAC
641	ACAGGAAGAA	GTCCGCGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TTGGAGAGCC
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAAGTGC
801	ACCAAATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCCTCACCAG	CAACAACTA	CTGAGGTACT	CATGGTTTTT
881	CTTTGATGTA	CTGATCAAA	CTATGGCTCA	GCATTGTGATA	GAGAACTCCA	AAGTTAAGTT	GCTGCCGAAAC	CAGAGATTTT
961	CTGCATCCTA	TCATCATGCA	GCGGAAACCG	TTGTAAATAT	GCTGATGCCA	CACATCAGTC	AGAAGTTTGG	AGATAATCTA
1041	CAGGCATCTA	AGAAGCGGAA	TCATAGCCTT	GCTGTCTTCA	TCAAGAGATG	TTTCACTTTC	ATGGACAGGG	GCTTTGTCTT
1121	CATGATGATC	AACAACATCA	TTAGCTGTTT	TGCTCCTGGA	GACCCAAAGA	CCCTCTTTGA	ATACAAGTTT	GAATTTCTCC
1201	GTGTAGTGTG	CAACCATGAA	CATTATATTC	CGTTGAACTT	ACCAATGCCA	TTTGAAGAAC	GCAGGATTCA	AAGATACCAA
1281	GACCTCCAGC	TTGACTACTC	ATTAACAGAT	GAGTTCTGCA	GAACCACTTT	CTTGGTGGGA	CTGTTACTGA	GGGAGGTTGG
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGCTGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATAAAG	CATTCTTTTG
1441	ATCACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACCTCTTAC	CTGCCCTCTGT	TTGGTCTGCT	GATTGAAAGC
1521	GTCCAGCGCA	TCAATGTGAG	GGATGTGTCA	CCCTTCCCTG	TGAACGCGGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT
1601	ACCAGCTGTG	AATCCGCTGG	TGACGCGCGCA	GAAGGGAAGC	ACCCTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGCGCCCA
1681	TCTCCGCGAT	GTCTTCTCCA	TATACAACTT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCCAG	AGGATCTCTC
1761	ATAAGCACAG	ATTCGGGTAA	CAGCCTTCCA	GAAAGGAATA	GTGAGAAGAG	CAATTCCTCG	GATAAGCACC	AACAAAGTAG
1841	CACATTGGGA	AATTCGCTGG	TTCGCTGTGA	TAAACTTTAG	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCTCTTACA
1921	TCTTAAAGAG	CATGTCTGAT	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACTTATGGA	GAATTTTACA
2001	ATATCTGAAG	TCTGCCCTGCA	CCAGTTCCAG	TACATGGGGA	AGCGATACAT	AGCCAGGAAC	CAGGAGGGGT	TGGGACCCAT
2081	AGTTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTTCCCTG	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAAAC	ACAGCTATGG	CCACTCGGAC	GCAGATGTTT	TGCACAGATC	ATTAATTTGA
2241	GCCAACATTTG	CTACTGAGGT	TTGCCCTGACA	GCTCTGGACA	CGCTTTCTCT	ATTTACATTG	CGCTTTAAGA	ACCAGCTCCT
2321	GGCGGACCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGTCTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA
2401	CGGCTTTAAA	AAATGTCTTC	ACTGCCCTTAA	GGTCTCTTAA	TTATAAGTTT	CCCTCAACAT	TCTATGAAGG	GAGAGCGGAC
2481	ATGTGTGCGG	TCTGTGTTA	CGAGATTCTC	AAGTGTCTGA	ACTCCAAGCT	GAGCTCCATC	AGGACGGAGG	CCTCCAGCT
2561	GCTCTACTTC	CTGATGAGGA	ACAACTTTGA	TTACACTGGA	AAGAAGTCTC	TTGTCCGGAC	ACATTTGCAA	GTCAATCATAT
2641	CTGTGAGGCA	GCTGATAGCA	GACGTTGTTG	GCATTGGGGA	AACAGATTTC	CAGCAGTCCC	TGTCCATCAT	CAACAAGTGT
2721	CCCAACAGTG	ACCGGCTTAT	TAAGCACACC	AGCTTCTCCT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTCT
2801	AATGGCCACC	GCCAGATGA	AGGAGCATGA	GAACGACCCA	GAGATGCTGG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT
2881	ATGCCAGCAC	GCCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG
2961	GCAGCAATGT	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	AAATCTCTAC	ACGGAAAGGC	GTGTTTAGAC	AAGGATGCAC
3041	GCGCTTCAGG	CTATTATCCC	CAACATTCGA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCAGGAT	GTCCATTTC
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	AGTGCGCAGA	TGGAGTCTGG	AAAGCCGAGC	GCTACGAGCT	CATGCCCGAC
3201	ATCTACAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAGGAGCT	ATATTACAA
3281	GGAAACCCAA	CTCACACCGC	TGTCGGAAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGGT	TCTGAAATG
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCCTAAGGA	TCTGGATTCT	AAGTATGCAT	ACATCCAGGT	GACTCACGCT
3441	ATCCCTTTCT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCAC	AACATCCGCC	GCTTCATGTT
3521	TGAGATGCCA	TTTACGCGAG	CGGGAAGAG	CAGGGCGGG	GTGGAAGAGC	AGTGCAAAAC	GCGCACCATC	CTGACAGCCA
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCTAT	GTACCAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGCTFCTC	GGCCGAGGTG	GACATGATCA	AACTGCAGCT
3761	CAAACCTCCAG	GGCAGCGTGA	GTGTTCAAGT	CAATGCTGGC	CCACTAGCAT	ATGCGCGAGC	TTTCTTAGAT	GATACAAACA
3841	CAAAGCGATA	TCCTGACAA	AAAGTGAAGC	TGCTTAAGGA	AGTTTTTCAGG	CAATTTGTGG	AAGCTTGCGG	TCAAGCCTTA
3921	GCGGTAAACG	AACGTCTGAT	TAAAGAAGAC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACTACAGGG	AAATGGCGAA
4001	GGAGCTTTCT	GAATCATGTC	ATGAGCAGAT	CTGCCCTCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTCC	CTTCACATCT
4081	TCAACGCCAT	CAGTGGGACT	CCAACAAGCA	CAATGGTTCA	CGGGATGACC	AGCTCGTCTT	CGGTCTGTGT	ATTACATCTC
4161	ATGGCCCGTG	TGTGGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA
4241	CAGGAGGAGC	CAAGGGGAAG	GGGAGAGAAA	GGAAATAAAG	AACAACGTTA	TTTCTTAACA	GACTTTCTAT	AGGAGTTGTA
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAAGTTTC	ATTGTCTCTT	AACAAGGTG	TGGTAGACAC
4401	TCTTGAGCTG	GACTTAGATT	TTATTCTTCC	TTGCAGAGTA	GTGTTAGAA	AGATGGCCTA	CAGAAAAAAA	AGTTTCTGGG
4481	ATCTACATGG	CAGGGAGGGC	TGCACTGACA	TTGATGCTG	GGGGACCTTT	TGCTTCGACT	CGTGCCGGAA	ATCTGATCGT
4561	AATCAGGGTA	CAGAACTTAC	TAGTTTTGTC	TAGGAGTATG	TTGATGCTG	AGGATTGTG	CTATTATCTC	ATTCAACAAC
4641	ATAGAGCAAG	AATAGTGAGC	TAACTGAGCT	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGAACT	TTGTCAATTA
4721	TCATCGACTC	CGGACCGGTC	ATATATGTAT	TACATTTCTA	CATTTTTAAT	ACTCACATGG	GCTTATGCAT	TAAAGTTAAT
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGGTTTA	TTCTTGTCAT	AAAAATGTGC	AATATGGAGA
4881	TGTATACAAG	TCTTTACT						4898
	10	20	30	40	50	60	70	80

FIG.10E

	10	20	30	40	50	60	70	80					
1	MEGHVMI	AF	PTILNQL	FRV	LTRATQ	EEVA	VNVTRV	IIHV	VAQCHEEGLE	SHLSYVKYA	YKAEPYVASE	YKTVHEELTK	80
81	SMTTILK	PSA	DELT	SNKLLR	YSWFFD	VLI	KSMAQH	LIEN	SKVKILRNQR	FPASYHAAE	TVVNMLMPHI	TQKFGDNPEA	160
161	SKNANH	SLAV	FIKRCF	TFMD	RGFVFK	QINN	YISCFAP	GDP	KTLEFYKFEF	LRVVCNHEHY	IPLNLPMPFG	KGRIQRYQDL	240
241	QLDYSL	TDEF	CRNHFL	VGLL	LREVGTA	LQE	FREVRL	IAIS	VLKNLLIKHS	FDDRYASRSH	QARIATLYLP	LFGLLIENVO	320
321	RINVRD	VSPF	PVNAGM	TVKD	ESLALPA	VNP	LVTPQK	GSTL	DNSLHKDLLG	AISGIASPYT	TSTPNINSVR	NADSRGSLIS	400
401	TDSGNS	LPER	NSEKSN	SLDK	HQQSST	LGN	VVRCDK	LQDS	EIKSLIMCFL	YILKSMSDDA	LFTYWNKAST	SELMDDFTIS	480
481	EVCLHQ	FQYM	GKRYIAR	NQE	GLGPIV	HDRK	SQTLFV	SRNR	TGMHARLQQ	LGSLDNSLTF	NHSYGHSDAD	VLHQSLLEAN	560
561	IATEVCL	TAL	DTLSLE	TLAF	KNQLLAD	HGH	NPLMKK	VFDV	YLCFLQKHQS	ETALKNVFTA	LRSLIYKFPS	TFYEGRADMC	640
641	AALCYE	ILKC	CNSKL	SSIRT	EASQLLY	FLM	RNNFDY	TGKK	SFVRTHLQVI	ISVSQLIADV	VGIGETRFQQ	SLSIINNCAN	720
721	SDRLIK	HTSF	SSDVKD	LTKR	IRTVLM	ATAQ	MKEHEND	PEM	LVDLQYSLAK	SYASTPELRK	TWLDSMARIH	VKNGDLSEAA	800
801	MCYVHV	TALV	AEYLTR	KGVF	RQGC	TAFRVI	TPNIDEE	EASM	MEDVGMQDVH	FNEDVIMELL	EQCADGLWKA	ERYELIADIY	880
881	KLIIPY	EYKR	RDFEDED	GK	EYIYKE	PKLT	PLSEIS	QRLL	KLYSDKFGSE	NVKMIQDSGK	VNPKDLDSKY	AYIQVTHVIP	960
961	FFDEKE	LQER	KTEFERS	SHNI	RRFMF	FEMPFT	QTGKRQ	GGVE	EQCKRRTILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID	1040
1041	EMSKKV	AEALR	QLCSSA	EVDM	IKLQLK	LQGS	VSVQVN	AGPL	AYARAFLDDT	NTKRYPDNKV	KLLKEVFRQF	VEACQALAV	1120
1121	NERLIK	EDQL	EYQEMK	ANY	REMAKEL	SEI	MHEQIC	PLEE	KTSVLPNSLH	IFNAISGTPT	STMVHGMTSS	SSV	1195

FIG. 10E (cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCACA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCTC	GAGGTCGACG	GTATCGATAA	GCTTGATATC
81	GAATTCGGCA	CGAGTTTTC	ACCATCACCA	AAACCCAGAA	TTTTATGATG	AGATTAAAT	AGAGTTGCC	ACTCAGCTGC
161	ATGAAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTGAC	CTGTGACAA	TCAAGTAAAG	GAAGCAGGAA	GAAGAGGGAT
241	GTCGTTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCTTCC	TGAAAGACGG	AAGGGTGGTG	ACAAGCGAGC	AGCACATCCC
321	GGTCTCGGCG	AACCTTCCTT	CGGGCTATCT	TGGCTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAATTAAT
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTACATA
481	ATTTTTCCTA	GTACTGTGAC	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAGAGTCTG
561	CATGCGATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTC	CGAGTCCTCA	CCAGAGCCAC
641	ACAGGAAGAA	GTCGCGGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TTGGAGAGCC
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAACTG
801	ACCAATATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCTTCACCAG	CAACAACTA	CTGAGGTACT	CATGTTTTTT
881	CTTTGATGTA	CTGATCAAAT	CTATGGCTCA	GCATTTGATA	GAGAACTCCA	AAGTTAAGTT	GCTGCGAAAC	CAGAGATTTT
961	CTGCACTCTA	TCATCATGCA	GCGGAAACCG	TTGTAAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTTG	AGATAATCCA
1041	GAGGCATCTA	AGAACCGGAA	TCATAGCCTT	GCTGTCTTCA	TCAAGAGATG	TTTCACCTTC	ATGGACAGGG	GCTTTGTCTT
1121	CAAGCAGATC	AACAACTACA	TTAGCTGTTT	TGCTCCTGGA	GACCCAAAGA	CCCTCTTTGA	ATACAAAGTT	GAATTTCTCC
1201	GTGTAGTGTG	CAACCATGAA	CATTATATTC	CGTTGAACCT	ACCAATGCCA	TTTGGAAAAG	GCAGGATTTCA	AAGATACCAA
1281	GACCTCCAGC	TTGACTACTC	ATTAACAGAT	GAGTCTGTCA	GAAACCACTT	CTTGGTGGGA	CTGTTACTGA	GGGAGGTGGG
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACTT	GCTGATAAAG	CATTCTTTTT
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACCCCTCAG	CTGCCCTGTG	TTGGTCTGCT	GATTGAAAAC
1521	GTCCAGCGGA	TCAATGTGAG	GGATGTGTCA	CCCTTCCCTG	TGAACGCGGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT
1601	ACCAGCTGTG	AATCCGCTGG	TGACGCCGCA	GAAGGGAAGC	ACCCTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGCGGCCA
1681	TCTCCGGCAT	TGCTTCTCCA	TATACAACTT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCGAG	AGGATCTCTC
1761	ATAAGCAGAG	ATTCGGGTAA	CAGCCTTCCA	TACATGAACT	GTGAGAAAGG	CAATTCCCTG	GATAAGCACC	AACAAGTAG
1841	CACATTGGGA	AATTCCTGTG	TTCCGTGTGA	TAACTTGAC	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCTCTTACA
1921	TCTTAAAGAG	CATGCTGTAT	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACTTATGGA	TTTTTTTACA
2001	ATATCTGAAG	TCTGCCTGCA	CCAGTTCCAG	TACATGGGGA	AGCGATATAC	AGCCAGGAAC	CAGGAGGGGT	TGGGACCCAT
2081	AGTTTATGAT	AGACATTTCC	TGTTTCCCGT	TTTCTCCCTG	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG
2161	GCAGCCTGGA	TAACCTCTCT	ACTTTTAAAC	ACAGCTATGG	CCACTCCGAC	GCAGATGTC	TGCACAGGAT	ATTACTTGAA
2241	GCCAAACATT	CTACTGAGGT	TTGCCCTGCA	GCTCTGGACA	CGCTTTCTCT	ATTTACATTG	CGCTTTAAGC	ACCAGCTCCT
2321	GGCCAGCCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGTCTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA
2401	CGGCTTTAAA	AAATGCTTTC	ACTGCCCTAA	GGTCTTAAAT	TTATAAGTTT	CCCTCAACAT	TCTATGAAGG	GAGAGCGGAC
2481	ATGTGTGCGG	CTCTGTGTTA	CGAGATTCTC	AAGTGCTGTA	ACTCCAAGCT	GAGCTCCATC	AGGACGGAGG	CCTCCCAGCT
2561	GCTCTACTTC	CTGATGTTGA	ACAACCTTGA	TTACATGGGA	TCTGGATTCT	TTGTCCGGAC	ACATTTGCAA	GTCCATCATAT
2641	CTGTGAGCCA	GCTGATAGCA	GACGTTGTTG	GCATTTGGGA	AACCAGATTC	CAGCAGTCCC	TGTCCATCAT	CAACAACGTG
2721	GCCAAACAGT	ACCGGCTTAT	TAAGCAGACC	AGCTTCTCTT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTGCT
2801	AATGGCCACC	GCCAGATGTA	AGGAGCATGA	GAACGACCCA	GAGATGCTGG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT
2881	ATGCCAGCAC	CGCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG
2961	GCAGCAATGT	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	TTATATCTAC	ACGGAAGGCG	GTGTTTAGAC	AAGGATGCAC
3041	CGCCTTCAGG	GTCATTACCC	CAAAACATCA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCAGGAT	GTCCATTCTA
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	CGCTGCGAGA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCGCCGAC
3201	ATCTACAAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTTACA
3281	GAACCCAAA	CTCACACCGC	TGTCGGAAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGGT	TCTGAAAATG
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCTTAAGGA	TCTGGATTCT	AAGTATGCAT	ACATCCAGGT	GACTCACGTC
3441	ATCCCCCTCT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCAC	AACATCCGCC	GCTTCATGTT
3521	TGAGATGCCA	TTTACGCGAA	CCGGGAAGAG	GCAGGGCGGG	GTGGAAGAGC	AGTGCAAAGC	GCGCACCATC	CTGACAGCCA
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCTAT	GTACCAAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGCTCCTC	GGCCGAGGTG	GACATGATCA	AACTGCAGCT
3761	CAAACTCCAG	GGCAGCGTGA	GTGTTCAAGT	CAATGCTGGC	CCACTAGCAT	ATGCGCGAGC	TTTCTTAGAT	GATACAAACA
3841	CAAAAGCGATA	TCCTGACAAAT	AAAGTGAAGC	TGCTTAAGGA	AGTTTTCTAG	CAATTTGTGG	AAGCTTGGCG	TCAAGCCTTA
3921	CGGGTAAACG	AACGCTCTGAT	TAAAGAAAGC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACTACAGGG	AAATGGCGAA
4001	GGAGCTTTCT	GAAATCATGC	ATGAGCAGAT	CTGCCCTCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTCC	CTTCACATCT
4081	TCAACGCCAT	CAGTGGGACT	CCAACAAGCA	CAATGGTTCA	CGGATGAGCC	AGCTCGTCTT	CGGTCTGTGT	ATTACATCTC
4161	ATGGCCCGTG	TGTGGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	AGACTGGGG	AGACCGAGCA
4241	CAGGGAGGAC	CAAGGGGAAG	GGGAGAGAAA	GGAATAAAG	AACAACGTTA	TTTCTTAACA	GACTTTCTAT	AGGAGTTGTA
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAAGTTTT	ATTGTGTCTT	AACAAAGGTG	TGGTAGACAC
4401	TCTTGAGCTG	GACTTAGATT	TTATTCTTCC	TTCAGAGTGA	GTGTTAGAAT	AGATGGCCTA	CAGAAAAAAA	AGGTTCTGGG
4481	ATCTACATGG	CAGGGAGGGC	TGCACTGACA	TTGATGCCTG	GGGGACCTTT	TGCCCTGACT	CGTGCCGGAA	ATCTGATCGT
4561	AATCAGGGTA	CAGAATTTAC	TAGTTTTGTC	TAGGAGTATG	TTGTATGATC	AGGATTTTGT	CTATTATCTC	ATTCAACAAC
4641	ATAGAGCAAG	AATAGTGAGC	TAACTGAGCT	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGACT	TTGTCATTAA
4721	TCATCGCAAG	CAAGGACGTC	ATATATGTAT	TACATTTCTA	CATTTTAAAT	ACTCACATGG	GCTTATGCAAT	TAAGTTTAAAT
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGTTTTA	TTCTGTGCAT	AAAATGTGTC	AATATGGAGA
4881	TGTATACAAG	TCTTTACT						4898

FIG.10F

	10	20	30	40	50	60	70	80	
1	MEGHVMI AFL	PTILNQLFRV	LTRATQEEVA	VNVTRVIIHV	VAQCHEEGLE	SHLSYVKYA	YKAEPYVASE	YKTVHEELTK	80
81	SMTTILKPSA	DELTSTNKLRL	YSWFFFDVLI	KSMAQHLLIEN	SKVKLLRNQR	FPASYHHAAE	TVVNMLMPHI	TQKFGDNPEA	160
161	SKNANHSLAV	FIKRCFTFMD	RGFVFKQINN	YISCFAPGDP	KTLEFYKFEF	LRVVCNHEHY	IPINLPMFFG	KGRIQRYQDL	240
241	QLDYSLTDEF	CRNHFLVGLL	LREVGTLAQE	FREVRLIAIS	VLKNLLIKHS	FDDRYASRSH	QARIATLYLP	LFGLLIENVQ	320
321	RINVRDVSPF	PVNAGMTVKD	ESLALPAVNP	LVTPQKGSTL	DNSLHKDLIG	AISGIASPYT	TSTPNINSVR	NADSRGSLIS	400
401	TDSQNSLPER	NSEKSNSLSDK	HQSSSTLGNS	VVRCDKLDQS	EIKSLLMCFL	YILKSMSDDA	LFTYWNKAST	SELMDEFTIS	480
481	EVCLHQFQYM	GKRYIARNQE	GLGPIVHDK	SQTLFVSRNR	TGMHARLQQ	LGSLDNSLTF	NHSYGHSDAD	VLHQSLLEAN	560
561	IATEVCLTAL	DTLSLFTLAF	KNQLLADHGH	NPLMKKVFDV	YLCFLQKHQS	ETALKNVFTA	LRSLIYKFPS	TFYEGRADMC	640
641	AALCYEILKC	CNSKLSSIRT	EASQLLYFLM	RNNFDYTGKK	SFVRTHLQVI	ISVSQLIADV	VGIGETREFQ	SLSIINNCAN	720
721	SDRLIKHTSF	SSDVKDLTKR	IRTVLMATAQ	MKEHENDPEM	LVDLQYSLAK	SYASTPELRK	TWLDSMARIH	VKNGDLSEAA	800
801	MCYVHVLTALV	AEYLTRKGVF	RQGCTAFRVI	TPNIDEEASM	MEDVGMQDVH	FNEDVIMELL	EQCADGLWKA	ERYELIADIY	880
881	KLIIPYIEKR	RDFEDEDGK	EYIYKEPKLT	PLSEISQRLL	KLYSDKFGSE	NVKMIQDSGK	VNPKDLDSKY	AYIQVTHVIP	960
961	FFDEKELQER	KTEFFERSHNI	RRFMFEMPFT	QTGKRQGGVE	EQCKRRTILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID	1040
1041	EMSKKVAELR	QLCSSAEVDM	IKLQLKLQGS	VSVQVNAGPL	AYARAFLDDT	NTKRYPDNKV	KLLKEVFRQF	VEACGQALAV	1120
1121	NERLIKEDQL	EYQEEMKANY	REMAKELSEI	MHEQICPLEE	KTSVLENSLH	IFNAISGTP	STMVHGTMSS	SSVV	1195
	10	20	30	40	50	60	70	80	

FIG. 10F (cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCACTA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCTC	GAGGTCGACG	GTATCGATAA	GCTTGATATC
81	GAATTCGGCA	CGAGTTTAC	ACCATCACCA	AAACCCAGAA	TTTTATGATG	AGATTAAAAA	AGAGTTGCC	ACTCAGCTGC
161	ATGAAAAGCA	CCACCTGTTG	CTCACATTCT	TCCATGTGAG	CTGTGACAAC	TCAAGTAAAG	GAAGCAGGAA	GAAGAGGGAT
241	GTCGTTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCTCCT	TGAAAGACGG	AAGGGTGGTG	ACAAGCGAGC	AGCACATCCC
321	GGTCTCGGCG	AACCTTCCTT	CGGGCTATCT	TGGCTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAAATTTAAAT
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCGTGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTTACATA
481	ATTTTTTCCA	GTACTGTCAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAGAGTCTG
561	CATGCGATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTC	CGAGTCTCA	CCAGAGCCAC
641	ACAGGAAGAA	GTCGCGGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TTGGAGAGCC
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAACTG
801	ACCAAAATCCA	TGACCCAGAT	TCTCAAGCCT	TCTGCCGATT	TCCTCACGAG	CAACAAACTA	CTGAGGTACT	CATGGTTTTT
881	CTTTGATGTA	CTGATCAAAT	CTATGGCTCA	GCATTGATA	GAGAATCCCA	AAGTTAAGTT	GCTCGCAAAC	CAGAGATTTC
961	CTGCATCCTA	TCATCATGCA	GCGGAACCG	TTGTTAAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTGG	AGATAATCCA
1041	GAGGCATCTA	AGAACGCGAA	TCATAGCCTT	GCTGTCTTCA	TCAAGAGATG	TTTCACCTTC	ATGGACAGGG	GCTTTGTCTT
1121	CAAGCAGATC	AACAACATCA	TTAGCTGTTT	TGCTCCTGGA	GACCCAAGA	CCCTCTTTGA	ATACAAGTTT	GAATTTCTCC
1201	GCTGTAGTGT	CAACCATGAA	CATTATATTC	CGTTGAACCT	ACCAATGCCA	TTTGGAAGAG	GCAGGATTCA	AAGATACCAA
1281	GACCTCCAGC	TTGACTACTC	ATTAACAGAT	GAGTTCTGCA	GAAACCACTT	CTTGGTGGGA	CTGTACTGA	GGGAGGTGGG
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATAAAG	CATTCTTTTG
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACCCCTAC	CTGCCCTCTG	TTGGTCTGCT	TTCCGAAAC
1521	GTCCAGCGGA	TCAATGTGAG	GGATGTGTCA	CCCTTCCCTG	TGAACGCGGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT
1601	ACCAGCTGTG	AATCCGCTGG	TGACGCCGCA	GAAGGGAAGC	ACCCTGGACA	ACAGCCTGCA	CAAGGACCTG	CTGGGCGCCA
1681	TCTCCGGCAT	TGCTTCTCCA	TATACAACCT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCGAG	AGGATCTCTC
1761	ATAAGCACAG	ATTCCGGTAA	CAGCCTTCCA	GAAAGGAATA	GTGAGAAGAG	CAATTCCCTG	GATAAGCACC	AACAAAGTAG
1841	CACATTGGGA	AATTCCGTGG	TTCGCTGTGA	TAAACTTGAC	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCTCTCTACA
1921	TCTTAAAGAG	CATGCTGTAT	GATGCTTTGT	TTACATATTG	GAACAAGGCT	TCAACATCTG	AACTTATGGA	TTTTTTTACA
2001	ATATCTGAAG	TCTGCCTGCA	CCAGTTCCAG	TACATGGGGA	AGCGATACAT	AGCCAGGAAC	CAGGAGGGGT	TTGGAGCCAT
2081	AGTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTCCCGT	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAAAC	ACAGCTATGG	CCACTCGGAC	GCAGATGTTT	TGCACCACTG	ATTACTTGAA
2241	GCCAACATTG	CTACTGAGGT	TGCGCTGACA	GCTCTGGACA	CGCTTTCTCT	ATTTACATTG	GCGTTTAAAG	ACCAGCTCCT
2321	GGCCGACCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGTCTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA
2401	CGGCTTTAAA	AAATGTCTTC	ACTGCCTTAA	GGTCCTTAAT	TTATAGAGTT	CCCTCAACAT	TCTATGAAGG	GAGAGCGGAC
2481	ATGTGTGCGG	CTCTGTGTTA	CGAGATTCTC	AAGTGTGFTA	ACTCCAAGT	GAGCTCCATC	AGGACGGAGG	AGTCCCAAGT
2561	GCTCTACTTC	CTGATGAGGA	ACAACCTTGA	TTACACTGGA	AAGAAGTCCT	TTGTCCGGAC	ACATTGCAA	GTCATCATAT
2641	CTGTCAGCCA	GCTGATAGCA	GACGTTGTTG	GCATTGGGGA	AACCAGATTG	CAGCAGTCCC	TGTCCATCAT	CAACAACCTG
2721	GCCAACAGTG	ACCGGCTTAT	TAAGCACACC	AGCTTCTCCT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTCTG
2801	AATGGCCACC	GCCAGATGTA	AGGAGCATGA	GAACGACCCA	GAGATCTGGG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT
2881	ATGCCAGCAC	GCCCGAGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG
2961	CGAGCAATGT	GCTATGTCTA	CGTAACAGCC	CTAGTGCCAG	AATATCTCAC	ACGGAAAGGC	GTGTTTAGAC	AAGGATGCAC
3041	CGCCTTTCAGG	TCATTATCCC	CAAACATCGA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCAGGAT	GTCCATTTC
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	AGTGCCGACA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCGCCGAC
3201	ATCTACAAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTACAA
3281	GGAAACCCAA	CTCACACCCG	TGTCGGAAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGGT	TCTGAAAATG
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCCTAAGGA	TCTGGATTCT	AAGTATGCAT	ACATCCAGGT	GACTCACGTC
3441	ATCCCTTTCT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCAC	AACATCCGCC	GCTTCATGTT
3521	TGAGATGCCA	TTTACGCAGA	CCGGGAAGAG	GCAGGGCGGG	GTGGAAGAGC	AGTGCAACG	GCGCACCATC	CTGCACGCCA
3601	TACTACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCTAT	TGACGAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCCGGCAGC	TGTGCTCCTC	GGCCGAGGTG	GACATGATCA	AAGTGCAGCT
3761	CAAACTCCAG	GGCAGCGTGA	GTGTTTCTG	CAATGCTGGC	CCACTAGCAT	ATGCGCGAGC	TTTCTTAGAT	GATACAAACA
3841	CAAGCGGATA	TCCTGACAAAT	AAAGTGAAGC	TGCTTAAAGGA	AGTTTTTCAGG	CAATTTGTGG	AAGCTTGGCG	TCAAGCCTTA
3921	GCGGTAAACG	AACGTCTGAT	TAAAGAAGAC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACATCAGGG	AAATGGCGAA
4001	GGAGCTTTCT	GAAATCATGC	ATGAGCAGAT	CTGCCCTCTG	GAGGAGAGAA	CGAGCGTCTT	ACCGAATTCG	CTTCACATCT
4081	TCAACGCCAT	CAGTGGGACT	CCAAACAGCA	CAATGGTTCA	CGGGATGACC	AGCTCGTCTT	CGGTCTGTGT	ATTACATCTC
4161	ATGGCCCGTG	TGTGGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA
4241	CAGGAGGAGC	CAAGGGGAAG	GGGAGAGAAA	GGAAATAAAG	AACAACGTTA	TTTCTTAACA	GACTTTCTAT	AGGAGTTGTA
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAGTTTTC	ATTGTGCTTT	AACAAGGTG	TGGTAGACAC
4401	TCTTGAGCTG	CAGTTAGATT	TTATTCTTCC	TCTGCAGATG	GTGTTAGAAAT	AGATGGCCCTA	CAGAAAAAAA	AGGTTCTGGG
4481	ATCTACATGG	CAGGAGGGGC	TGCACTGACA	TTGATGCTTG	GGGGACCTTT	TGCTTCGACT	CGTGCCGGA	ATCTGATCGT
4561	AATCAGGGTA	CAGAACTTAC	TAGTTTTGTC	TAGGAGTATG	TTGTATGACT	AGGATTTGTG	CTATTATCTC	ATTCAACAAC
4641	ATAGAGCAAG	AATAGTGAGC	TAACTGAGCT	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGAACT	TTGTCATTAA
4721	TCATCGACTC	CGGGACGGTC	ATATATGTAT	TACATTTCTA	CATTTTTTAA	ACTCACATGG	GCTTATGCAT	TAAGTTTAAAT
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGGTTTA	TTCTTGTCAT	AAAAATGTGC	AAATATGGAGA
4881	TGTATACAAG	TCTTTTACT						4898

FIG.10G

	10	20	30	40	50	60	70	80	
1	MEGHVMI AFL	PTILNQLFRV	LTRATQEEVA	VNVTRVIIHV	VAQCHEEGLE	SHLSYVKYA	YKAEPYVASE	YKTVHEELTK	80
81	SMTTILKPSA	DELTSNKLRL	YSWFFFDVLI	KSMAQH LIEN	SKVKLLRNQR	FPASYHHAAE	TVVNMLMPHI	TQKFGDNPEA	160
161	SKNANHSLAV	FIKRCFTFMD	RGEVFKQINN	YISCFAPGDP	KTLFEYKFEF	LRVVCNHEHY	IPLNLMPFPG	KGRIQRYQDL	240
241	QLDYSLTDEF	CRNHFLVGLL	LREVGTALQE	FREVRLIAIS	VLKNLLIKHS	FDDRYASRSH	QARIATLYLP	LFGLLIENVQ	320
321	RINVRDVSPF	PVNAGMTVKD	ESLALPAVNP	LVTPQKGSTL	DNSLHKDILG	AISG IASPYT	TSTPNINSVR	NADSRGSLIS	400
401	TDSGNSLPER	NSEKSNSLDK	HQQSSTLGNS	VVRCDKLDQS	EIKSLIMCFL	YILKMSDDA	LFTYWNKAST	SELMDEFTIS	480
481	EVCLHQFYQM	GKRYIARNQE	GLGP IVHDRK	SQTLPVSRNR	TGMHARLQQ	LGSLDNSLTF	NHSYGHSDAD	VLHQSLLEAN	560
561	IATEVCLTAL	DTLSLFTLAF	KNQLLADHGH	NPLMKKVFDV	YLCFLQKHQS	ETALKNVFTA	LRSLIYKFPS	TFYEGRADMC	640
641	AALCYEILKC	CNSKLS SIRT	EASQLLYFLM	RNNFDYTGKK	SFVRTHLQVI	ISVSQLIADV	VGIGETREFO	SLSIINN CAN	720
721	SDRLIKHTSF	SSDVKDLTKR	IRTVLMATAQ	MKEHENDPEM	LVDLQYSLAK	SYASTPELRK	TWLD SMARIH	VKNGDLSEAA	800
801	MCYVHV TALV	AEYLTRKGVF	RQGCTAFRVI	TPNIDEEASM	MEDVGMQDVH	FNEDVLMELL	EQCADGLWKA	ERYELIADIY	880
881	KLIPIYIEKR	RDFEDEDGK	EYIYKEPKLT	PLSEISQRLI	KLYSDKFGSE	NVKMIQDSGK	VNEPKDLDSKY	AYIQVTHVIP	960
961	FFDEKELQER	KTEFERSHNI	RRFMFEMPFT	QTGKRQGGVE	EQCKRRTILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID	1040
1041	EMSKKVAELR	QLCSSAEVDM	IKLQLKIQGS	VSVQVNAGPL	AYARAFLDDT	NTKRYPDNKV	KLLKEVFRQF	VEACQOALAV	1120
1121	NERLIKEDQL	EYQEEMKANY	REMAKELSEI	MHEQICPLEE	KTSVLPNSLH	IFNAISGTPT	STMVHGMTSS	SSVV	1195
	10	20	30	40	50	60	70	80	

FIG. 10G (cont.)

	10	20	30	40	50	60	70	80
1	AATTGTAATA	CGACTCATA	TAGGGCGAAT	TGGGTACCGG	GCCCCCCTC	GAGGTCGACG	GTATCGATAA	GCTTGATATC 80
81	GAATTCGGCA	CGAGTTTAC	ACCATCACCA	AAACCCAGAA	TTTATGATG	AGATTAAAA	AGAGTTGCC	ACTCAGCTGC 160
161	ATGAAAAGCA	CCACCTGTTG	CTCACATCT	TCCATGTGAC	CTGTGACAAC	TCAAGTAAAG	GAAGCAGGAA	GAAGAGGGAT 240
241	GTCGTTGAAA	CCCAAGTTGG	CTACTCCTGG	CTTCCCTCC	TGAAAGACGG	AAGGGTGGTG	ACAAGCGAGC	AGCACATCCC 320
321	GGTCTCGGCG	AACCTTCCTT	CGGGCTATCT	TGGCTACCAA	GAGCTTGGGA	TGGGCAGGCA	TTATGGTCCG	GAAATTAAT 400
401	GGGTAGATGG	AGGCAAGCCA	CTGCTGAAAA	TTTCCACTCA	TCTGGTTTCT	ACAGGGATAC	TCAGGATCAG	CATTTACATA 480
481	ATTTTTTCCA	GTACTGTCAG	AAAACCGAAT	CTGGAGCCCA	AGCCTTAGGA	AACGAACTTG	TAAAGTACCT	TAAGAGTCTG 560
561	CATGCCATGG	AAGGCCACGT	GATGATCGCC	TTCTTGCCCA	CTATCCTAAA	CCAGCTGTTT	CGAGTCTCA	CCAGAGCCAC 640
641	ACAGGAAGAA	GTCGCGTTA	ACGTGACTCG	GGTCATTATT	CATGTGGTTG	CCCAGTGCCA	TGAGGAAGGA	TTGGAGAGCC 720
721	ACTTGAGGTC	ATATGTTAAG	TACGCGTATA	AGGCTGAGCC	ATATGTTGCC	TCTGAATACA	AGACAGTGCA	TGAAGAACTG 800
801	ACCAAAATCCA	TGACCACGAT	TCTCAAGCCT	TCTGCCGATT	TCCTCACCAG	CAACAACTA	CTGAGGTACT	CATGGTTTTT 880
881	CTTTGATGTA	CTGATCAAAT	CTATGGCTCA	GCATTTGATA	GAGAAGTCCA	AAGTTAAGTT	GCTGCGAAAC	CAGAGATTTC 960
961	CTGCATCCTA	TCATCATGCA	GCGGAAACCG	TTGTAAATAT	GCTGATGCCA	CACATCACTC	AGAAGTTTGG	AGATAATCCA 1040
1041	GAGGCATCTA	AGAAGCGGAA	TCATAGCCTT	GCTGTCTTCA	TCAAGAGATG	TTTACCTTTC	ATGGACAGGT	GCTTTGTCTT 1120
1121	CAAGCAGATC	AACAACCTACA	TTAGCTGTTT	TGCTCCTGGA	GACCAAAAG	CCCTCTTTGA	ATACAGTTT	GAAATTTCTC 1200
1201	GTGTAGTGTG	CAACCATGAA	CATTATATTC	CGTTGAACCT	ACCAATGCCA	TTTGGAAAAG	GCAGGATTCA	AAGATACCAA 1280
1281	GACCTCCAGC	TTGACTACTC	ATTAAACAGAT	GAGTTCTGCA	GAAACCACTT	CTTGGTGGGA	CTGTTACTGA	GGGAGGTGGG 1360
1361	GACAGCCCTC	CAGGAGTTCC	GGGAGGTCCG	TCTGATCGCC	ATCAGTGTGC	TCAAGAACCT	GCTGATAAAG	CATTCTTTTG 1440
1441	ATGACAGATA	TGCTTCAAGG	AGCCATCAGG	CAAGGATAGC	CACCTCTTAC	CTGCCTCTGT	TGCTGTCTGT	GATTGAAAAC 1520
1521	GTCCAGCGGA	TCATGTGTAG	GGATGTGTCA	CCCTTCCCTG	TGAACGCGGG	CATGACCGTG	AAGGATGAAT	CCCTGGCTCT 1600
1601	ACCAGCTGTG	AATCCGCTGG	TGACGCCGCA	GAAGGGAAGC	ACCTCTGGCA	ACAGCCTGCA	CAAGGACCTG	CTGGGCGCCA 1680
1681	TCTCCGGCAT	TGCTTCTCCA	TATACAACTT	CAACTCCAAA	CATCAACAGT	GTGAGAAATG	CTGATTCTG	AGGATCTCTC 1760
1761	ATAAGCAGAG	ATTCGGGTAA	CAGCCTTCCA	GAAGGGAATA	GTGAGAAGAG	CAATTCCCTG	GATAAGCACC	AACAAAGTAG 1840
1841	CACATTGGGA	AATTCGGTGG	TTGCTGTGTA	TAAACTTGAC	CAGTCTGAGA	TTAAGAGCCT	ACTGATGTGT	TTCTCTTACA 1920
1921	TCTTAAAGAG	CATGCTGTAT	GATGCTTTGT	TTACATAATG	GAACAAGGCT	TCAACATCTG	AACCTATGTA	TTTTTTTACA 2000
2001	ATATCTGAAG	CTGCGCTGCA	CCAGTTCCAG	TACATGGGGA	AGCGATACAT	AGCCAGGAAC	CAGGAGGGGT	TGGGACCCAT 2080
2081	AGTTCATGAT	CGAAAGTCTC	AGACATTGCC	TGTTTCCCGT	AACAGAACAG	GAATGATGCA	TGCCAGATTG	CAGCAGCTGG 2160
2161	GCAGCCTGGA	TAACTCTCTC	ACTTTTAAAC	ACAGCTATGG	CCACTCGGAC	GCAGATGTTT	TGCACCACTG	ATTAATTGAA 2240
2241	GCCAAACATTG	TACTGAGGT	TTGCCGTGCA	GCTCTGGACA	GCCTTTCTCT	ATTACATATG	GCCTTTAAGA	ACCAGCTCCT 2320
2321	GGCCGACCAT	GGACATAATC	CTCTCATGAA	AAAAGTTTTT	GATGTCTACC	TGTGTTTTCT	TCAAAAACAT	CAGTCTGAAA 2400
2401	CGGCTTTTAA	AAATGTCTTC	ACTGCCCTAA	GGTCTTAAAT	TTATAGGTTT	CCCTCAACAT	TCTATGAAG	GAGAGCGGAC 2480
2481	ATGTGTGCGG	CTCTGTGTTA	CGAGATTCTC	AAGTGTGTA	ACTCAAGCT	GAGCTCCATC	AGGACGGAGG	CCTCCAGGCT 2560
2561	GCTTCTATTC	CTGATGAGGA	ACAACCTTGA	TTACACTGGA	AAGAAGTCTT	TTGTCCGGAC	ACATTTGCAA	GTCTCATAT 2640
2641	CTGTGAGCCA	GCTGATAGCA	GACGTTGTTG	GCATTGGGGA	AACCAGATTG	CAGCAGTCCC	TGTCCATCAT	CAACAACCTG 2720
2721	GCCAAACAGTG	ACCGGCTTAT	TAAAGCACAC	AGCTTCTCCT	CTGATGTGAA	GGACTTAACC	AAAAGGATAC	GCACGGTGCT 2800
2801	AATGGCCACC	GCCCAGATGA	AGGAGCATGA	GAACGACCCA	GAGATGCTGG	TGGACCTCCA	GTACAGCCTG	GCCAAATCCT 2880
2881	ATGCCAGCAC	GCCCAGGCTC	AGGAAGACGT	GGCTCGACAG	CATGGCCAGG	ATCCATGTCA	AAAATGGCGA	TCTCTCAGAG 2960
2961	GCAGCAATGT	GCTATGTCCA	CGTAACAGCC	CTAGTGGCAG	AATATCTCAC	ACGGAAGGCG	GTGTTTAGAC	AAGGATGCAC 3040
3041	CGCCTTCAAG	GTCATTACCC	CAACATGACA	CGAGGAGGCC	TCCATGATGG	AAGACGTGGG	GATGCAGGAT	GTCCATTTCA 3120
3121	ACGAGGATGT	GCTGATGGAG	CTCCTTGAGC	AGTGCGCAGA	TGGACTCTGG	AAAGCCGAGC	GCTACGAGCT	CATCGCCGAC 3200
3201	ATCTACAAAC	TTATCATCCC	CATTTATGAG	AAGCGGAGGG	ATTTCTTTGA	AGATGAAGAT	GGAAAGGAGT	ATATTTACAA 3280
3281	GGAACCCAAA	CTCACACCGC	TGTCGGAAAT	TTCTCAGAGA	CTCCTTAAAC	TGTACTCGGA	TAAATTTGTT	TCTGAAAATG 3360
3361	TCAAAATGAT	ACAGGATTCT	GGCAAGGTCA	ACCCTAAGGA	TCTGGATTCT	AAGTATGCAT	ACATCCAGGT	GACTCACGTC 3440
3441	ATCCCTTCTT	TTGACGAAAA	AGAGTTGCAA	GAAAGGAAAA	CAGAGTTTGA	GAGATCCCAC	AACATCCGCC	GCTTCATGTT 3520
3521	TGAGATGCCA	TTTACGCAGA	CCGGGAAGAG	GCAGGGCGGG	GTGGAAGAGC	AGTGCAAACG	GCGCACCATC	CTGACAGCCA 3600
3601	TACACTGCTT	CCCTTATGTG	AAGAAGCGCA	TCCCTGTCTAT	GTACCAGCAC	CACACTGACC	TGAACCCCAT	CGAGGTGGCC 3680
3681	ATTGACGAGA	TGAGTAAGAA	GGTGGCGGAG	CTCGGCGAGC	TGTGCTCCTC	GGCCGAGGTG	GACATGATCA	AACTGCAGCT 3760
3761	CAAACTCCAG	GGCAGCGTGA	GTGTTCAGGT	CAATGCTGGC	CCACTAGCAT	ATGCGCGAGC	TTTCTTAGAT	GATACAAACA 3840
3841	CAAAGCGATA	TCCTGACAAAT	AAAGTGAAGC	TGCTTAAAGGA	AGTTTTCAGG	CAATTTGTGG	AAGCTTGGCG	TCAAGCCTTA 3920
3921	CGCGTAAACG	AACGCTGTGAT	TAAAGAAGAC	CAGCTCGAGT	ATCAGGAAGA	AATGAAAGCC	AACCTACAGG	AAATGGCGAA 4000
4001	GGAGCTTTCT	GAAATCATGC	ATGAGCAGAT	CTGCCCTCTG	GAGGAGAAGA	CGAGCGTCTT	ACCGAATTCC	CTTCACATCT 4080
4081	TCAACGCCAT	CAGTGGGACT	CCAACAAGCA	CAATGTTTCA	CGGGATGACC	AGCTCGTCTT	CGGTCTGTGT	ATTACATCTC 4160
4161	ATGGCCCGTG	TGTGGGGACT	TGCTTTGTCA	TTTGCAAACT	CAGGATGCTT	TCCAAAGCCA	ATCACTGGGG	AGACCGAGCA 4240
4241	CAGGGAGGAC	CAAGGGGAAG	GGGAGAGAAA	GGAAATAAAG	AACAACGTTA	TTTCTTAAAC	GACTTTCTAT	AGGAGTTGTA 4320
4321	AGAAGGTGCA	CATATTTTTT	TAAATCTCAC	TGGCAATATT	CAAAAGTTTC	ATTGTGCTTT	AACAAAGGTG	TGGTAGACAC 4400
4401	TCTTTGAGCTG	GACTTAGATT	TTATTTCTCC	TTGCAGAGTA	GTGTTAGAAAT	AGATGGCCTA	CAGAAAAAAA	AGGTTCTGGG 4480
4481	ATCTACATGG	CAGGGAGGGC	TGCACTGACA	TTGATGCCTG	GGGGACCTTT	TGCTCGACT	CGTGCCGGAA	ATCTGATCGT 4560
4561	AATCAGGGTA	CAGAACTTAC	TAGTTTGTCT	TAGGAGTATG	TTGTATGACT	AGGATTGTGT	CTATTATCTC	ATTCAACAAC 4640
4641	ATAGAGCAAG	AATAGTGAAG	TAACTGAGCT	AGACACTCAA	TTAATCCGCT	ACTGGCTTCA	AGTCAGAACT	TTGTCAATTA 4720
4721	TCATCGACTC	CGGGACGGTC	ATATATGTAT	TACATTTCTA	CATTTTAAAT	ACTCACATGG	GCTTATGCAT	TAAGTTTAAAT 4800
4801	TGTGATAAAT	TTGTGCTGGT	CCAGTATATG	CAATACACTT	TAATGGTTTA	TTCTGTGCTAT	AAAAATGTGC	AATATGGAGA 4880
4881	TGTATACAA	TCTTTACT						4898

FIG.10H

	10	20	30	40	50	60	70	80
1	MEGHVMIAFL	PTILNQLFRV	LTRATQEEVA	VNVTRVIIHV	VAQCHEEGLE	SHLRSYVKYA	YKAEPYVASE	YKTVHEELTK 80
81	SMTTILKPSA	DFLTSNKLLR	YSWFFFDVLI	KSMAQH LIEN	SKVKLLRNQR	FPASYHHAAE	TVVNMMPHI	TQKFGDNPEA 160
161	SKNANHSLAV	FIKRCFTFMD	RGFVFKQINN	YISCFAPGDP	KTLFEYKFEF	LRVVCNHEHY	IPLNLPMPFG	KGRIQRYQDL 240
241	QLDYSLTDEF	CRNHFLVGLL	LREVG TALQE	FREVRLIAIS	VLKNLLIKHS	FDDRYASRSH	QARIATLYLP	LFGLLIENVQ 320
321	RINVRDVSPF	PVNAGMTVKD	ESLALPAVNP	LVTPQKGSTL	DNSLHKDLLG	AISGIASPYT	TSTPNINSVR	NADSRGSLIS 400
401	TDSGNSLPER	NSEKSNSLDK	HQQSSTLGNS	VVRCDKLDQS	EIKSLIMCFL	YILKSMDDA	LFTYWNKAST	SEIMDFFTIS 480
481	EVCLHQFQYM	GKRYIARNQE	GLGPIVHDRK	SQTLEPVSRR	TGMMHARLQQ	LGSLDNSLTF	NHSYGHSDAD	VLHQSLLEAN 560
561	IATEVCLTAL	DTLSLFTLAF	KNQLLADHGH	NPLMKKVF DV	YLCFLQKHQS	ETALKNVFTA	LRS LIYKFPS	TFYEGRADMC 640
641	AALCYEILKC	CNSKLSSIRT	EASQLLYFLM	RNNFDYTGKK	SFVRTHLQVI	ISVSQLIADV	VGIGETRFQQ	SLSIINN CAN 720
721	SDRLIKHTSF	SSDVKD LTKR	IRTVLMATAQ	MKEHENDPEM	LVDLQYSLAK	SYASTPELRK	TWLD SMARIH	VKNGDLSEAA 800
801	MCYVHV TALV	AEYLTRKGVF	RQGCTAFRVI	TPNIDEEASM	MEDVGMQDVH	FNEDVLMELL	EQCADGLWKA	ERYELIADIY 880
881	KLIIPYIEKR	RDFEDEDGK	EYIYKEPKLT	PLSEISQRLL	KLYSDKFGSE	NVKMIQDSGK	VNPKDLD SKY	AYIQVTHVIP 960
961	FFDEKELQER	KTEFERSHNI	RRFMFEMPFT	QTGKRQGGVE	EQCKRR TILT	AIHCFPYVKK	RIPVMYQHHT	DLNPIEVAID 1040
1041	EMSKKVAELR	QLC SSAEVD M	IKLQKLQGS	VSVQVNAGPL	AYARAFLDDT	NTKRYPDNKV	KLLKEVFRQF	VEACQALAV 1120
1121	NERLIKEDQL	EYQEEMKANY	REMAKELSEI	MHEQICPLEE	KTSVLENSLH	IFNAISGTPT	STMVHGMTSS	SSVV 1195

FIG. 10H (cont.)

Exon 1A (-182 to -102)

GCAGGGGAAAAACCTGGCCCCATGATTCACTTACTTCCCACCGGATCTCTCCCATGACACGTGAGGATTA
TTACAATTTAA -102

Exon 1B (-219 to -102)

TTATCCCTTTTACTACTTGGCAAGTGAGTTCGGTAGATGGGAGTGGAGAAGAGAACCTTAGAATCATTGTTTGTAGTCTTCAT
CTTTCACAGCTCAGGCTGAAGGCCTTTCCTTGCTGAGA -102

Exon 1C (-143 to -102)

GCGGCAGAGCGTGTCTGAGGTGGTGCGCGGCTCCGTGCTCCT -102

Exon2 and the rest of human CLASP2 cDNA

-101 -79
GGCAAAGCCAAAGCTAATTGAGC

-78

-1

AAGCTAATTGAGGCACTGCACTATGAAAATGTCATCGTCCAGAAGAAGACTCAGATCCTGAACGACTGTTTACGGGAG

1/1

31/11

ATG CTG CTC TTC CCT TAC GAT GAC TTT CAG ACG GCC ATC CTG AGA CGA CAG GGT CGA TAC
Met leu leu phe pro tyr asp asp phe gln thr ala ile leu arg arg gln gly arg tyr

61/21

91/31

ATA TGC TCA ACA GTG CCT GCG AAG GCG GAA GAG GAA GCA CAG AGC TTG TTT GTT ACA GAG
ile cys ser thr val pro ala lys ala glu glu glu ala gln ser leu phe val thr glu

121/41

151/51

TGC ATC AAA ACC TAT AAC TCT GAC TGG CAT CTT GTG AAC TAT AAA TAT GAA GAT TAC TCA
cys ile lys thr tyr asn ser asp trp his leu val asn tyr lys tyr glu asp tyr ser

181/61

211/71

GGA GAG TTT CGA CAG CTT CCG AAC AAA GTG GTC AAG TTG GAT AAA CTT CCA GTT CAT GTC
gly glu phe arg gln leu pro asn lys val val lys leu asp lys leu pro val his val

241/81

271/91

TAT GAA GTT GAC GAG GAG GTC GAC AAA GAT GAG GAT GCT GCC TCC CTT GGT TCC CAG AAG
tyr glu val asp glu glu val asp lys asp glu asp ala ala ser leu gly ser gln lys

301/101

331/111

GGT GGG ATC ACC AAG CAT GGC TGG CTG TAC AAA GGC AAC ATG AAC AGT GCC ATC AGC GTG
gly gly ile thr lys his gly trp leu tyr lys gly asn met asn ser ala ile ser val

361/121

391/131

ACC ATG AGG TCA TTT AAG AGA CGA TTT TTC CAC CTG ATT CAA CTT GGC GAT GGA TCC TAT
thr met arg ser phe lys arg arg phe phe his leu ile gln leu gly asp gly ser tyr

421/141

451/151

AAT TTG AAT TTT TAT AAA GAT GAA AAG ATC TCC AAA GAA CCA AAA GGA TCA ATA TTT CTG
asn leu asn phe tyr lys asp glu lys ile ser lys glu pro lys gly ser ile phe leu

481/161

511/171

GAT TCC TGT ATG GGT GTC GTT CAG AAC AAC AAA GTC AGG CGT TTT GCT TTT GAG CTC AAG
asp ser cys met gly val val gln asn asn lys val arg arg phe ala phe glu leu lys

541/181 571/191
ATG CAG GAC AAA AGT AGT TAT CTC TTG GCA GCA GAC AGT GAA GTG GAA ATG GAA GAA TGG
met gln asp lys ser ser tyr leu leu ala ala asp ser glu val glu met glu glu trp

601/201 631/211
ATC ACA ATT CTA AAT AAG ATC CTC CAG CTC AAC TTT GAA GCT GCA ATG CAA GAA AAG CGA
ile thr ile leu asn lys ile leu gln leu asn phe glu ala ala met gln glu lys arg

661/221 691/231
AAT GGC GAC TCT CAC GAA GAT GAT GAA CAA AGC AAA TTG GAA GGT TCT GGT TCC GGT TTA
asn gly asp ser his glu asp asp glu gln ser lys leu glu gly ser gly ser gly leu

721/241 751/251
GAT AGC TAC CTG CCG GAA CTT GCC AAG AGT GCA AGA GAA GCA GAA ATC AAA CTA AAA AGT
asp ser tyr leu pro glu leu ala lys ser ala arg glu ala glu ile lys leu lys ser

781/261 811/271
GAA AGC AGA GTC AAA CTT TTT TAT TTG GAC CCA GAT GCC CAG AAG CTT GAC TTC TCA TCA
glu ser arg val lys leu phe tyr leu asp pro asp ala gln lys leu asp phe ser ser

841/281 871/291
GCT GAG CCA GAA GTG AAG TCA TTT GAA GAG AAG TTT GGA AAA AGG ATC CTT GTC AAG TGC
ala glu pro glu val lys ser phe glu glu lys phe gly lys arg ile leu val lys cys

901/301 931/311
AAT GAT TTA TCT TTC AAT TTG CAA TGC TGT GTT GCC GAA AAT GAA GAA GGA CCC ACT ACA
asn asp leu ser phe asn leu gln cys cys val ala glu asn glu glu gly pro thr thr

961/321 991/331
AAT GTT GAA CCT TTC TTT GTT ACT CTA TCC CTG TTT GAC ATA AAA TAC AAC CGG AAG ATT
asn val glu pro phe phe val thr leu ser leu phe asp ile lys tyr asn arg lys ile

1021/341 1051/351
TCT GCC GAT TTC CAC GTA GAC CTG AAC CAT TTC TCA GTG AGG CAA ATG CTC GCC ACC ACG
ser ala asp phe his val asp leu asn his phe ser val arg gln met leu ala thr thr

1081/361 1111/371
TCC CCG GCG CTG ATG AAT GGC AGT GGG CAG AGC CCA TCT GTC CTC AAG GGC ATC CTT CAT
ser pro ala leu met asn gly ser gly gln ser pro ser val leu lys gly ile leu his

1141/381 1171/391
GAA GCC GCC ATG CAG TAT CCG AAG CAG GGA ATA TTT TCA GTC ACT TGT CCT CAT CCA GAT
glu ala ala met gln tyr pro lys gln gly ile phe ser val thr cys pro his pro asp

1201/401 1231/411
ATA TTT CTT GTG GCC AGA ATT GAA AAA GTC CTT CAG GGG AGC ATC ACA CAT TGC GCT GAG
ile phe leu val ala arg ile glu lys val leu gln gly ser ile thr his cys ala glu

1261/421 1291/431
CCA TAT ATG AAA AGT TCA GAC TCT TCT AAG GTG GCC CAG AAG GTG CTG AAG AAT GCC AAG
pro tyr met lys ser ser asp ser ser lys val ala gln lys val leu lys asn ala lys

1321/441 1351/451
CAG GCA TGC CAA AGA CTA GGA CAG TAT AGA ATG CCA TTT GCT TGG GCA GCA AGG ACA TTG
gln ala cys gln arg leu gly gln tyr arg met pro phe ala trp ala ala arg thr leu

1381/461 1411/471
TTT AAG GAT GCA TCT GGA AAT CTT GAC AAA AAT GCC AGA TTT TCT GCC ATC TAC AGG CAA
phe lys asp ala ser gly asn leu asp lys asn ala arg phe ser ala ile tyr arg gln

1441/481 1471/491
GAC AGC AAT AAG CTA TCC AAT GAT GAC ATG CTC AAG TTA CTT GCA GAC TTT CGG AAA CCT
asp ser asn lys leu ser asn asp asp met leu lys leu leu ala asp phe arg lys pro

1501/501 1531/511
GAG AAG ATG GCT AAG CTC CCA GTG ATT TTA GGC AAT CTA GAC ATT ACA ATT GAT AAT GTT
glu lys met ala lys leu pro val ile leu gly asn leu asp ile thr ile asp asn val

1561/521 1591/531
TCC TCA GAC TTC CCT AAT TAT GTT AAT TCA TCA TAC ATT CCC ACA AAA CAA TTT GAA ACC
ser ser asp phe pro asn tyr val asn ser ser tyr ile pro thr lys gln phe glu thr

1621/541 1651/551
TGC AGT AAA ACT CCC ATC ACG TTT GAA GTG GAG GAA TTT GTG CCC TGC ATA CCA AAA CAC
cys ser lys thr pro ile thr phe glu val glu glu phe val pro cys ile pro lys his

1681/561 1711/571
ACT CAG CCT TAC ACC ATC TAC ACC AAT CAC CTT TAC GTT TAT CCT AAG TAC TTG AAA TAC
thr gln pro tyr thr ile tyr thr asn his leu tyr val tyr pro lys tyr leu lys tyr

1741/581 1771/591
GAC AGT CAG AAG TCT TTT GCC AAG GCT AGA AAT ATT GCG ATT TGC ATT GAA TTC AAA GAT
asp ser gln lys ser phe ala lys ala arg asn ile ala ile cys ile glu phe lys asp

1801/601 1831/611
TCA GAT GAG GAA GAC TCT CAG CCC CTT AAG TGC ATT TAT GGC AGA CCT GGT GGG CCA GTT
ser asp glu glu asp ser gln pro leu lys cys ile tyr gly arg pro gly gly pro val

1861/621 1891/631
TTC ACA AGA AGC GCC TTT GCT GCA GTT TTA CAC CAT CAC CAA AAC CCA GAA TTT TAT GAT
phe thr arg ser ala phe ala ala val leu his his his gln asn pro glu phe tyr asp

1921/641 1951/651
GAG ATT AAA ATA GAG TTG CCC ACT CAG CTG CAT GAA AAG CAC CAC CTG TTG CTC ACA TTC
glu ile lys ile glu leu pro thr gln leu his glu lys his his leu leu leu thr phe

1981/661 2011/671
TTC CAT GTC AGC TGT GAC AAC TCA AGT AAA GGA AGC ACG AAG AAG AGG GAT GTC GTT GAA
phe his val ser cys asp asn ser ser lys gly ser thr lys lys arg asp val val glu

2041/681 2071/691
ACC CAA GTT GGC TAC TCC TGG CTT CCC CTC CTG AAA GAC GGA AGG GTG GTG ACA AGC GAG
thr gln val gly tyr ser trp leu pro leu leu lys asp gly arg val val thr ser glu

2101/701 2131/711
CAG CAC ATC CCG GTC TCG GCG TAC CTT CCT TCG GGC CAT CTT GGC TAC CAA GAG CTT GGG
gln his ile pro val ser ala tyr leu pro ser gly his leu gly tyr gln glu leu gly

2161/721 2191/731
ATG GGC AGG CAT TAT GGT CCG GAA ATT AAA TGG GTA GAT GGA GGC AAG CCA CTG CTG AAA
met gly arg his tyr gly pro glu ile lys trp val asp gly gly lys pro leu leu lys

2221/741 2251/751
ATT TCC ACT CAT CTG GTT TCT ACA GTG TAT ACT CAG GAT CAG CAT TTA CAT AAT TTT TTC
ile ser thr his leu val ser thr val tyr thr gln asp gln his leu his asn phe phe

2281/761 2311/771
CAG TAC TGT CAG AAA ACC GAA TCT GGA GCC CAA GCC TTA GGA AAC GAA CTT GTA AAG TAC
gln tyr cys gln lys thr glu ser gly ala gln ala leu gly asn glu leu val lys tyr

2341/781 2371/791
CTT AAG AGT CTG CAT GCG ATG GAA GGC CAC GTG ATG ATC GCC TTC TTG CCC ACT ATC CTA
leu lys ser leu his ala met glu gly his val met ile ala phe leu pro thr ile leu

2401/801 2431/811
AAC CAG CTG TTC CGA GTC CTC ACC AGA GCC ACA CAG GAA GAA GTC GCG GTT AAC GTG ACT
asn gln leu phe arg val leu thr arg ala thr gln glu glu val ala val asn val thr

2461/821 2491/831
CGG GTC ATT ATT CAT GTG GTT GCC CAG TGC CAT GAG GAA GGA TTG GAG AGC CAC TTG AGG
arg val ile ile his val val ala gln cys his glu glu gly leu glu ser his leu arg

2521/841 2551/851
TCA TAT GTT AAG TAC GCG TAT AAG GCT GAG CCA TAT GTT GCC TCT GAA TAC AAG ACA GTG
ser tyr val lys tyr ala tyr lys ala glu pro tyr val ala ser glu tyr lys thr val

2581/861 2611/871
CAT GAA GAA CTG ACC AAA TCC ATG ACC ACG ATT CTC AAG CCT TCT GCC GAT TTC CTC ACC
his glu glu leu thr lys ser met thr thr ile leu lys pro ser ala asp phe leu thr

2641/881 2671/891
AGC AAC AAA CTA CTG AGG TAC TCA TGG TTT TTC TTT GAT GTA CTG ATC AAA TCT ATG GCT
ser asn lys leu leu arg tyr ser trp phe phe phe asp val leu ile lys ser met ala

2701/901 2731/911
CAG CAT TTG ATA CAG AAC TCC AAA GTT AAG TTG CTG CGA AAC CAG AGA TTT CCT GCA TCC
gln his leu ile glu asn ser lys val lys leu leu arg asn gln arg phe pro ala ser

2761/921 2791/931
TAT CAT CAT GCA GCG GAA ACC GTT GTA AAT ATG CTG ATG CCA CAC ATC ACT CAG AAG TTT
tyr his his ala ala glu thr val val asn met leu met pro his ile thr gln lys phe

2821/941 2851/951
GGA GAT AAT CCA GAG GCA TCT AAG AAC GCG AAT CAT AGC CTT GCT GTC TTC ATC AAG AGA
gly asp asn pro glu ala ser lys asn ala asn his ser leu ala val phe ile lys arg

2881/961 2911/971
TGT TTC ACC TTC ATG GAC AGG GGC TTT GTC TTC AAG CAG ATC AAC AAC TAC ATT AGC TGT
cys phe thr phe met asp arg gly phe val phe lys gln ile asn asn tyr ile ser cys

2941/981 2971/991
TTT GCT CCT GGA GAC CCA AAG ACC CTC TTT GAA TAC AAG TTT GAA TTT CTC CGT GTA GTG
phe ala pro gly asp pro lys thr leu phe glu tyr lys phe glu phe leu arg val val

3001/1001 3031/1011
TGC AAC CAT GAA CAT TAT ATT CCG TTG AAC TTA CCA ATG CCA TTT GGA AAA GGC AGG ATT
cys asn his glu his tyr ile pro leu asn leu pro met pro phe gly lys gly arg ile

3061/1021 3091/1031
CAA AGA TAC CAA GAC CTC CAG CTT GAC TAC TCA TTA ACA GAT GAG TTC TGC AGA AAC CAC
gln arg tyr gln asp leu gln leu asp tyr ser leu thr asp glu phe cys arg asn his

3121/1041 3151/1051
TTC TTG GTG GGA CTG TTA CTG AGG GAG GTG GGG ACA GCC CTC CAG GAG TTC CGG GAG GTC
phe leu val gly leu leu leu arg glu val gly thr ala leu gln glu phe arg glu val

3181/1061 3211/1071
CGT CTG ATC GCC ATC AGT GTG CTC AAG AAC CTG CTG ATA AAG CAT TCT TTT GAT GAC AGA
arg leu ile ala ile ser val leu lys asn leu leu ile lys his ser phe asp asp arg

3241/1081 3271/1091
TAT GCT TCA AGG AGC CAT CAG GCA AGG ATA GCC ACC CTC TAC CTG CCT CTG TTT GGT CTG
tyr ala ser arg ser his gln ala arg ile ala thr leu tyr leu pro leu phe gly leu

3301/1101 3331/1111
CTG ATT GAA AAC GTC CAG CGG ATC AAT GTG AGG GAT GTG TCA CCC TTC CCT GTG AAC GCG
leu ile glu asn val gln arg ile asn val arg asp val ser pro phe pro val asn ala

3361/1121 3391/1131
GGC ATG ACC GTG AAG GAT GAA TCC CTG GCT CTA CCA GCT GTG AAT CCG CTG GTG ACG CCG
gly met thr val lys asp glu ser leu ala leu pro ala val asn pro leu val thr pro

3421/1141 3451/1151
CAG AAG GGA AGC ACC CTG GAC AAC AGC CTG CAC AAG GAC CTG CTG GGC GCC ATC TCC GGC
gln lys gly ser thr leu asp asn ser leu his lys asp leu leu gly ala ile ser gly

3481/1161 3511/1171
ATT GCT TCT CCA TAT ACA ACC TCA ACT CCA AAC ATC AAC AGT GTG AGA AAT GCT GAT TCG
ile ala ser pro tyr thr thr ser thr pro asn ile asn ser val arg asn ala asp ser

3541/1181 3571/1191
AGA GGA TCT CTC ATA AGC ACA GAT TCG GGT AAC AGC CTT CCA GAA AGG AAT AGT GAG AAG
arg gly ser leu ile ser thr asp ser gly asn ser leu pro glu arg asn ser glu lys

3601/1201 3631/1211
AGC AAT TCC CTG GAT AAG CAC CAA CAA AGT AGC ACA TTG GGA AAT TCC GTG GTT CGC TGT
ser asn ser leu asp lys his gln gln ser ser thr leu gly asn ser val val arg cys

3661/1221 3691/1231
GAT AAA CTT GAC CAG TCT GAG ATT AAG AGC CTA CTG ATG TGT TTC CTC TAC ATC TTA AAG
asp lys leu asp gln ser glu ile lys ser leu leu met cys phe leu tyr ile leu lys

3721/1241 3751/1251
AGC ATG TCT GAT GAT GCT TTG TTT ACA TAT TGG AAC AAG GCT TCA ACA TCT GAA CTT ATG
ser met ser asp asp ala leu phe thr tyr trp asn lys ala ser thr ser glu leu met

3781/1261 3811/1271
GAT TTT TTT ACA ATA TCT GAA GTC TGC CTG CAC CAG TTC CAG TAC ATG GGG AAG CGA TAC
asp phe phe thr ile ser glu val cys leu his gln phe gln tyr met gly lys arg tyr

3841/1281 3871/1291
ATA GCC AGG AAC CAG GAG GGG TTG GGA CCC ATA GTT CAT GAT CGA AAG TCT CAG ACA TTG
ile ala arg asn gln glu gly leu gly pro ile val his asp arg lys ser gln thr leu

4741/1581
GAT CTC TCA GAG GCA GCA ATG TGC TAT GTC CAC GTA ACA GCC CTA GTG GCA GAA-TAT CTC
asp leu ser glu ala ala met cys tyr val his val thr ala leu val ala glu tyr leu

4771/1591
4801/1601
ACA CGG AAA GGC GTG TTT AGA CAA GGA TGC ACC GCC TTC AGG GTC ATT ACC CCA AAC ATC
thr arg lys gly val phe arg gln gly cys thr ala phe arg val ile thr pro asn ile

4831/1611
4861/1621
GAC GAG GAG GCC TCC ATG ATG GAA GAC GTG GGG ATG CAG GAT GTC CAT TTC AAC GAG GAT
asp glu glu ala ser met met glu asp val gly met gln asp val his phe asn glu asp

4891/1631
4921/1641
GTG CTG ATG GAG CTC CTT GAG CAG TGC GCA GAT GGA CTC TGG AAA GCC GAG CGC TAC GAG
val leu met glu leu leu glu gln cys ala asp gly leu trp lys ala glu arg tyr glu

4951/1651
4981/1661
CTC ATC GCC GAC ATC TAC AAA CTT ATC ATC CCC ATT TAT GAG AAG CGG AGG GAT TTC TTT
leu ile ala asp ile tyr lys leu ile ile pro ile tyr glu lys arg arg asp phe phe

5011/1671
5041/1681
GAA GAT GAA GAT GGA AAG GAG TAT ATT TAC AAG GAA CCC AAA CTC ACA CCG CTG TCG GAA
glu asp glu asp gly lys glu tyr ile tyr lys glu pro lys leu thr pro leu ser glu

5071/1691
5101/1701
ATT TCT CAG AGA CTC CTT AAA CTG TAC TCG GAT AAA TTT GGT TCT GAA AAT GTC AAA ATG
ile ser gln arg leu leu lys leu tyr ser asp lys phe gly ser glu asn val lys met

5131/1711
5161/1721
ATA CAG GAT TCT GGC AAG GTC AAC CCT AAG GAT CTG GAT TCT AAG TAT GCA TAC ATC CAG
ile gln asp ser gly lys val asn pro lys asp leu asp ser lys tyr ala tyr ile gln

5191/1731
5221/1741
GTG ACT CAC GTC ATC CCC TTC TTT GAC GAA AAA GAG TTG CAA GAA AGG AAA ACA GAG TTT
val thr his val ile pro phe phe asp glu lys glu leu gln glu arg lys thr glu phe

5251/1751
5281/1761
GAG AGA TCC CAC AAC ATC CGC CGC TTC ATG TTT GAG ATG CCA TTT ACG CAG ACC GGG AAG
glu arg ser his asn ile arg arg phe met phe glu met pro phe thr gln thr gly lys

5311/1771
5341/1781
AGG CAG GGC GGG GTG GAA GAG CAG TGC AAA CGG CGC ACC ATC CTG ACA GCC ATA CAC TGC
arg gln gly gly val glu glu gln cys lys arg arg thr ile leu thr ala ile his cys

5371/1791
5401/1801
TTC CCT TAT GTG AAG AAG CGC ATC CCT GTC ATG TAC CAG CAC CAC ACT GAC CTG AAC CCC
phe pro tyr val lys lys arg ile pro val met tyr gln his his thr asp leu asn pro

5431/1811
5461/1821
ATC GAG GTG GCC ATT GAC GAG ATG AGT AAG AAG GTG GCG GAG CTC CGG CAG CTG TGC TCC
ile glu val ala ile asp glu met ser lys lys val ala glu leu arg gln leu cys ser

5491/1831
5521/1841
TCG GCC GAG GTG GAC ATG ATC AAA CTG CAG CTC AAA CTC CAG GGC AGC GTG AGT GTT CAG
ser ala glu val asp met ile lys leu gln leu lys leu gln gly ser val ser val gln

5551/1851

5581/1861	5611/1871
GTC AAT GCT GGC CCA CTA GCA TAT GCG CGA GCT TTC TTA GAT GAT ACA AAC ACA AAG CGA	
val asn ala gly pro leu ala tyr ala arg ala phe leu asp asp thr asn thr lys arg	
5641/1881	5671/1891
TAT CCT GAC AAT AAA GTG AAG CTG CTT AAG GAA GTT TTC AGG CAA TTT GTG GAA GCT TGC	
tyr pro asp asn lys val lys leu leu lys glu val phe arg gln phe val glu ala cys	
5701/1901	5731/1911
GGT CAA GCC TTA GCG GTA AAC GAA CGT CTG ATT AAA GAA GAC CAG CTC GAG TAT CAG GAA	
gly gln ala leu ala val asn glu arg leu ile lys glu asp gln leu glu tyr gln glu	
5761/1921	5791/1931
GAA ATG AAA GCC AAC TAC AGG GAA ATG GCG AAG GAG CTT TCT GAA ATC ATG CAT GAG CAG	
glu met lys ala asn tyr arg glu met ala lys glu leu ser glu ile met his glu gln	
5821/1941	5851/1951
ATC TGC CCC CTG GAG GAG AAG ACG AGC GTC TTA CCG AAT TCC CTT CAC ATC TTC AAC GCC	
ile cys pro leu glu glu lys thr ser val leu pro asn ser leu his ile phe asn ala	
5881/1961	5911/1971
ATC AGT GGG ACT CCA ACA AGC ACA ATG GTT CAC GGG ATG ACC AGC TCG TCT TCG GTC GTG	
ile ser gly thr pro thr ser thr met val his gly met thr ser ser ser ser val val	
5941/1981	5971
TGA TTA CAT CTC ATG GCC CGT GTG TGG GGA CTT GCT TTG TCA TTT GCA AAC TCA GGA TGC	
..A	
6001	6031
TTT CCA AAG CCA ATC ACT GGG GAG ACC GAG CAC AGG GAG GAC CAA GGG GAA GGG GAG AGA	
6061	6091
AAG GAA ATA AAG AAC AAC GTT ATT TCT TAA CAG ACT TTC TAT AGG AGT TGT AAG AAG GTG	
6121	6151
CAC ATA TTT TTT TAA ATC TCA CTG GCA ATA TTC AAA GTT TTC ATT GTG TCT TAA CAA AGG	
6181	6211
TGT GGT AGA CAC TCT TGA GCT GGA CTT AGA TTT TAT TCT TCC TTG CAG AGT AGT GTT AGA	
6241	6271
ATA GAT GGC CTA CAG AAA AAA AAG GTT CTG GGA TCT ACA TGG CAG GGA GGG CTG CAC TGA	
6301	6331
CAT TGA TGC CTG GGG GAC CTT TTG CCT CGA CTC GTG CCG GAA ATC TGA TCG TAA TCA GGG	
6361	6391
TAC AGA ACT TAC TAG TTT TGT CTA GGA GTA TGT TGT ATG ACT AGG ATT TGT GCT ATT ATC	
6421	6451
TCA TTC AAC AAC ATA GAG CAA GAA TAG TGA GCT AAC TGA GCT AGA CAC TCA ATT AAT CCG	
6481	6511
CTA CTG GCT TCA AGT CAG AAC TTT GTC ATT AAT CAT CGA CTC CGG GAC GGT CAT ATA TGT	
6541	6571
ATT ACA TTT CTA CAT TTT TAA TAC TCA CAT GGG CTT ATG CAT TAA GTT TAA TTG TGA TAA	
6601	6631
ATT TGT GCT GGT CCA GTA TAT GCA ATA CAC TTT AAT GGT TTA TTC TTG TCA TAA AAA TGT	
6661	
GCA ATA TGG AGA TGT ATA CAA GTC TTT ACT	

A. Allelic variations: single nucleotide changes (polymorphism) between CLASP-2 cDNA isoforms

Isoform	Difference	Nucleotide(s)	Consequence
1	polymorphism	862	A to G change; mis-sense mutation
2	polymorphism		A to C change; mis-sense mutation changing codon from histidine to proline
3	polymorphism	2210	A to G change; mis-sense mutation changing codon from asparagine to glutamic acid
4	polymorphism	2225	C to T change; mis-sense mutation changing codon from histidine to tyrosine

B. Alternative splices

Isoform	Difference	Nucleotide(s)	Consequence
1	exon deletion	209-291	premature, in-frame stop codon leading to the production of a truncated, most likely soluble protein

These differences may be found separately or together in various combinations in the different human CLASP-2 isoforms

FIG. 11B

human CLASP-2




FIG. 11C

1st exon (nucleotides 335 to 445)

TGTCTTGCTTATCTTTTCGCCCTCCAGGCAAAGCCAAAGCTAATTGAGCCACT
CGACTATGAAAATGTCATCGTCCAGAAGAAGACTCAGATCCTGAACGACTGT
TTACGGGAGATGCTGCTCTTCCCTTACGATGACTTTCAGGTAAGTAACGTTAT
GTTTCTATCCGTAGAACCACG

2nd exon (nucleotides 7101-7190)

TTACCCAAGGCTTTTCCTCCTGTTTTTGTTCAGACGGCCATCCTGAGACGA
CAGGGTCGATACATATGCTCAACAGTGCCTGCGAAGGCGGAAGAGGAAGCA
CAGAGCTTGTTTGTACAGAGGTAAGGCTCTTTCCTGCATTAATTACATTTT
GAAGTCATTTTCCCCTAACTGCCTCC

3rd exon (nucleotides 11439 to 11521)

TTTTCTATTTTAAATCCCCCTTCAATAGTGCATCAAAACCTATAACTCTGAC
TGGCATCTTGTAAGTATAAATATGAAGATTACTCAGGAGAGTTTCGACAGC
TTCCGAAGTGAGTAAGCTATATTATACACATAGGGAAAAGTCTTT

4th exon (nucleotides 13987 to 14056)

CTAAAACAAATTTTCTTTGTTGTTTTATAGCAAAGTGGTCAAGTTGGATAAA
CTCCAGTTCATGTCTATGAAGTTGACGAGGAGGTGCGACAAAGATGAGGTGG
GATACCTGCTTGCTGTTGCTTCTCTTTTCACTCTAGATTAA

5th exon (nucleotides 15212 to 15307)

GGAGGTGACTGCTGGTGTTTTCTTCTCTCCTAGGATGCTGCCTCCCTTGGTT
CCCAGAAGGGTGGGATCACCAAGCATGGCTGGCTGTACAAAGGCAACATGA
ACAGTGCCATCAGCGTGACCATGAGGGTGAGGACGCACATCACTTTGCCCTC
CCCTCTCACAAGCCCTTTC

6th exon (nucleotides 16269 to 16404)

TGAAAGAATAGCTGTGTGTATATTTTTCTCTCAGTCATTTAAGAGACGATTTT
TCCACCTGATTCAACTTGGCGATGGATCCTATAATTTGAATTTTTATAAAGAT
GAAAAGATCTCCAAAGAACCAAAAGGATCAATATTTCTGGATTCCTGTATGG
GTGTCGTTTCAGGTAAATATGAAAAGAGTTTTACCATTATGTTTTCTTA

7th exon (nucleotides 19459 to 19633)

AAGTATGTCTGTTTATCCTTTTTTTCATTTTCAGAACAACAAAGTCAGGCGTTTT
GCTTTTGAGCTCAAGATGCAGGACAAAAGTAGTTATCTCTTGGCAGCAGACA
GTGAAGTGGAAATGGAAGAATGGATCACAATTTCTAAATAAGATCCTCCAGCT
CAACTTTGAAGCTGCAATGCAAGAAAAGCGAAATGGCGACTCTCACGAAGGT
AGATAGGCTTGGCTTCCCCCAGGCACATACACACTCT

8th exon (nucleotides 20567 to 20634)

ATTACAAGTGATTCCGATAATCTGTTTTGCCATTTTAGATGATGAACAAAGCA
AATTGGAAGGTTCTGGTTCCGGTTTAGATAGCTACCTGCCGGAAC TTGCCAAG
GTAACATCGTCTTATATCTTCTGCTCTTCGTTGAATGC

9th exon (nucleotides 30257 to 30331)

GATTGTGTTAAATGTAATTTTCATGTATCTTGTTATCAGAGTGCAAGAGAAGC
AGAAATCAAAC TAAAAAGTGAAAGCAGAGTCAAAC TTTTATTGGACCCA
GATGCCCAGGTAAGAACTATCTAAATGTTTAATATTTAAAACCAAAT

10th exon (nucleotides 31851 to 31991)

CATAACTTATTTATATGTTTACATTTTCTTTTAAAGAAGCTTGACTTCTCATCA
GCTGAGCCAGAAGTGAAGTCATTTGAAGAGAAGTTTGGAAAAAGGATCCTTG
TCAAGTGCAATGATTTATCTTTCAATTIGCAATGCTGTGTTGCCGAAAAATGAA
GAAGGACCCACTACAAATGTAATTTTCATTTTAAAAATAAACATTAATAAAAA
AAAATAGGCAG

11th exon (nucleotides 32472 to 32675)

CCATGGTGATCATTGGATTGTTTTGTTTTGTTTCAGGTTGAACCTTTCTTTGTTA
CTCTATCCCTGTTTGACATAAAATACAACCGGAAGATTTCTGCCGATTTCCAC
GTAGACCTGAACCATTTCTCAGTGAGGCAAATGCTCGCCACCACGTCCCCGG
CGCTGATGAATGGCAGTGGGCAGAGCCCATCTGTCCTCAAGGGCATCCTTCA
TGAAGCCGCCATGCAGTATCCGAAGCAGGTGGGGAGTATGAGCCAGCATTC
CCACTACTCAGACTCACTTTGCATGC

12th exon (nucleotides 33063 to 33185)

GAATTCTGCTTACTGAAGAAAATGTTTGCCTCCTAGGGAATATTTTCAGTCA
CTTGTCCTCATCCAGATATATTTCTTGTTGGCCAGAATTGAAAAAGTCCTTCAG
GGGAGCATCACACATTGCGCTGAGCCATATATGAAAAGTTCAGACTCTTCTA
AGGTATGAATGGCTTTTACGCTTTGGGGTGGTAAAAAGCAATCTGAA

13th exon (nucleotides 36702 to 36784)

CAGTATCTCATAGCTTTATTCTCATGTCTTCAAGGTGGCCCAGAAGGTGCTGA
AGAATGCCAAGCAGGCATGCCAAAGACTAGGACAGTATAGAATGCCATTTGC
TTGGGCAGCAAGGTAAGGAACACCTTTTATACCTTTTAAATCGATATAGATA
GGTGCATGG

14th partial exon (nucleotides 37353 to 37475)

GAAACCCAGTTTAGAAATGTTGCTTTGCCATTTCAAGGACATTGTTTAAGGATG
CATCTGGAAATCTTGACAAAAATGCCAGATTTTCTGCCATCTACAGGCAAGA
CAGCAATAAGCTATCCAATGATGACATGCTCAAGTTACTTGCAGACTTTTCGG
AA

1 TACCAAGGGCAACTCTGGCACACCCCTAAAGTCTGGAAAGGGGACATAGCTAGTCAGGGATGACCCGAGAAATGACTGGAGCTCCACCAGAA
93 TGCAGAGCTTCCTTTGTGCTTAAATACTGAACAAGCATCACTCTGTGTAGCAGACACCACCCAGCATTTTTTGTCCCTTTGGAACAACCT
185 CTTATTTCTGTTCTTTGTGATACCAAACTAGCATACTCTAATTGTAGAAAATACAAACATAGAGTAGAACATACTAAGTTCTTTATCTT
277 AAGAAATGGCATTTGTGTATGAGATGTCTTGCTTATCTTTTCGCCCTCCAGGCAAGCCAAAGCTAANTGAGCCACTCGACTATGAAAATG
369 TCATCGTCCAGAGAAGACTCAGATCCTGAACGACTGTTTACGGGAGATGCTGCTCTTCCCTTACGATGACTTTTCAGGTAAGTAACGTTATG
461 TTCTATCCGTAGAACCACGTGTTGATCTTAAACAAGCAGTATTTTTCTATGTATTGATTATTTGTTGGTTAGTTAATTATTATTATTATT
553 ATTTTTTTTGTAGACACAGTCTTGCTCTGTGACGCGAGGCTTGAGTGCAGTGGTCCCATCTTGGCTCAACGGCAACCTCCGCCCTCTGGGTTCA
645 AGCATTTCTCCTGCCCTCAGCCTCTCAAGTAACCTGGGATTACAGGCGTGTGCCACCATGCTGGCTAATTTTTGTCTTTGTATTAGAGACAGG
737 GTTTTGGCACGGTGGCCAGGCTGCTCCAACTCTGGCCCTAAGTGATCTATCTGCCCTTGGCTCTCAAAATGTTGGGATTATAGGCATGA
829 GCCACTGTGCCCGGCTAATTATGGTTTTTAAAGATGAAATAAGATGTTATTTAAGAAAGAAAAGTTATTTTATATTCTTCCAGCATCC
921 TTCATGAGTGTATTAATTTTAAAGGTTATTTTGTGATTTAATTTATAAGTATGCCAAATGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT
1013 GTGTATCTTGAATAAAGTGTATACCTCTGTCGGGCTGATTAGTGGGCACAAGTGCCCTTTCTGCTTTTGTAGATTGTTTGTAGTTAGAT
1105 TTTTGTGGCAAGTACCTTAGAATCTTAATGATGGGCTTGTCTGGGAGCAGTGGGAGATTTCGTGATGCCCTTTTTTAGGATGGCATTTGGG
1197 AGCCTGCTTCAGAGGCCCTGCACTCTGTGTGGGCTCCAGCAAAGCGTTCAAGGTTAGCCAAGAATGCCCTGAAGTTTACCTCTGTAGTGT
1289 AATGTGGGTGCTGTTCCTTGGAGAAATGTGGAGGACTCAGCACAGCTCCTGCTGTGTGCCGCTCTTTCAGGCTATGGCTGTGGTTAAGAG
1381 ACTAACAAGAAGCTGTGAGGCTGTTGAGGAATGAGAATGACATTTCTTCCAGGAAACCCGGTGGTGTAAATGCCCTTGCAGCGAGCCACC
1473 TTGGTCCATTTGGAGGTTCTGGTTACTTTCTGCTCTCTGGGACCTGATCCTGGCACTTCTTTCTCTCTCTTTGATGCTCTTTAGTTGG
1565 ACACITCTCTACCTGATGCTGTACCATTTAAGCCCTGTCTTTTGTGAATCGAGCTGCCTTTTTTTTTAAAGCTTCACTGATCTTTGTGT
1657 TTGATTCCAAAGTGTGTACATCCATGTACAAAAGATAAATGAGAGGGAATATTGAAATAATTGCATGAAAAGCCCTCCACGCCCTCTA
1749 ATCCCATCCACAGAACATGACTTAACGTATACAGCTCTGTGCATCTTGTCTTTAGAACTTCCATGTTAATAGAAATGTAAATTAACG
1841 ATCCTTGAAGGTTTTTTTCCACCAATTTAAGCGACTCCAGCTTACAACAGAGGTGAGAAATTTCAAAATGTTCACTCTTTCTAAGTTGT
1933 AGAGATAOCTGGGCCCAAAATGATTATTTCTTTAGCTCTGTCTGCTAATAAAGGAATGCCATGGGAATGAAATGACCATTCGTGTGGTGT
2025 TGCTACCAAGTAACAGGTAAATGGGTTGAGGTCATGCCAAACAATACCATGCTTTGCATCTTCTATTTCACTGACTAAACTGCATGGGAACG
2117 GACTAATTAATGAGAACCTCTGAATGATGCTTTTGCCTGTGATTTGGCAACAAATGAAAAGCAAATCAATGATTATAAATGTACTGCA
2209 TGTGTGACAAGATTTTCTGTAGTGTGTCTGAGGAAGCTAAAGGTTATCTCAATTTCTCTCAACATGAAGTATGTCTCTCTTGGTATTA
2301 ATTAAAGTAACAACCTTTTTGAGTTTGAACCTTAGAATGAAAATTTCTATTTGTATGACTGAGATAAAATGCTTAAGAAACAACCAAGAA
2393 ACGAGATACAGTTAGTTGAGTGTATCTTTATCCAGGGAACAGGTATCTGGATGTTAAGCAGTTGCAGAAATGAGAACTCACTAGCATTTTGT
2485 GAAAACCTTCTGTGTCCCTTTGCTTTTAACTACTCTGGTGATAGCAGGCACAAATATCTAGGAAAGGCAAGAACTCACTAGCATTTTGT
2577 GGCTAAGGTGATGAGCAATATTAATTTCTGTTTGGGAGAAAGTTTCCCTAGAGATTAGGAGCTTGAATGGAGCTTTAATCCTCATACA
2669 GGAATTTGTATGGGCCCCAGTGAAGTTTGGGTACAATTAATTTGTTTTCTTATAGACTCCACTTTCTTATCAGGTAAGGCCAATGACTCTGT
2761 GCTTTCTTGTAAATGTCTCAGTGATGTTATTAAGTGTCTAATTAGCTGGATGAGTGAAAGGTCTTAACAGTGCCACAGATCTTTCTATC
2853 TGTGTTTTCTTAGGCAGAAATAAGAGCAGAAATTAATGTATTATTAGAGGCAGAGGGAACAAATTAGATTGGGAAAGTGTTTTATTTCATAG
2945 GAAAAGTAATACCAAGTTGGTTAGGAATGGCAGCAGCAAAACCCATGCTGAGGGGTGATTTACTGCATTAATAATTTAGCAGTATAAGT
3037 TAACTATTAAAATAATAGAACTTGGTGCCATTTCTGCCAATATTAATTGAAATGACAAATTACTAAAATATAAGCATGGATAGTGGTGATG
3129 CTTGTGTACATTTTCAAGTAGGCACATGTTGATCTTGAGCCTTTACTGGTCAGATCCTAAAGGCATCTACATGTTCTCTAAAATGAGTTG
3221 TGTCAAGAAAGATTTGCGGGTGCATGATGTGCTGAGGATGACAGAAAGTAGTTACTACAACAGCAGCAAGAGAGAGACATGAAGT
3313 AAACGTGGATTTTTAAAATCAAAAGAAATAGGCCAGGCGCACTGGCTCATGCCTGTAATCCAGCACTTTGGGAGGCGAGGTGGGCAGATC
3405 ACAAGGTCAAGGATTTGAGACCAGCCTGGCCAAATATGGTGAACCCCATCTCTACCAGAAATACAAAATAGCCAGGCATGGTGGTGCATG
3497 CCTGTAATCCAGCTACTCGGGAGGCTGAGGCAGGAGAAATCTTGAACCCAGGAGACAGAGGTTGCAAGTAACTGAGATCGTGGCACTGCA
3589 TTCCAGCCTGGGCGACAGAGTGAGACTCCATCTCAAAACAACAAAAATCAAAATAATAGTTCCAGCCATCAGGTTATGATGAAGTAGG
3681 CTGGGACCGTGGCTCACACCTGTAATCCAGCACATGGGAGTCCGAGGCAGGTGGATCACTGAGGTGAGGTGTTGAGACCAGCCTGGC
3773 CAACATGGCAAAACCCGCTCTACTAAAAATACAAAATAGCCAGGCATGGTGGTGGGACCTATAATCCAGCTACTTGGGAGGCTGAG
3865 GCAAGAGAAATCGCTTGAACCTGGGAGGTGGAGGTTGCAAGTGAAGCAAGATCGGCCATTGCACTCCAGCCTGGGGGACAAGAGGAAAATCC
3957 ATCTCAAAAAAAGGGAATATTAATGAAGTAAAGTACATGTGATCTGCCATGGCCAGGACAGGAATGCCATGGGCGCTGCAGCCGTCA
4049 CTAGCTGATGGCCCTCTTTTGCAGAACTCAGATCCTGTGCTTGGGATCTCTGCCATCTGTGCTTTGGCTTCATGGTTCTCTTGGCCAGC
4141 AGCATCTTCTCTCTTAGATCTTTCTACCTTTAGAGACCCTTGAATCCCATATTGTCTGAAGCTATTTAAGTCCACAGAACTTTTCCC
4233 CCCATGCTCTCAATTCTTTCTACTGCTGTCTGCACCGTGCACATAAACACTTGAGTATGTGGTCTTGGCTGTTACGACCTACTTCTTA
4325 GGCTTCTTGCACGCGAGCATCCCGCCCGTGTGTGGTCTGAGAGGGCTGGCTTTGAGCCTCTGTCTCCACCCACCTGCCACCTTACA
4417 CATGCACAAAATCCCTTTCTGTAGGTGCTAGGTTGAATACCATGCTTACCTTACTAATAGTAAATTTTACAAGCAATAGGTTAT
4509 TTCTTTGATTTCATCAAGTAAATATTAATACTGTTTGAACATGTGATAGTCCAGCGACTAGATTGTAAAAATATTGAGGATCAATGAT

FIG. 12B
1 of 10

FIG. 12B
· 2 f 10

9201 GGGTGGGGGTTGCTTGCCATAATCTGTGCTTCTGCTTTGGAATTAGCCAATGCCTGCTATGTAGATGGTCAACACTGGCTTGTAAATCPA
9293 TGAATCTCTAAATACTTAGCCAGGTTCACTGTGGAGTTTTTTTTTGTGTTTTTTTTTTTTTTGATGTGCGATTCTTTATAATTATAAATATA
9385 TGAACATAATAAATGTAATTATTTCCCATTTTTGCTAAGGTATATAAATCTTGGCCAGGCGCAGTGGCTCAGCCTGTAAATCCAGTAC
9477 TTTGGGAGGCCGAGGGCGGTGGATCAGAGGTCAGGAGTTCGAGACCAGCCTGACCAACATTGTGAACCCCTGTCTCTACTAAAAATACAAA
9569 AAGTAGCCAGGCGTCTGTGGTGTTCGCCCTGTAAATCCAGCTACTCGGGAGGCTGAGGTAGGAGAATCGCTTGAACCCGGGAGGCAGGGTTGC
9661 AGTGAGCCAAGATTGCACCCTGCCTCCAGCCTGGGTAAACAGAGCAAGATTCCGTCTTAAAAAAAATAAATAAATAAATAAATCTGGG
9753 CTCATTTCATAAATGCATTTATAATATACAAATTTATGCTGGTTTATTTAGTTTTTAATTGGGCAGGTACATTTTATCTTCAGATAGTTT
9845 TAATGCTCTGCAGGAGACTTGGTGGTACAATTTCACTTCTAGAGCTTGTGAAAAACCGCTGTAGCTCTTTGAAGAGATAGGATCACACCC
9937 ACAAGACCACGCTGTACATAGTCAGGCTGACTACTTAGCTGTGCCAGTGGACCTAGGGGCAGTAGTGGGAAATCCAAAATATGGATTAT
10029 TCTGAAATCTGACTGGTTTTGGAATGAAATTTGAGTGTGAATGTATGATACTCAAAATGTGAAATGCTTTGAGAAATCTAAAAAATCAT
10121 TTTCAAGGATAAAATAGCATTTTAAAAATTTCTTAACACAGAGTAAAAAATCAAAAAGTTTGCAGCTCCAAAATCTGACTTTGCGCTTA
10213 GCGCTTCTCTGCCATTTATTCACCTCTCAGATAATTTTAAAACTTACATTACCTTCACATCACACATCACACCTTGGCAGTAAAGTGC
10305 TTATGTAGAGCCTGGCACAAAATAAGCACTTGTCTGTGTTGAATCTTACTTTTCATGAGTTTGGAGGTGGAGGGAGTTTGCCTCAGAG
10397 ACCTTCAGGCATGGCCTGTGACACAGATAGAAAGGTTGACAGTAGAAGCATGAAGTGAAGAAGCAGATAAGAGACGTTGTTTCATGTAGT
10489 TACAGCCAGAGCCAAATAAATAGATAAGGCATAAATACATTTATCGACCTATGTTTTCACTGATTCATCTCATGCTTTTTTCCACCCACAC
10581 TGAAGTCTGCACATAAATAGTCAAGTCTGACTGGACAGAGATCAATTTGAACCTAAGCTGATTTTAACTAAAATCTTTAGCTTTTGAATTT
10673 GACTCATTCATATTTGAAGATCTGGTGGATGTGCATGTGCGCAGGTAGTTTGAATGCAGTCTGTGTAGGTGAGGTCACTAATAGAATCTTGGCAGTG
10765 GACACCAGGCTGGCCCATTTGAATCTGTCAATGTGCGCAGGTAGTTTGAATGCAGTCTGTGTAGGTGAGGTCACTAATAGAATCTTGGCAGTG
10857 GAAGACTACATTAACCATTTCTGTACACACACACAGATGAAATGTCAATGGTCTAATATTAAATGGAATGCATATGTTCTCTCAATTAAT
10949 TCACCTCTTTCTTATGAATGGAAGAAAACTGAGGGCAGATTTATATGGTAATTAATACTAACAGAGCTAGCTAGTCTCAATATAGTAA
11041 ATACCATGAAGGCGCTCTTAAAGTTCACGCATCTTATTCATTTTATGTCTAGTAGAGGCCACATGGCTTTTAAAGAACAGGTGCTTACTC
11133 TGAAATAGTCCATTGTTAAGTAAATGTCACAAGGTTAGGTGAAGTTGTCTCTTGTAAAACCTGCTCTCAGATCATTAATGATGATAACTTA
11225 AAGTGATACTACCCCCAAGGTAATGTTTCAGTGGTTCAAAGTCTCAAGCTTCACAGGAATCTCTGGTGGTGAATTCAGGCAATAGCTCG
11317 TTGATGGGAAAAAATTTTTTGTATGTTCTATGGGTATGCCCTTCAAACTTTCCCAAGTGTTTTCCAAAATCACGCTTTCTGTTTTTA
11409 TTTTCTATTTTTAAAAATCCCCCTTCAATAGTGCATCAAACCTATACTCTGACTGGCATCTTGTGAATATAAATATGAAGATTACTCAGG
11501 AGAGTTTCGACAGCTTCCGAAGTGAGTAAGCTATATTATACACATAGGGAAAACTTGTGACTTGAATGCTTGGGGGAGGTATGTAAT
11593 TCATATGCAATCAGAGTAATGAGGAAAAATTTTTAGATGGTTTATGTATGTGGTGAATGTTTACAGGGCTTTTATTGTAAGA
11685 CTCAAACATGCTACTTTGGTATTGATAGGTAGTAATAACATGTGGATGGTTTAAATATGTCCAGTGGTTCTTTTCAGGGTACCCTGATAA
11777 ATAAATAGGTGAAATTTTTCTACTACAAAATATAAAACAAACAAAGCTAAGTGAATGAAATATGCTGCAGTGCATTTTTTCTTGAAG
11869 AATATTTTGAAGAAGATAATATAAGAGCATTTTTACATGAAATAAATTTATGTTTTTAAAAAAGTAAATCAAGAAACAAGCATTTTCCC
11961 AGTTAAATTTTTTTTTTATCTCCTTTAATGTATTAACCTTATCTAGACTATACCAAAGCAAGTGAATTTAGATTGAATAGTTGTGGCCAA
12053 GTGAATCGCGGTAGCTAGGTATGCTTTCAGACACTATCTTTTATAAAGGTTCCATTGTGTGCTTTTAAAGGAATACCTGAGACTGGGT
12145 AATTTCTAAAGAAAGAGATTTATTTATTTATTTTATTTATTTAGTTTGTAGACAGAGTCTTGTCTGTCCAGGCTGGATACATGA
12237 CACGATCTCTGCTCACTGCAACCTCCACCTCCCAAGTTCAGCAACTCTCCCGCTCAGCTCCCGAGTTGCTGGGATTACAGCCACTTGCC
12329 ACCACGCCCGCTAATTTTGTATTTTATGTAGAGACAGGTTTACCCTGTGTGGCCAGGCTGGTCTCGAATCTCTGAATCAGGTGATCCAC
12421 CTGCTCTGCTCTCCAAAGTGTGGGATTACAAGCATTAGCCACCGAGACTAGCCAAAGAAAGAGATTTATTTGCTTATAGTTCTGCAGGT
12513 TGTACAGAAGCATAGCACTGGCTTCTGCTCAGCTTCTGTTGAAGACTTTTGTGCTGCTTCAAAACATGATGGAAGGTCAGAGGAAAGT
12605 TGGCACTTGTGAAAGAGACCAAGAGGAGGAGAACTCACTTTATAACAACCCATTCTCTTGGGTACTAATCCATTTCAGCAACAAGTAA
12697 TCCCATCTTGCCAGGAACAAGATTCATCTACTGTGAGAACAGCACCAGCCTCTCATGATGGATCTGCCCCATAACCCACTAGGCCOA
12789 ATTATGCAACAACGGGGACCAATTTCAATCTGAGTTTTGGTGGGAATAAAACCATATCCAAACCATAGCAGTTGCCAAATGAATTATAC
12881 TTGATTTTGGGAAATTAAGCAAAATAGTGATGGATTATGTTTTAAACTTAACCATCACCATGAAAATATTAATGAGACTGATTATAG
12973 TATTTTCTTTGCGTTTGTATCATCTTTGAGCTGGAACCTTTATGCGGTTCTCAGTAGACCTAGCTGTTTTGTTTTCTCTCTGTGGCT
13065 TTGCCACTCTTAAGAAATGTTGGCCAAATCTCCCGATTGCCCTTTTTTAAACCTCAGCCAGGAACACTCCCTCCTAGTATTATCTCTCCAG
13157 ATGGGTAGCCCTTTAGTTCTATATTTACCAATCTCCTTTAGGGATTTTTAATCTTCCCATTTGGATTGGCTTAACCCATCTTTGTGAT
13249 CCTCTGTTCTAGTCTCAGGTTGAAAGATATCAGACTATGTCATGTCGTATACTTACTATCTAATAGACTGCTGGTACATTTCTCTCTT
13341 GGCATTAATAGAAATTTCCAAATGTGATGAGAAAGAGAGGAGAAATGTAACAGTGGTGAAGAACACAGATTCTGTGTTCTGCTCAGAC
13433 GACTCTGCCGCTCTGTCAGCCTGATTTCTCTCATCTGTTAAATGGCCTTAAACAACAGTCACCATGATTAAAGGATTAATGAGAGGGCAGATG
13525 AGAAGTGGCAGGGCTAGCACGTCAATCCATTGAGTAAATGTAAGCTGCTTATGGTATTGGTGTCTGTTTTGTGTGGTTAATACCA
13617 TCATTGTTAATCGGTTTCAACGCAACAACCTCAATCTCTTTTTTCTCATAACTTTGTATAAAAAGTTATCTACCAAGCTTTTGTGTTA
13709 TTAAAACTAATCCACTTTCTTATTTTAGTATGCTGCTAACTCCCCAGAGCTATGCTGTCTTTTCCACATAGCTTTTGGAGCTTTCTT

FIG. 12B
3 of 10

13801 ACTCAAGTCTCTTGCCTTACCCACCTTGAAAAGCAAGGGCATAGATGGTTTTATTCTTTGCTGATAAAGAAGCTGGGCCATCTTTGGATT
13893 TAGTAAAGCCCTGGCCTATGATGGAGGAAGAAATGCAAGCCCTCTTCTTGACTAGGCCATTTCTAAACAAATTTCTTTGTGTTTTAT
13985 AGCAAAAGTGGTCAAGTTGGATAAACTTCCAGTTCATGTCATGAGMTGACGAGGAGGTGACAAAGATGAGGTGGGATACCTGCTGCTGT
14077 TGCTTCTCTTTTCACTCTAGATTTAAACATCAATTTTACAGACTTAGAAGATTAGTTAGAAAATTACCGACATTTAGCCAAAACAGGCATTG
14169 GAGTCTTACATGAACGGGATAATTTTTTAAAAATGTTATGATTGATTGGAATAAGGTCTCTGTTTCAACTTTACTGCTTAGCATTTTCAT
14261 GTTCTCTTGGTTGTGTTTTATTGTTCTGAGATCATTTTCAAAGACTTGGATCAGATCTGGCTACATTGTTAAAGATATCAAGATGACTTAG
14353 ACCTTGAAATTAGCTTGTTTTTCAACAGATCTCGAAACAGCTGCCAGCCAGTAGATTAAATGGCTATTTCTTCAATGATTGCTTTTAGTGA
14445 AGTCTGATTGATCAAGGCCACTCCCCCTATTCTAGAGGAAAGCTCATGGCTAAAGAACTATATAAGGGAGTAGGGCAITGAGATGAGTC
14537 TGCCCACTGAGTGAAGGAACCTCACAAGAAGACAATGCCCATCTCTGCATTTCTCATCTCCCCATTGATTGTTAAGTGTCCCAITGTGAG
14629 TTTAGGTTTTCTCTCTTTAAAAAATTGTGAGTGAAGTATACCACTAGCCACTCATTAAGCAATGTGCATGTAGCAAAATATTTTTATTTC
14721 CCCCCATATTTTATCTCTCTTCTGATTGCTCAATTTCTCTCCCTTGCTTTATTCACCTTTCCCTGAACTAAGCCCTCTGGGAAGGTTTC
14813 CAGGAATGTGATGCTCTTTCTCTCTGACTATAGGGGAGTGTCAATTTGAAACATTTTTCGTGAACCCAGGCAAGACCTTCCAAAGTGA
14905 GTGGTCACTTACCTAGCTCTCTTTCTGCTCTTTGTTGGCTCATTAACACTGACAAATAAAATTTGGACAGGAGCTAGCTTTGCCCTTTAA
14997 TGGAAATAAGTTTTTCAAAATGAGGGGGTCTCTCTCTTTCAACGCTAAGTGGACTTTTATGTGACTTGTAGGCATTGGTGTCAATGGGTG
15089 CTTGAGTAAATTAATGGAACAATTCGCACAAAGGGAATGACCTTCCCATTTGACCAAACTCAGCAAGCAACCCAGGTAATAACGGGAG
15181 GTTGACTGCTGGTGTCTCTCTCTCTAGGATGCTGCCCTCCCTTGGTTCCAGAAAGGGTGGGATCACCAGCATGGCTGGCTGTACAAAG
15273 GCAACATGAACAGTCCCTTACCTTACCATTGAGGGGTGAGGAGCCACATCACTTTGCCCTCCCTCTCACAAGCCCTTTCTGCCATAGAGCT
15365 CGAGAACATCTCTGAGTAAATGATGCTGTTCTTCCCACAAAGGGACATTTGCTGATTCCTAGGATGCTCCCTGGTGTGATGAGCACC
15457 CCCCATTGGCACAATTAATCCACCACTTTCCCTGACTGCTCTTCTGACCACCAAGGAGACATCCCTGGGCTGGTGTGAGGTGCGAG
15549 ACACATGACATAAGCTTTCTCTCTGTAATAACTGAAAGTGCTCTTTGGTACCTCAGCAATGTCAACAGGGGGCTATCTGTCATGCCAATC
15641 CTGAGCACTTCTGTGAATGTAATGAGCAAAAGTCAAGTAAAGCAAAATTTAGGACGAGAAAAGAAAATAGTTGCATAGAAAGAGAGGTT
15733 GCAGACAGAGAACTCAAAACATAGAGAGCTATTGAGGAGAAAGTGGGACAGAGGAACATCAGGATTAATAACAAAGGGAAGAGAAC
15825 AAGGGAGTCAAGGATAAAATTAAGTAGGAATGTGACTGTCTTACCCTAAGGCTGGAATATCATTGAGCGTCATGAGGCAAAAATAG
15917 TTCCCATTTCTGTGCAAGAAACCTCGGATTTTAGAGAAAGTTCTGTCTTCTGTGCTGCATCCCAATTTGGAAGTCCCTGCACTGCTT
16009 TTGGGTAGTTATGTAATAATCTCTGATTCGCTGGGTGAGAAAATGACCCATGGATATTAGGGGAACACCTCCTCAGAACTGAGATGCACTG
16101 AGCTTCTTAGATGCTATTAAGTCTTGACCCACAGTGACCTGGAGCATCAGCTAGAGTGAACCGAAACAGGTTTTATGTATGTATGTA
16193 GTCATAAGTGGTTATGATAGAGATTGTGACCTCTTCATTTTGAAAGATAGCTGTGTGATATTTTTCTCTCAGTCATTTAAGAGACGA
16285 TTTTCCACCTCATCACTTGGCGATGGATCCTATAATTTGAATTTTATAAAGATGAAAAGATCTCCAAGAACCAAAAGGATCAATATT
16377 TCTGGATTCTGTATGGGTGTCTGTTGAGTAAATATGAAAGAGTTTACCATTATGTTTCTTATCTGCACTAGTGCTTATGTGTAAATTA
16469 GCAGATTTAAGCAAACTTCAAAAATGGCAATATGATGTTAGAAATATAACATATACTTTAATGAGGCAAGCCTTGTTTTTCATCAT
16561 TGTAGAAGATGGAGTATTAATGTAGAGGCAGAAATATGCTGTGGCAGGCAGGAGCACTCTGGCTCGGCCACTTTATAGCTCGGTGACCTTT
16653 AACAGGCTACTTAATTCAGATAATGAGATGTTTCTTTAATACGGCAATGAGTACATTTGGATGATCAGTGAGGAAAATATTTAAACAC
16745 TTCTATAGTATCTCACTGCTGATTTTATCGCTAGCATTTAGTACAGTGGCGGTGATGATCAGTAAAGAGATTAGGTTTCAGCGCAGATTG
16837 AGTTCAAAATCCCTGCTCTCCACTTACCAACTGTGTAACTTGGAGATGTTATTTAACTCTCTGTACCTCAGTTTCTTCATTGTTAAATA
16929 AGGATAATGGCAGTACCAATATGGTACTGAGAGGGTTCAATTCATTACACATGTAAAAGCTTAGAACAGTGCCAAACAATGGTAGCATT
17021 TGGTCAGTATAGATAGTTTTGTTATCATAGGGCTGTTGTACTTTATATCATAGGGCTTATGTACTTATCTTTAAAAATTATGTTAATTA
17113 AAGATAACACATGAATGATTTTTCTTGTAAAAATCAGCCAAATACAGATAAAGTGAAGTCTCTGAGACTCTCCCTCCTCAGTGTCT
17205 CTTTCTGAGGGGAGCTACTACAGTTTTGCATGCATCTTCTGTAGCTTTTTAGCATGTCTTTGGAAGAGAGTTGTCAATTTCCCTGTCC
17297 ATCATCTGTCCATCCATCCATCCATCCATCTGTCCACCCCTCATTTCATCCAGCCTTGCCACTTTCAAGGAAGATTTAAGGCAGCAGC
17389 TTATAAGCATAACACAGGACATGGGATAGCATAAATTTAAAGTGGGGGTGAAAGCAGAAAGATGAACAGGGGATTTGGGATAAGGGTGAGAGAA
17481 AATAGAGTTAAGGAGAAAGCCTATGTTTTGAAGATCTAACACCTTGCTGTGGGTGGGCCACCACCTGGGCTCTATGCTTTCTCACTTGGAGAC
17573 CTGTTTAGTCACGCAATTCACAGTGCATATGAGATAAAGGCATGATGCTGTTTAGTCCAGCTCTAGAAGCACCCCTGACTTTAAAAAGAGT
17665 TAAAGCAAACTAAATGATTTTGGCAACCTCATTTTTTAAAGTAGGAAGTAATTTTGTGTTTATAAGAGAGTTTTGCTGCCCTGTTCTGT
17757 GCCCAGGGCAGATGTTTATAAGTACAACCTGCCCTGAGCTATCAATTAGTCTCCGGGGTGCAITTCAAAATCTAAGGTTCTGACTTCAATGG
17849 AAGTCTCTTCTTCAAAATGCTTTGCGAGATGCGAGTGTGTTTCAATTTAATAAAGTGTATCCAAGGCTTCAAAAAGTAAATAAT
17941 TGTTTTTATCTGTCTCTGTTTGTAAACTAAGCATCAAAAGTGTGATTTAAGTGTTTTTAATAATTTATCTATGGAATTTATAAATAAT
18033 TAGTTGACTGGTGTGTGAATTAATAAAGTGCCTAAACTAAAAAATTTTGAAGCATTTAGAACCTTTGAAATTTATATATCTTTATTITGC
18125 AGATGAGAAAATGAGGCTCAGAAACAGAAATTTAGAATTGAGGCCAATGTTTCTTCCACTTTTAACTTTCTCTTTTCATGATTGTGA
18217 GTATGCGGGGAAAGGAGGAGAGAAATTCATTTTGTTCAGGCCCTTGACTTCTTCCCTGGTTCTTGCTTGAAGTTTAAAGTGAATCCAAA
18309 GTGGCAATTACTGAGCCACAGCGACAGTCTGTGCACAAGAGTGTGTGGCTTTGCCAAGGGGAGCACTTGACTTTGCATTTCTAAGAACTG

FIG. 12B
4 of 10

18401 TGCTGCAGAAATCACAGAGACTTTTGGGAGGGTTCCTGTCCCTGAGACCTCCACCAAGGAACCTCTTAGAGAGAGTGTGGATAACCCAGTAG
18493 GATTTTAGTGGCTATGCGGGGGCTGTCTGCTGGCTCAGGTTAGTGGGAGTGTGTTGATTTTCATATCGCTCAGCCTGTCTTACAGGGGATC
18585 TTGTGCCATGATCCTCAGAGCTGAACCTCTGTCTACTGCGGCCAACCTGGGGAGATTTTGTCCCTGGAGGACATCTTGAATGTCTGAAG
18677 CTGGCATCTATTGGCTTAAGGCCATAAATTTTCGCTAAACATTTGACATGTCATGACCGACCCACTCACAACAAAGAAATGGCTGCCCAAGTG
18769 TCAGTAGTACCGAGATTGAGAAATCCTGSCCTAGTGCATGTTCACTTCCGTCTGTACTGCACATGGACTACTGTTCTTGTCTGTGAGCC
18861 AGTCACCTCTTTGCAGGCATGAAACCTGGAGGCATGAGGCAAGGCCACCGACAGGAGTCCAAATACCTTTTGGGATTCTAAAGGATGGGA
18953 AAGTTCAGATAAGTAAGCCAAACATAGTAATAGATAATGGTGGCTTTTAAAAATGTAATACCATACACTACTTTCATTAATAAATAGGAG
19045 CTGAGAAATATGAAAATTTTACATGAATTTTCATTTATTCAACAAATATTTTCAAATACCCACTATGTGCAAGTCACTGTAGAGTCCATA
19137 GAGACTAAGGATGTGTAGCACTGACAPAAATGGGAGCACTGAGGAGGTTTCATTCCACTGCAGGGACACACAGTGAATCAGATGAGTATGTA
19229 AAGCAGCTAATGAGTCAGAAAGGAAATAAAGCTTGCAGAAAGTGAAGCAGGGAAGGTGGACGGTAATGGGATTTCATGGGGGGGGCTTTC
19321 ATGATAGGGGGCAAGCTATTTTAAATAGCTTGGTCTAAATGCCAATGAGATATCACTCACCACAAGAGAGAGTAATATTTTAAAGCAG
19413 TTCTAATCTTTTAAAGTATGTCTGTTTATCCTTTTTCATTTCAGAACAAAGTCAGGCGTTTGTCTTTGAGCTCAAGATGCAGGACA
19505 AAGTACTTATCTCTTGGCAGCAGACAGTGAAGTGAAGTGAAGAAATGGATCACAATTCATAATAGATCCTCCAGCTCAACTTTGAAGCT
19597 GCAATGCAAGAAAAGCGAAATGGCGACTCTCAGGAAGGTAGATAGGCTTGGCTTCCCCCAGGCACATACACACTCTGTGGGTGTCTTTATTT
19689 TTGCCAGGTGGGTATAGAAGGAGACCTGTGTTACACAAGTACATGAGAGGTGGGACGGATAGGAGCTCTTTACAAATATCCTGTGACAAA
19781 GGTTTTGTCTCAATTATAACTTACTTCCCTGACATTTCTGATATGGAATCATGTAATGGGAAGAACCAAGCTTTGGAGGCAGAAAGGGAGA
19873 CCTGGGTTTGAGTGCCATAAATACTGTATTTGAGCTGTGTAGCCCTGGGTAAACAACCTTATGTTTCTGAGCCTCAGTGTACTCACTTAA
19965 AATGCAATAAACAATGAAATTTGCTGGGAAGATGGGAAGTGTAAATAGAAGTGAATCTCAAGTATCTGGCATAGAAATTTACTGTATTA
20057 AAAATATTAGTAATAATTAGAATGCATGGGAGCCTCAGATTAAATTTGGTGAGAAAAATCTGGCTATGTTCTTGACAAATCATGTTTACTTC
20149 AACCCTTAGGTGATCCCAACCCCTGGCTTCCCTTAGAAGTACCTGGGAGCTTTTAAAAATAACATTTACCTGGTCCCAAAAAGATTCTG
20241 ATTTAGTTGGTCTGGGGTGGAGCTGGGAGGCTGACTTTTAGGGGGTCTCATGGAGCTGTCCATGTGGGCTGTGTTTCATAGCTAGTGTCT
20333 AGTTCTAATTGGACGGTGTCCATGCTATACCAGCTGCTCAGTGTGTTGACTTTTCATCACTGAGCCTGTGGATCAGTATTTTTCAAAGCACC
20425 CCAAGTGTTCAGGAGCATCCAGAGTGGGGAACCACTGTGTTCAATTTGAAGGCACCTAAGAGAAACGGCTTCCCTCCTCTGTTTCAAT
20517 GAAATGCTATGAATTACAAGTGATCCGATAATCTGTTTGGCATTTTAGATGATGAACAAGCAAAATTTGAAGGTTCTGGTTCCGGTTTAG
20609 ATAGCTACCTGCCGAACCTGCCAAGGTAAACATGCTTATATCTTCGCTCTTCGTTGAATGCTGTGGAAGTATGTCTCATTTTCACTGGTT
20701 TGTCAGAAATGGAATCTGTTGAAATCATAAAAAATACATTTGTGATTACCTCTCTCTTTTCTGACCTGATTACGAGGTGACGTGTACTCATG
20793 CAGTATGATTTAGGCTGTCTTCTAAAAAGTACCTACAAAGCATCTCCTTTTATTATTATTTTAAAGTGTTTTTTCCCTGATAATGCTT
20885 AACACTGCATCACAGGTAAGTGAAGAAATAACTGAAATATGCAGGCAGATGTTCTCATATAGCATCTGACTTTCTATGTTGATACATGTGCT
20977 CTCCCTTACTCAGGTAATAGACAGGTTCCAAAGAGGAAGGACCTGGTAATCTTGCACGAAACCCGGGGTGTGCTGAGTTACAGAAAT
21069 GTTTCGGGTCACTCTTACTGGAAAAAATAAGCTATTCTGTGTCTTACAATTTTGAAGTTTGAAGTTTGAAGTTTGAAGTTTGAAGTTTGAAGTTT
21161 GCAATCAACCATACTGCTACTTCCAGATTAATATCTATTATGATGTTGCTTTTGAAGTTTGAAGTTTGAAGTTTGAAGTTTGAAGTTTGAAGTTT
21253 TTATAGTTGAAGTCTTTTAAACACTGTCTTATTCOCATCTCTTCTTCTCCCAACCCCAATCAGAAACAAGCACTATTAAAAAGTTT
21345 ACTTTCATTTTATATCTTTTAAACACTGTCTTATTCOCATCTCTTCTTCTCCCAACCCCAATCAGAAACAAGCACTATTAAAAAGTTT
21437 TACCATATGATCTGCACTTTTAAACACTGTCTTATTCOCATCTCTTCTTCTCCCAACCCCAATCAGAAACAAGCACTATTAAAAAGTTT
21529 CATTTTTTCTATTATAAAACAAAGCTACAATGACTGTCTTGTACTTGTGTCTGTGTATCTGAATGATTCTTCTTAAATGAGAGAAA
21621 TATCTTTGTCTTCTTAGCCCTTTGCACTCTTACTCTGTACTGCTTCTTATCTTTTGTACTAGAGTGAATGGCGACCTCCACAC
21713 CCACATCTTAAACACTATAATAGAAACATGGTTTATCTATATAGGATTATAAATAGACCAGCAATCAGCAATGACCTTTATTTTAAAGCAAC
21805 ATGGCTGTCTCAAGTGTAAATCTCCTCCTGGCTAGGGCTTTAGAGCAATGTTTTCTTTAGGACTTGACTGCTACCAAGTATCTTTT
21897 TAGCACTGCTTATTAAGCTAATTTTAGTGCACCAATGTAAACCACTCCTAGTCTGGGAAGGTTTGGCTTGTGTGTTTGTGTTATGA
21989 ATGTCTGTGATCATATTTGCAATGCAATTTGCTTTTGTGTTCTGGATGTTGGGGTTTCAATTTCTCAAACAAATATTTGTGCCC
22081 ATTTGGGTTTAGTTGTGTGAGCAGGTAATATATGTATGCCATCTAGAATTGAGAAAGTAACCTTCTGCACTTACTGGGTGAACGGAAATG
22173 GATACCTAGGAGAGATTGATGTTTGTAGCCTAATGTTGATTAATTAATAATCTATGCTTTTCTATGAGGATATACAGGAACGGTCCC
22265 CTCCTTCTTACTACCCAGCCAAATATAATTCAGTATTGTTGATCCCAAGACCTAGGGAGATTTTAAAGATATACATATATTTAATATAA
22357 ATGTATACATTTATGTATATATACTTTTTTATAAGTATAATGTATATTTTGTCAATTAATAATATTTGGAACCTATTTTAAACATAGGGTTA
22449 CAAGTTAGGTTAAGCCATTTTAGTTGGTGAAATCAGTTTGAATTCACCCCTGCTTCTTTTGTGTTTGTGTTTGTGTTTGTGTTTGTGTTT
22541 TTATTTAGGTTAATTTGCAATAGCAGAAATGCTCAACATGAATTTAGAGCTCACTGGAGTTTGGATTTGTACACTGATATAAGCAGCCCT
22633 CAGGCTAGCTTGTACTGAGCCCTCTTATTTAGCTTAAACATGTGATTCACTTTAAGAAGTTTTTTTAGGTCGGGCATAGTGGCTCATGCTGT
22725 TACATGTACTCTTTGTGAGGCTTATTTAGCTTAAACATGTGATTCACTTTAAGAAGTTTTTTTAGGTCGGGCATAGTGGCTCATGCTGT
22817 AATCCAGCACTTTGGGAGGCTGAGGTGAGCGGATCTTTGAGGTTAGGAGTTCAAGACCAGCCTTGCCAAACATGGTATAAAACCCCTGTCTC
22909 TACTAGAAATACAAAATTAGCTAGGCGTGGTGGCAGGTGCTGTATCCAGCTACTTGGGAGGCTGAGGCAGGAGATCATTTGAACCTG

FIG. 12B
5 of 10

23001 GAAGGCAGAGATTGCAGTGAGCTGAGATCATGCCACTGCCTCTAGCCTGGGTGACAGAGCAAGACTGTGTCTCAAAAAAAAAAAGGGTC
23093 CGTTTTATGAAATAAAATGGAATGGAGAATATGAAAGTACACTGCCCTTAATATGACATTATTTTTATATAAAATACTGTCAATTATAT
23185 TTTGGTGGCACCTGCCACCATGCCCTAGCTAATTTTTGTATTTCTAGTAGAGACAGGGTTTTATACCATGTTGGCGAGGCTGGTCTTGAACCTC
23277 CTAACCTCAAAAGATCCACCCACTCAGCCTCCCAAAGTGCTGAGATTACAGGCATGAGCCACTACGCCCGACCTGAAAAAACTTTTTAA
23369 AGTGAATTACATAAATTTTTTACATAAATAATGTCAATTATTAAGGGCAATGTACTATTATACATATAGTGTGTATGTGTCTTGCATAGT
23461 GATATAAAAGATATTTGTTTTCTTAGTGTCTATTATGTATATTTATTTACTTTTCATTGGTATATAATGTACCTATTTTGGGAGTTCATGT
23553 GATACTTTGATATCTGTATACATGTGTGTATGATCAATCAGGATAATTGGGATAGCCATCACCCTCAAACTTTATCTTTGTGTTGGGAATT
23645 TGAACATTTCTTACCAGGAGTCATGGTCAAAACCTGAAAAATGAATCCTTGTGTAGAGGCTTTTACTCTTTCCCTGGCTTTGAGGTGTTT
23737 TACAAATACTTTTATTTAGGAAGGTAGAAAGGTGAAAAGTAATTTTTTGAGGGGAAAAGAATGAAGAAAATGGAGATGAGTATTCACTCA
23829 GCACATGGGTATCTGTGGGCTTTGCCCTTTAAAGCCAGCTTTGGTGTGAGTGTGAGCAGCCAGGCAGTAAGGGGAGACCTGTGTCCCAT
23921 CCCCAGCCTTGAGCAAAAATGCAGTTTGGCTGTTTATCATCCCTCTCAGGGTGTCTGAACATTTGCACCGGTTGAGAAGCCAAAGAAGT
24013 TGACCTGATAACTGTTGGTCATCCCATTAGGAAGGATGGATTCATGCTTACAGAAATCAGAGACTGAAGTATGCAGAGGGAGGGGTGGGAG
24105 AGAGAACTGTGCAAGCAGTTTACCCAGGGTATGAAGAGGTAAAGAGGTGAGTATCAGGGAAGGAAGGTGCAAGAAAGGTGAGGCTGGGAGG
24197 CTGGGCCACAGTTCAGTAAGATTACAAAGAAGGGCCTAGAACAATGAGGGCAGGCAGAAGGTGGCTGAAGGTGTAATTTCAATGGCAGGTTC
24289 TTTTCTAATCAGCTCCTCTAACCTCCTTCATCTGTGCCCCGGCTTTTGTTTTCCACTGTGACTAAGACATAGCCAAACAGGATATGACCG
24381 ACAGGAAGTTGTTTCAGTGCAAAAATACTGATGTCTCATTCTGGAATATTATGGAAGGGCTCAATACTTACAGTGTGAGTGTAAACCC
24473 AGGTTTTCAGAAATATTTGTATAATCTTGGAGCTTATGTTGTACATTTAGTACTGAACATCTGTATTGTTTTCTTATAGAGACACACTG
24565 TATTTACCTTAAACTGGTCTTTTCTCTCTATGTCTATTATGGAACCAACAATTTTATTGTAAATGTAAAGTGTGTAGCATCAGTCT
24657 TATAAATATTTTAGTTTGATACAAACCGTAGTTCAAGTTAGTTAATTGATTTCTTCCCTAGAAAGTCAAGGAGTAACATATATCAGCTTAT
24749 AAACCTCATTACTAGTTATTTAATAATTTATTTCTCTGGTTACATTTATATCTTAGGTGACATCAGAACATATATGTCACCTCCTTAAAGAT
24841 AGTGTGAAGAAAACCCACCTTATGTTTTCTTCCACAGCTTTTCTGTTTGTGAGCTTTTATTTTTGTACTCAAAGAAATAGCATCAACTTTTA
24933 CTTTGGTTTCCCATGTGGTCTGAAAGAGAAGTAGAATTTCTTAAATCCGGAATTGCTCACATCCTTTTACCTTTTAACTTTGTTTTAA
25025 GCAATGAACCTTATGTTCCAGGTAAATCTTCCACAGTTGCATGCAAGGGAAAGTATGATGTCTCAGACTTTATAGTCTCATGGAGATGGAG
25117 TGAGGATCAAGGGCCATGCTCAGCAGAACTTGTGAGGACCCAGCAGTTTCACGGACACCTTTTCTTAAATTTTAAACCAAGTCTATAATAAG
25209 TGCTTTCTTCCCTAGATTCCAATCCAGAAACAATATCATTCACATTTATACAAAGGAGCTGGCTAGGCTTGTGTCTGTGGGGTCAAGCTGG
25301 TGTTCATTTCTGGGCTCCTTTGTGAAGAGGATGAATGATGGTCTGAGAAGTTAGGTGTCTTGAAGTAGTGGAAATAAATCATGATAA
25393 CTCTTTAATTAAGATTATATATTTTGGCCTCAAAACATTTTGCAAAGTCTCCTATTCCAACCAATCTGTTTAAATGACCCAAACATTCA
25485 ACACATTGTTTCTGATAATTCATCCTCAGAAATAGATGCTGTGGCCATAATCTTTGTCTCTAGATTGTTTTATCTACTCGAAATAAATTT
25577 AAGACACAGAGTATGCCATAAGCCTACAGCAGACTTTCTGAAACTCTTGAATGTTTGGTCCATAACTACTTCTTAAAGCAAGAAGAAAC
25669 CTTGTGAGGTGTGTCTATTAGTGTGATGTAGGGTTTACAGGATGGGGTGGGGTGGGGGAATCGCCCTTGGTTTAGATGAATCATCTTT
25761 TCCTTGTCTTCTCAGCAACACCAAGTTTCTACAGAGAACAGCTCTGCCATTGTGCATTTTCTGTCTCCATTTTCTCTCTCTCTCTCCA
25853 CGAACCACAGAGTAGTCAAGTGGGCTTTGGGCAGGAAAGTGGCAACAGGGTGTCTGGGGAAAGCCAGTTGGCTCTTCTTACCATCACAATAT
25945 AGACTGACCACAGGTATTTTAAAGAGCAGAGCTGGTTTCCATCCTCTGAGAAGTGTCAACTACAGACTTTGGGATGATATTTGTTATAGC
26037 TGTATTTTCTCCACTCTTAGATTGTGAAAGTACATATTACAAGTATTTATTTTATTATCTTTACTAAAATTTTAAATAAAAAGAGCGTGCT
26129 TGCCGCAATAAGTAAAAATACCCAAAGTTGTTTAAAGAAAAGTTACCTTTTCCCTTCATCCTCCATCCCACTTCTGAGAACACTGAAG
26221 TTAATAACCGGTTGCAATCCCCCTTCCACCAACTGATTGCTCATAGAAATATAGATAAACATATGAAGGTTTTTAAAGTTTTTAAATAAA
26313 AATATGTTCAATGATATATACATTATCTGAAATTTCTGTATCACTTAAAAATATTTTATAGATGTCCCTCTGGGCCAGTGGAGATCTGGT
26405 TCCCCCTTACATACATATCAGCAAGCTGCATGATTTTCAACTATTGCTACTGCACAGTTTATTCAGCCATCTCCCTATTAATGAACATTTA
26497 AGGTTTTTTTTTTCAGTTTTTACGCTCTACAAACAGTACACAATAAACACATGACATTAAATACTTGTGCTCTTATTTCACTAGGAGAAAT
26589 CCCCATGTGGAATTTTAAAGTCAAGTTTATTTGTGTTTTTAAATGCTTTAAACATTGCCAGGTACCGTCCCAAAAGGCTATAACAATTCAC
26681 ATTTCTGTTTCTCTGCATCTTACCAGACGAGTGAAATGGTATTTTACTGTGCTTTTCAATTTATATTTTGTGGTTATTTAGTATATTTT
26773 CATATTTTCAATATATTTATTTGCCATTGTGTTTTTCTTCTGACTTGCTTTGTCACATTGTTTACCTTGTCTTCTCTCTGTGTAGT
26865 GTAATAGTTTAGACTCTGAAGCCAGGCAACCTGAGTTAGAAGCCAGGCTCTATTTCATGATGTAGGTCTTTGGGCAAGTACCTAACATTC
26957 ATGCCCTAGTGTCTTCTTTTAAATGAGCAGGATAAATAAGTACCTGCCCTCCTAAGGTTGTATAAAATTAATGGGCACTTAGGGTAAT
27049 ATCTAGCAGGTAGATATTTGGCTATTATCAATAGTACTCTTATCGTTACTATTTCTCCAGATACTGTTTCTGACTCTGGGGCAAGTCTCTG
27141 CTACCCCTGAACCACATTTTCTACCTCTTAGATTACTTGGTAATTCATCAGCCACTGTGGGCACTCCTCTGTGTTAATGCATCATCT
27233 TAGACCTTAGGAGGGATGGGAGGAACCTTAAAGAGCCGAATTTGCTTTTTATTTATCTGTAGCAGAGCAATAGATGTATATTAGGTAGATT
27325 ACAAGCTTTTAGGTTATTTTGCATCTAAAGCTGTCCCTTCTTTTCCAAATAAATGATGTCTGTGGTAAAGAAATATACTGTGGGTGTTAGT
27417 GACAAATCAGAATGCTTTGTGTCTATTTGGCTAGTAGTTAATGTTTTCTTTTATTGTGTCTGCATTCTTATTGTTCTTTAATATATAC
27509 CGAGCTCATTAGCAGTTATCTTGTCTTTATTCATTTCTTATCTCTTAGCATAGTCAGCTCAAGACAACAAGCATCTTTCAGAAAGCCACTAG

FIG. 12B
6 of 10

27601 GAATTGCATCTACATTAAAGAACCCACTCCTCTGCTCTAGCGTCTGAGAACATAACACAGTATTTGCCCTTTGTTGAAGGGCTCAAGCGGACGA
27693 TTCAGAACTAGCAATAGATAACTGTTGGTGGTGGCTTGCAGCACCTAGTACTGATGGTTTGTAGGAAACACACAGGAGTATACAATGTA
27785 AAGCAAGTTCTTTGGGTTTCAAGTAAATAGATTCCACTTGCACGTGTTCTCACTCTTTTGGTGTGAAAATTAGGAAAGGTGATAGGCAGGAT
27877 AGAATAAAATGAAGTGGTCCCCTCTCCATATGGAGAGCGCTGCCCTCCACCACAGACACATGTTTGGCCCTGGAAGCATAAACAGAAGATT
27969 GCAGGAACGCCCTTCACTCCATAGCCTTCAGGCTCCCATCGATAGCATCAAGATAAACTTGGTGTGGCAACAGACTTGAGCCATCATCTT
28061 GTTTAACATTTTACCTGGAAGTGAATGGAATCCAGAGAGGCTAAGTAGCTTGCACAGCTACTTAAATGAAGTACAGACCAAGACCCAGTT
28153 TCTTAGTATCTGGTGGCCATTTTATATAAACATAGACAGCTGATCATGGTAGTTTGGATCATTGCTAAAGACCTATTATATACAACATCG
28245 TATGGCTAAGAAGATTAAGATGCTTATTTCTCGTTATCCATGATTAGAAATGTAATAAGAGACTTTGTTCTGTCAACCAACAGAAGAGACAA
28337 GACGCCCTTTCTGCTGTGGTTGTGAGTGGCCCATCAAGTCTGAGATGTGACTCATGTCTTTGGGAAAGTAGCAGATGAGAGAGTACATTC
28429 TATTCGACTCCATAGACTTAACTGTGAGAGCTGAGAGTAGTGGAGAGCTCAGACTGACAAGAAGAACAGAAATAGACTTAGAGGCACCCAG
28521 GCATGAAAAATCATAAAGATGGAGATGACTTCTATCTTGTGAATGTCTGAAAGGCTCTCTCTCTTTCTGTTCCCAATCTTTCATCTCCAG
28613 CCCTCATCTCTGCCACCATCTGTGGTTTCTTCCGTGGCTCTCTTCTGTCCACCTCTTCAATGTTGGGCCCCAGGATTTGTCTTCAGA
28705 CTCACCTTTCACCCCTCTCCCTCAGGACATCTCATCTCCCAAGGCTGGAAGTCCAGCCCTTCTCTCAGCCTTGGGCTTAAGTTTCATCCCT
28797 TTAGCTGTCTATTGGATGTTTCCAGTTGGAAATCCACAGGCATCTCTACCCAGGCGTCTCTACACAAGATGATAAAAATATATCATCTT
28889 TTTGGCAGATTTGTTTTCTGAGTTTCTGTGCTGTTCATGGATTCAACATTCAAGAGGCTGCCAAGCTAGAAATGTGGGATTTGTAAAT
28981 CGGCCCTGTGTGAGACTGGGAGGCTGGTGTGATAACCAAGAAAGACATTGTGGGCTCTGGGCTTATCCCTTTTAAACCAATTGCCCTCAG
29073 COCTGCCCCGAGTGATGCTTCTGAAATGTGGAAGTCAITTTATCTCTTACCTAAAAGGTTGCAGGATATTTAGATCTGAAATGAAATCCCC
29165 AATCCTTTTTTTTTTTTTTAAAGTGAAGCAAGTTTATTAAGAAAGTAAAGAAATGAGAAATGGCTATGCCATTTGGCAAAGCAGCCCTGTGG
29257 GCTGCTGGTTGCCCATTTTATGTTTTTCTGTGATGATGCTAACAAGGGGTGGGAAATCCCCAATCCTTGGCATTCAAATCCCAGACT
29349 TATTCTTTTTCACMTTTTTTTTTTAAATCATGCCCTGCCCTTTCAGTTATGATGAGTACTTGGTCAITGCTCAGGTGTGACTTGTCCCTTTGC
29441 ACCTGCTGTGCCCTCTCTCTAGAATGCCGTTTTCTCCTTTCCTGGCCAAAGTGTCTTGTCTTCCAAATGGGCCCTTCCCTTGGGAGGTGGTTT
29533 CTGACGACCACCCCTAGTCCAAGTCAGCTCCCACTGTACTTTAACTTTCTCTTGTCTTCTTATTAACCTGTGTATGCTCTCTCCCCAC
29625 CTGGTGTCTCTTGGGACTAGGGACTTCCCTTCAATCACATTTACATAAAGTGTAGGGCTGGCTCATAAGAGGTGCTTAATGAATATTATATG
29717 AATTAATAGCATCTGTCTTCAAGATCAGCCATCAITTTCTCTATCTCATCAATCAAATATATCTCTCTCTTCCCTTCTTGCACCC
29809 AGTCACAGACTGGACTCTATTAATCCTGTCTATCATCTGGGCTCAITTTCCATCCTCAGTGTCTGTCTGTGCATCCTTTTCATTACGCCAGG
29901 GATGTTCAAGTCTGATCTGCCCCCTCAITTCAGGCTGTCCCATATTCATTACTTTATGAAGGCTTTCTGTACACACAGATGCTTAATAT
29993 TCTCTTTTGCITTTCTTGTGTGACTTTGACTCTGCCACTGGTGTGAGCTTCAGAAGGGCAGGGATCTCACTTCACTTCTTTTTCTCCT
30085 AGTGCTTTCTTGTGTGCTGCACACTCCCTGGCACACACAGCGGCTCTCCAACAGGAGGACAGGCTTCCAGCAGCCTCAACCTTCAGGACT
30177 GGGCAGCTTTTAAATGATTTTGGGCACCTTTGCAAGAAAGGATTTGTGTTAAATGTAATTTTCATGATCTTGTATCAGAGTGCAAGAGAA
30269 GCAGAAATCAAACTGAAAGTGAAGCAGAGTCAAATTTTTATTTGAGCCAGATGCCAGGTAAAGAACTATTAATGTGTTTAAATATTA
30361 AAACCAATGTGGGAGAGAAATCATCGATGGGCTTATTTGTTTATTTGTTTATTTTGGAAAAACAAGCAAAATAACTATAGA
30453 AAATTTGGGAAAAAGAGGAAAAATAAAATGATATCTTATCACCAATAGCATTACTATTGTGAATATTTGATATCAATAGATGTTTGA
30545 AITGGGAGAGATTTATGAAAGACATCTCAAGTTCACAAAGAACATCAATTTACCTGTAAATAAACCATCAGAAACAACAGGTATCAC
30637 TGCAGTTGCCCTGGGAGTCAGTGATAATCCCGACTAGCCAGGCTCAGGCTCAAATACAAACCTTTTCCATTAACTCTAACGATAAGTACT
30729 TTTCTGTTTCTCACAAACCTCATAACCATACGTATGTGTGTTTATATGCTATATTTTTTATTTGCTTTTAAAGAGTTTTGTGTTATCAIT
30821 GTAAATATACATAAATAAATTTACCAITTTAACCAITTTTAAAGTGACGGTTCAGTGGCATTAAATACATCTCATTGTTGTACAAACCA
30913 TTACCACCATCCATTTCCAGAACTTCTTCAITTTCCACACGGAACTTTGTATCAATGATAACCTTCTTCTCTCTCCCCCATCCCCCT
31005 AGTAACCTCTGTCTACTCTGTGAACCTGCCTATTTTAGGAACCTCATAAATGTGAATCATAAGATTTGTCTTGTGTTCTGGCTTCTT
31097 AAACCTAACATGTTTCAAGGTCATCCATGTTGTAGCATGTGTCAGAAATTTCTTCTTCTGTGGCTGAATATCCATTGTATGTATATA
31189 CTACATTTTATATATCCTTGTAAATCTGTGTAGGACACTTGGTTGGATACTTGATGGACATTGGTTTTGTTGTTTCAATGATCATAATTTTCA
31281 AGCTCTGTATTTTTTTCAGTTTCACTTCAITGAGTATACCATCATGTCTTTTTTTTTTGTCTTTTTTTTTTTTTTTTTTTTGGGCAGAG
31373 TCTTGTCTGTGCGCCAGGCTGGAGTGCAGTGGTGAATCTCGGCTCACTGCAAGCTCCGCCCTCTGGGTTACACCAITCTCCTGCCCTCAG
31465 CCTCAGCCTCCCGAGTAGCTGGGACACAGGTGCCCACTACCACACCTGGCTAAITTTTTTGTATTTTTTGTAGAGACGGGGTCTCACTGGG
31557 TTAGCCAGGATGGTCTCGATCTCCTGACCTGGTGAAGCCGAGCCTCGGCTCCCAAAGTGTGGAATACAGGCGTGAGCCAGCCTGCCCG
31649 GCCCATGTCTTTGACCAITGTTATAAACTATGTGTGAACACTATAAACCATAGAAACCGATTATATAATAGCAACACTATTGTGAGTAAA
31741 TAAGTGTATATAGCTTTTCCATATTTTATCCGTTTCCCTTGGATGCAITTTATGATTTTTTTTAAATTAAGAGCAATACTTATTTATATGT
31833 TTACATTTCTTTTAAAGAGCTTGACTTCTCATGCTGAGCTGAGCCAGAGTGAAGTCAATTTGAAGAGAGTTTGGAAAAAGGATCTTGTCAA
31925 GTGCAATGATTTATCTTTCAATTTGCAATGCTGTGTTGCCGAAATGAAGAGGACCCACTACAATGTAATTTTCAITTTTAAAAATAAAC
32017 ATTAATAAAAAAATAGGCAGAGGTTTCAAGATGACCTTTACAGTGCAGCCTGGATAAGAAATCTAGTCCCTGGTATCAAGAGGTGCAGTG
32109 TTTGATCAGGATATGGAGGTTGTAGCCTGCAAGGACAGGATGTTCTGTATGGAAGATGAGGGTGCAGGTTTGTCTCAGCTTTCCAGGA

FIG. 12B

7 f10

FIG. 12B
8 of 10

36801 ATACCTTTTAAATCGATATAGATAGGTGCATGGATGGGTCAATAGGCCATTCTGTTTGTGTTTCAGAGACAAAGAGGATTGAATGTGTAA
36893 AACTGAGAAATACATAAGCCAGATTTTGA AAAAATCATTGGTAGAGTACAGAGAGGATAGACACTGTCTGGAGAAGTGCTACTGGAAC
36985 TGGCAGGGTGCACGGTAGTGTAGCTGCAGAGCTGGGATTCAASSACCCACACATGCCTCCAGCTGGAAAGTCAGGGCAATCCAGTGAGGC
37077 CTGGGGTGATCTTTATCTCTTGA CTCTACTTGTAAAGCATTGTGCTGTATATGTGTTTCCTAAGCACAAAGCCATTGGCTGGAACTGTTTT
37169 CTATGTAAATGTATTTAGTTGTCTCATCCCCATAGATGTTTTCCATGTTTTTAGATAATGAGATTTCTGTTGGCTATAGCCATGGAATAA
37261 TAATTAGACTTCTCATAGAACTAGACTTTAAATAATGAATTGATTTTGGTGTGTTTGGAAACCCAGTTTAGAAATGTGCTTTGCCATTTTCAG
37353 GACATGTTTAAAGGATGCATCTGGAAATCTTGACAAAATGCCAGATTTTCTGCCATCTACAGGCAAGACAGCAATAAGCTATCCAATGATG
37445 ACATGCTCAAGTTACTTGCAGACTTTGGGAAGTGAGTTTCAAGGTCTTATTTCCACACCTGAAAAATAGAAGCTGTGTAGTGGGGAGGGAGG
37537 AACAGGGGAGCAGTCACCTAGGTTGCTCGATTAGACATCAGAGGGGATGGCAATGAGCGTGAAGCATTTCCTCAAACCCCTTGAGAAGAAA
37629 GATGGGGTGAAAAATCAGAAGAATAACCAAGTTAAATTTGAATCTGTAGAGGATGTTTTGGGTGGTGTGTGAAGGGTGGACTGGGTAAAGGATG
37721 AGCCTATGGTGGGGAGGAAACAGTTCAGGAACCTTGTCAAGAGGTGAGAAAGGACTCAGCAAAGCCACTGCAAGTGCAACAGGAAGAAGGG
37813 GACAAATTCAGTCTGTGCCAAAGAGATACGATGACTGGGTCTTGGCTTGGGGTGGTAATAGTCTAAGATAAATAACTTGCAAGGTTTCTAAC
37905 TTGGAGTTTCTGGCACCAGTTGTTGCTTAGCTTGTGGCAGCATGCTCCACTCTGCCTCTGGCCTCACATGGTCCCTTCTCTGTGTCTC
37997 TGTCTAAATTCCTTCTTAGAACCTAGTGATAAAGCATCAGGGCCAGGCTAGTGACCTCATCTTAAGTATTGATGTCATGTCAGAGACCCCTG
38089 TTTCCAAATAGGTCACATTTATAGGTACAAAGGGTTGGGACTTCACCATGCTTTTGGAGGACACAATCAACCCATAAATGAGGCAAGA
38181 GGGAGCAAGGAATGTGTGCAACATCAGAGGCCGGCAGCTTCCCAAGTCAGTCTCACCCGAGGGTCTGTGTTCTTAACCTCTATGCTGTTT
38273 TGCTGCTACATCTTAAGAGTTCACTCTGAACCTTTGAATCTGATTTTCTTGTGAGGAGATGGGTCTTAGAATTTTCTGGGGAATTCCT
38365 GCGAATGTGAAGAGCTGAGGGCGCTAGAAGATGTGAAGTGAAGAGAAATAGCTGAGAGCCAAATGCTAAGTATTCTATGCCAAAGGTATCCT
38457 TGTTTTTTTTTTTTTTGTGTCATATCAAATAGCAATCTTATCAGTTTGTCTAGAACTCAAGAATGATGCTTAGCTTTCTTTAACCTTATT
38549 TTACCTTTTTCTTATCTGCTTCAGTAGTAGGAATAGAAACGATATGAGTCATAGAAACAGGCTCAATAAGTTCTGAAAACACAGAGACGTG
38641 TTCTTAATCAGAAATCCAATCAGTCCATGTCAGCAGGGCGGCTTCAGCCTTCACAGCGACGTGAATCCCTTGTCAAGAGGCTCAAAAAGGTA
38733 GAAAGGATTCTCAAGGTCTCTTTCA GTTATGTGATTATACAGTTTTTGA CTGTCTTGATGTTTCCCTGTGTTGGAGCTTTAATGAGAAGTGC
38825 AACCTCAGTTTTTGCTAACATGCAGCTAAGGTTGGCTGTTCAGCAAAGCAGTGTGCATGCCCGCTGGGCTGATTGGAATGAACCTTTCACA
38917 GCTCACGTAGGGAATTGGAGAAGGGGAGAGGAGGATACTGGTGAAGGATGAGGCCCTGCTGGGTAGCCCTCCAGGGTTCCTGGACCATATA
39009 GGTGCCCCAAATTCAGTCACTATCTGACAGTTTATGACCTGGTAAGGACACAGGCTTTGGCCAGGGAGTGCCCTCGATCCCTATGAAT
39101 CTGTTATTCATGAAAGACTAAATAAAGAATAGTACCCTATTTTACTTTTAAATCATAGAGGTTCTTTAGTTTACVACATAATACATGTT
39193 CATTTTAGAAATTTTGAGAAATACAGAAGAATAAAGGATGAAAAAGGTTTACTACTAGTTTAAACCTTCGTGGTGAACTTTGGAGAAGT
39285 TTTTTTTTTTTTTTGAGATGGAGTCTCACTCTGTCCCTCAGGCTGGCGTACAGTAGCACGATTTCAAGTCACTGCAACCTCCGCTCCGAG
39377 TTCAAGCGATTCTCTGCCCTCAGCCTCCCAAGTAGCTGGGACTAGGGCCACCACCTACGCCCTGGCTAATTTTGTATTTTATAGTAGAGA
39469 TGGGGTTTCAACATATTGGCCAGGCTGATCTCAAAGTCTGACCTTGTGATCTGCCCGCTCAGCCTCCCAAAGTGTGGGATTACAGGCAT
39561 GAGCCACCGTGGCCAGCTAGGGGGGAACATTTTTTTTACGTTTATTCCTTTACATTTTATTTAGTTTATCTTATGTAGCTATGATCAT
39653 ACTAAATATGTAAATATTTCCTGACAACTCAAGTATTTTCTGAAAGTGTATATATACATTTTATAGACATCATTTTATATGCATAAATA
39745 TTATAATAGTCCATTGAGATAGACATAGATTATTTAACTCTTCCCCCATTTTTTGACTTTTTTTTTTTTTTCCGAGATGGAGTCTCGCTCT
39837 GTCGCCAGGCTGGAGTGCAGTGGCACCATCTCAGCTCACTGCAACCTCGCCCTCCAGGTTCAAGCAATCTCTCGCTCAGCTCTCTGAGT
39929 AGCTGGGACTCCTGAGTAGCTGAGTAGCGCATGCTGCCACGCCCGCTAATTTTTTTTTTTTTTTTTTTTTTTTGTATTTAGTAGAGA
40021 TACTAAATATCTCACCATCTTGCC CAGGCTGGTCTCAAAGTCTGACCTCAGGCAATCTGCCCGCTCAGCTTCCCAAAGTGCTGGGATTAC
40113 AGGTGTGAGCCACCATGCC CAGCCATTTTTGACTTTTAAATGTGTTTCTGATTTTTCAGAAATATACCTATAAGCCACAGTAGAATCTTTA
40205 AAAAAATCTCTCTATTGGTAGTGGTAATATATTATCATACATACTATATATCATATAGTAATATTGTCAATTTTTGAGTTTCAAGAAA
40297 ATTTCACTCTTACTAATTTTTTCAAAAACAGTCACTTTAGTTGATAGATTTCAATATTTCTCTCGCTCAACTACCATGCAACTCTTAA
40389 TAACCATGAGGTGGGTCTGCGTGACTTAGGAAAGTGAATACATATATTATTAAGGAAGAAAAATATATCTGTATTACTATATTTTTGA
40481 AAGAAATATATATTTCTTTTGTATGTAATGAAGAAATGGATAAGCAAGTAGCTATCTAGATGGAAAGATAGGCAATAAATAGCTATTTA
40573 GGATATATGCCAAATAATCATGGTTATCTCTGAGGGATGGGTGTAGTGGGTGATATTTCACTTCCACTTTATAACATTCTGTCAITTTTATA
40665 TGAACTTTTAAAAAATAACACTTTTATATTCAGACAAAACAAACATGAAGTTTTTATATGTGATGGAGTTGGAGCCCTGTCTCAGAAG
40757 TTACTTCTTAGGCTGGTTAGCTTGAGACTTCCCCACAGTGGCGGCTCCTCAGGGGCAGGCCAGTGCATGGTCTGTCTTTCAGTGGAGGCTG
40849 GGGAGTGGGGCTTCACATGGTCACTAATTTGAAAGTGTGAGTGCATGAGGAGGAGGCAATGGTGTGAAGGGTGTGCTCCAGGTGGCCAGTTAGAGACCC
40941 GAGTCTCTCTGACACCACTTCTTCATGCTGTGCTCAGCAGGAGGGCAATGGTGTGAAGGGTGTGCTCCAGGTGGCCAGTTAGAGACCC
41033 AGAAACCTGAAAAACAGGGATCCGATGGTGACAGCATAGAAGACACAGCAGGATAAGTGAAGGCCAGCTCCTCAATAAGTATTCAAGAAACT
41125 TTGGTGCCCACTCCCGTATTCTTCACAACAGAGTTAGGGGAGCTGGAGGATTCCTTTTTTCAATTTTTTAAAAATCTTTGCAATGCTATTTTT
41217 CTTTCTCTGTATATTTTACAGGAAATAATCTCATGTGCTAGTGGCTGGGCACCGGCTTGGATCCAAAGCTATTGTTTCTACCCCATGATG
41309 TCTCAAAATGTTATTTAATAATGCATGAAAAAATTTCTTCAGCTGTCTCAGTCTTAACAAAACAGCTGCCAAAGCTCATAGCCACTTTC

FIG. 12B
9 of 10

41401 CTTTTTCCCTTGCAATAATTACCCAGGGATATGTTCCAAGATTAGTAAGAAAGCGATTCTGTCCGATAGATGATATTGCTAACATTTTATA
41493 AGAAGCAGAGACTTGGTACTTTGTATTTGATTTGTTTCATGGTGGTATCTCATGGATAAGATGGTATCTCATCTTTTCCAACCTCTGCAGGAAA
41585 TGCGAAGACATGAAGGCAAAGTATAAAAATAGAACGTTTTCTTTAAACGTAGACCTTTTAAATGGTACTACGTTGGATAGTTTAGGTAATA
41677 ATACTACTAAAGTTTTTCCGTATGCAGCTTAATGTGTCTGTGTTATTTGTACACTCATCTTCTTTGCATCCAGGTTTTACAGTCTTACCCC
41769 GATTTGCTCTGGTTACACTGCACCTCAAGCCAAGTAGGGCTGCTTGACTTTCTCTAAACCCACTGGGGACTTCCCTCTGCCATGCTTTTCTC
41861 TCTGCCCAAATGTGTCCCCCTTCTGCTCATCAAGCAGCACATAAATCACAAACACATGCAGCATACACACTTCCCTTTTCCCTTTGTCTT
41953 TCTCAGGGAACCTCTACTCATCTTTCAAAGCCAGTCTGTGGCTCACTTCTGTGCTGGGAGTCTTGGAGGCGGTTACTTGGCTTCTCTGCCCTG
42045 AGCCCGCTCCTCTTTTAAGGGTGGATAATAACAGCCCTGCCOCCATAAACCCGTGGTGGGGAATAAATGCPAAAGGCATTAGGTGATTC
42137 TCCCACCATGAATACTGATCTCATCCCGTGTTCCTCTCGATAGATCTAGATACTCTGCCTTCTGGTAGAGGTTTGTACATACTCTGTGAAA
42229 GTGATTGCCCTCATATGCCGTAAGTAGCTTACAGTGTCTACTGGACTTTTGGCTTCTTGAGGAAAGAAATTATGTCTTGTGCAITCCTCC
42321 ATGGTCTTGAGTACATACATTGCAGCATATCTTAAGCACCTTGATAAATGCTTAITGAATTTTCTTCTAGACATAAAGTCAAGTGGTTTTGT
42413 TGAAACAAAAATATCTCAAATTTCTTTCAATCATATATAGTTGTTTTTTTTTAAGTGACACCAGCTTTTAGGGAATATTTCTTTTACAA
42505 AACACAGTTAGAAGATTAACTCACCACCAATAGCAGTCCAAACATACCTGTATTGCCAGCTAATCATTTTAAAGGACCAATACAGGAAGTC
42597 AGGAAGGGAAGACCGGCTGCAGAAACCTTAGATAAGGACCCCAAATCTGTTGGCATGGGAGGACTGCTAGTTGATGATACCATTCCTCATTT
42689 CCTCTGTGGGAATTGTTGAGTCAGCAAAAATGGATGGGAGTGGGAGGGAATAATTTCTAAGAGAGAGTTTGAGCCTCACTTCTACATTC
42781 ACACAGAGACAGGAGCAGTTCCAGAGGCCAGGCATCCTGCAAGTGTCTGTATTGCATGCTTACTTAATTCGTGTAATTTTAAAGTAGATTT
42873 TCAATGTTCAAGGATTATTTTATAAATTTTGCATAGAATATAGGTACTCTTTAGCAAAACAAAGCAAAAAACCAAACTATTTCTAGTCATG
42965 AAAGAATTCAAGTTTGTGTACACGGCACACAACCACCACTTTGGAAGTGCATAAAAAGGCAGTAAATCTTTATTTGCTGTGAGTGTGTTGATG
43057 TCTAATAAACCCAGATTCAACATAAACCATAACTTTTGAATGGGTTTGAAGATTGGGTTTTTAAAAACTTAAAGCTGGCAAAAAAACA
43149 ACTTTTAAAGCCCATGTGCTACATAATATGGAACTAACCTCAGAAATGTGCTTGGAAACACATGGAAAGAACGCTTTTACAGAGCAGCAA
43241 CTAGAAGTAAATCTCTCAGCAGAGGGAGGAAATAGAAATAAGAAATAACTATAGTTAGGCACAGAGGACACAATACTATAGGAAGATTT
43333 CCAGTGAAGATCATTTAATTAAATATATGTTGCTTAGAAACGATTTTAATTTGTGTTCCACCTCTCTCAAAAATTTATATGTGGAGGATGTTG
43425 GAGTGATCTTAAAAATGGTGATGAAGATGCCTGTTCAATCATAGGTGGAAATAATTAGGAGGGGTGAATCCATTACCTTGCATACTTAC
43517 TTATATTTAAAGTATAAATTTGTAATAAA

```

hCLASP4 -----MFPMEDISISVIGRQRRTVQ----- 20
hCLASP5 -----MTHLSLDVQLAQELG----- 16
hCLASP3 -----MAERRAFAQKISRTVAAEVRKQISGQYSGSPQLLNINIVG 41
hCLASP2 -----MLLFYDDFQTAILRRQGRYICS----- 23
hCLASP7 -----MAASERRAFAHKINRTVAAEVRKQVSRERSGSPHSSRRCSSSL 43
hCLASP1 MSFRGKVKFKREPSEFWKKRRRTVRRVIEEHRFSSQEKPRLLEPLDYETVIEELEKTYRN 60
      ...

hCLASP4 -----STVPEDA EKRAQSLFVKECIKTYSTDWHVVNYK 53
hCLASP5 -----DFT 19
hCLASP3 N-----ISHHTTVPLTEAVDPVDLEDYLITHPLAVDSGPLRDLIEFP 83
hCLASP2 -----TVPKAEAAAQSLFVTECIKTYNSDWHLVNYK 55
hCLASP7 G-----VPLTEVVEPLDFEDVLLSRPPDAEPGPLRDLVEFP 79
hCLASP1 DPLQDLLFFPSDDFSAATVSWDIRTLYSTVPEDA EHKAEENLLVKEACKFYSSQWHVVNYK 120
      ::

hCLASP4 YEDFSGDFRMLPCKSLRPEKIPNHVFEIDEDCEKDED-----SSSLCSQKGGVIKQG 105
hCLASP5 DDDL DVFTPKCRTLP--SLPEEGVELDPHVR-----DCVQTYIREWLI 63
hCLASP3 PDDIEVVYSPRDCRTLVS-AVPEE-SEMDPHVR-----DCIRSYTEDWAI 126
hCLASP2 YEDYSGEFRQLPNKVVKLDKLPVHVVEVDDEVDKDED-----AASLGSOKGGITKHG 107
hCLASP7 ADDLELLLPRECRTTEP-GIPKD-EKLDAAQVR-----AAVEMYIEDWVI 122
hCLASP1 YEQYSGDIRQLPRAEYKPEKLPSHSFEIDHEDADKDEDTTSHSSSKGGGGAGGTGVFKSG 180
      :: .      : * . : : * .

hCLASP4 WLHKANVNSTIT--VTMKVFKRRYFYLTQLPDGSIILNSYKDEKNSKESK-GCIYLDACI 162
hCLASP5 VNRKNQGSPEIC--GFKKTGSRKDFHKT-LPKQTFESETLECSEPAQA--GPRHLNVLC 118
hCLASP3 VIRKYHKLGTGF--NPNTLDKQKERQKG-LPKQVFESDEAPDGSYQDDQDDLKRMSI 183
hCLASP2 WLYKGNMNSAIS--VTMRSFKRRFFHLIQLGDGSYNLNFYKDEKISKEPK-GSIFLDSM 164
hCLASP7 VHRRYOYLSAAY--SPVTTDTQRERQKG-LPRQVFEQDASGDERSGPEDSNDSTRGSGSP 179
hCLASP1 WLYKGNFNSTVNNTVTVRSFKKRYFOLTQLPDNSYIMNFYKDEKISKEPK-GCIFLDSCT 239
      : :      : : : * : : : : :

hCLASP4 DVVQCPKMRRHAFELKMLDKYSHYLAETE QEEMEENLITLKKIIQINTDSLVEKKETVE 222
hCLASP5 DVSGKG PVTACDFDLRSLQPDKRLNLLQQVSAEDFEKQNEEARTN-----RQAE 169
hCLASP3 DDTPRGSWACSI FDLKNSLPDALLPNLLDRTPNEEIDRQNDQQRKSN-----RHKE 234
hCLASP2 GVVQNNKVRRAFELKMQDKSSYLLAADSEVEMEENITILNKILQLN-----FEAAMQEK 219
hCLASP7 EDTPRSSGASSIFDLRLNLAADSLPSSLERAAPEDVDRRNETLRRQH-----RPPA 230
hCLASP1 GVVQNNRLRKYAFELKMNLDLTYFVLAATESMDENIHTLNRILQISPEGFLQGRSTEL 299
      * : * :      : : : : :

hCLASP4 TAQDDETS-----QGKAENIMASLERSMHPELMKYGRETEQLNKL SRGDGRONLFSFDSE 278
hCLASP5 LFALYPSVD-----EEDAVEIRPVPECPEHLG-----N-----RILVKLLTLKFEIE 212
hCLASP3 LFALHPSPD-----EEPIERLSVPDIPKEHFG-----QRLVKCLSLKFEIE 277
hCLASP2 RNGDSHEDD-----EQSKLEGSGGLDSYLP ELAKSAREAEIK---LKSESRVKLFYLDPD 272
hCLASP7 LLTLYPAPD-----EDEAVERCSRPEPPREHFG-----QRILVKCLSLKFEIE 273
hCLASP1 TDLGLDSLDSVTCECTPEETDSSENNHLHADFAKYLTETEDTVKTRNMERLNLFSLDPD 359
      :      : : : : : : : : : :

hCLASP4 VQRLDFS-----GIEPDIKP-FEECKNKRFLVNCHDLTENILGQIGDNAKGPPTNVEPFFI 333
hCLASP5 IEPLFAS-----IALYDVKERKKISENFHCDLNSDQFKGFLRAHTPSVAASSQARS AVFSV 268
hCLASP3 IEPIFAS-----LALYDVKEKKKISENFYFDLNSEQMKGLLRPHVPPAAITTLARSAIFSI 333
hCLASP2 AQKLDIFS-----SAEPEVKS-FEEKFGKRI LVKCNLDLSFNLOCCVAENEEGPTTNVEPFFV 327
hCLASP7 IEPIFGI-----LALYDVREKKKISENFYFDLNSDSMKGLLRAGHGTHTPAISTLARSAIFSV 329
hCLASP1 IDTLKLQKKD LLEPESVIKPFEEKAAKRIMII CKALNSNLQGCVTENENDPITNIEPFFV 419
      : :      : : : : : : : : : : :

```

FIG. 13
1 of 6

3NSDOCID: <WO 0231117A2 I >

```

hCLASP4      FKSHLESTIYTQDLHVHKFFHHCQLIQS-----GSKEVPGELIKYLKCLHAM 794
hCLASP5      IEVQAVSSVHTQDNHLEKFFTLCHSLESQVTFFPIRVLDQKISEMALEHELKLSIICLNSS 715
hCLASP3      VEVVAVSSIHQDPYLDKFFALVNALDEH-LFPVRIGDMRIMENNLENELKSSISALNSS 780
hCLASP2      ISTHLVSTVYTQDQHLHNFQYCKTES-----GAQALGNELVKYLSLHAM 787
hCLASP7      VELTAVSSVHPQDPYLDKFFTLVHVLEEG-AFPFRLKDTVLSEGNVEQELRASLAALRLA 767
hCLASP1      VSTFVVSTVNTQDPHVNAFFQECQKREKD-----MSQSPTS NFIRSCNLLNVE 887
..      *:: : ** : : * : :

hCLASP4      EIQVMIQFLPVILMQLER-----VLTNMTH-----EDDVP 824
hCLASP5      RLEPLVLFLHLVLDKLEQLSVQPMVIAGQTANFSQFAFESVVAIANSLHNSKDLKSDQHG 775
hCLASP3      QLEPVVRFLHLLLDKLELLVIRPPVIAGQIVNLGOASFEAMASIINRLHKNLEGNHQHG 840
hCLASP2      ECHVMIAFLPTILNQLER-----VLT-RAT-----QEEVA 816
hCLASP7      SPEPLVAFSHHVLDKLVRLVIRPPIISGQIVNLGRGAFEAMAHVVSLSVHRSLEAAQDARG 827
hCLASP1      KIHAIMSFLPIILNQLER-----VLVQNE-----EDEIT 916
..      : : * : : * : :

hCLASP4      INCTMV-LLHIVSKCHEEGLDS-----YLRSEFIKYS-----FRPEKP 860
hCLASP5      RNCLLASYVHVFRLEPVQRDPVPSGAPTALLDPRSHTYGRTSAAAVSSKLLQARVMSS 835
hCLASP3      RNSLLASYIHVFRLEPNTYPNSSSPG-PGGLGGSVHYATMARSAVRPASLNLNRSRSLN 899
hCLASP2      VNVTRV-IIHVVAQCHEEGLES-----HLRSYVKYA-----YKAEPY 852
hCLASP7      HCPQLAAYVHYAFRLPGTEPSLPDGAFF----VTVOAATLARGSGRPASLYLARSKSISS 883
hCLASP1      TTVTRV-LPDIVAKCHEEQLDH-----SVQSYIKFV-----FKTRAC 952
..      : : : :

hCLASP4      SAPQAQLIH-----ETLATTMIAILKQS----- 883
hCLASP5      SNPDLAGTHSAADEEVKNIMSSKIADRNC SRMSYYCSGSSDAPSSPA----- 882
hCLASP3      SNPDISGTPPTSPDDEVRSIIIGSKGLDRSNSWNTGGPKAAPWGSNPSPSAESTQAMDRSC 959
hCLASP2      VASEYKTVH-----EELTKSMTTILKPS----- 875
hCLASP7      SNPD LAVAPGSVDDEVSRILASKLLHEELA-LQ----- 915
hCLASP1      KE---RPVH-----EDLAKNVTGLKSN----- 972
..      : :

hCLASP4      -----ADFLSINKLLKYS-----WFFFEIIAKSM 907
hCLASP5      -----APRPASKKHFEELALQ-----MVVSTGMVKSM 910
hCLASP3      NRMSSHTETSSFLQTLTGRLPTKKLFHEELALQWVCSGVSRESALQQA WFFFEIIMVKSM 1019
hCLASP2      -----ADFLTSNKLLRYS-----WFFFDVLIKSM 899
hCLASP7      -----WVSSSAVREAILQHA-----WFFFQLMVKSM 942
hCLASP1      -----DSPTVKHVLKHS-----WFFFAIILKSM 995
..      : : ***

Cadherin Cleavage
hCLASP4      ATYLLEENKIKLERGQRFPEYHHVLSLLLAIIPHVTIRYAEIPDE---SRNVNYSLAS 964
hCLASP5      AQHVHNMdkRDSERRTRFSDFMDDITTIVNVVTSEIAALLVKPQKENEQAEKMNISLAF 970
hCLASP3      VHHLYFNDKLEAERKSRFFERFMDDIAALVSTIASDIVSRFQKDTM---VERLNTSLAF 1076
hCLASP2      AQHLIENSKVKLLRNQRFPPASYHHAETVVNMLMPHITQKEGDNPEA---SKNANSLAV 956
hCLASP7      ALHLLLGQRLDTERKLRFPGRFLDDITLVGSGVGLVITRVHKDVEL---AEHLNASLAF 999
hCLASP1      AQHLIDTNKIQLERPORFPESYQNELDNLMVLSDHVIWKYKDALEE---TRRATHSVAR 1052
..      : : * : : * : : : : : : : : * : :

hCLASP4      FLKRCCLTMDRGFIINLINDYISGFSPKDP-----KVLAEYKFEFLQITCNHEHYIPLNL 1019
hCLASP5      FLYDLLSLMDRGFVFNIRHYCSQLSAKLSNL---ETLISMRLFLRILCSHEHYVTLNL 1027
hCLASP3      FLNDLLSVMDRGFVFSLIKSCYQVSSKLYSLPNPSVLVSLRLDFLRIICSHEHYVTLNL 1136
hCLASP2      FIKRCFTFMDRGFVFKQINNYISCFAPGDP-----KTLFEYKFEFLRVVCNHEHYIPLNL 1011
hCLASP7      FLSDLLSLVDRGFVFSLVRAHYQVATRLQSSPNPAALLTLRMEFTRILCSHEHYVTLNL 1059
hCLASP1      FLKRCFTFMDRGCVFNMVNNYISMFSGDL-----KTLQYKFDLQEVCOHEHFIPLCL 1107
* : : : : * : : : : * : : : : * : : : : * : : : :

```

FIG. 13

3 of 6

Cadherin EC motif

hCLASP4 PMAFAKPKLQR-----VQDS--NLEYSLSDEYCKHHFLVGLLRETSI 1060
hCLASP5 FFMNADTAPTSP--CPSISSQNSSSCSSFDQDKIASMFDLTSEYRQQHFLTGLFTELAA 1085
hCLASP3 PCSLLTPPASPSVSSATSQSSGFSTNVQDQKIANMFELSVFRRQHYLAGI VLTELAV 1196
hCLASP2 PMPFGKGRIQR-----YQDL--QLDYSLTDEFNRHFLVGLLREVTG 1052
hCLASP7 PCCPLSPPASPSVSSSTTSQSSTFSSQAPDPKVTSMFELSGPFRQOHFLAGILLTELAL 1119
hCLASP1 PIRSANIPDPLTP-----SES---TQELHASDMPEYSVTNEFCRKHFLIGILLREVTG 1157
. : : : : : * : : *

hCLASP4 ALQDN---YEIRYTAISVIKNLLIKHAFDTRYQHKNQQAIAQLYLPFVGLLLENIRL 1116
hCLASP5 ALDAEGEGISKVORKAVSAIHSLLSSHDLPKVCPEVKVIAALYLPVGLIILDALP-- 1143
hCLASP3 ILDPDAEGLFGLHKKVINMVHNLSSHSDPRYSDPQIKARVAMLYLPLIGIIMETV-- 1254
hCLASP2 ALQEFR---EVRLIAISVLKNLLIKHSFDDRYASRSHQARIATLYLPLFGLLIENVRI 1108
hCLASP7 ALEPEAEGAFILHKKASAVHSLLCGHDTDPYAEATVKARVAELYLPILLSIARDTLP-- 1177
hCLASP1 ALQEDQ---DVRHLALAVLKNLMAKHSFDDRYREPRKQAQIASLYMPLYGMLLDNMPRI 1213
* : : : : * : : * : : *

hCLASP4 AGRDTLYSCA-----AMPN-S---ASRDEFPCGFTSPANRGSLSLTDKDTAYGS 1160
hCLASP5 -----QL-----CDFTVADTRRYRTSGSD----- 1162
hCLASP3 -----QLY-----DFTETHNQRGRPICLATDD-- 1276
hCLASP2 NVRDVSPFPVNAGMTVKDESALPA-VNPLVTPQKGSTLDNSLHKDLGALSGIASPYTT 1167
hCLASP7 -----RLH-----DFAEGPGQRSRLASMLDSOTE 1201
hCLASP1 YLKDLVPFTVNTSNQGSRDDLSLSTNGGFQSQTAKHANSVDTSFSKDVLSIAAFSSIAIS 1273
. : : : : :

hCLASP4 FQ-NGHGIKREDSRGS LIPEGATGFPDQGN TGEN-----TRQSSTRSSVSQYNRLDQYE 1213
hCLASP5 -----EEQEGAGAINQNALAIAGNNFNLT-----SGIVLSSLPYKQYNMLNADT 1208
hCLASP3 -----YESESGSMISQTVAMAIAGTSVPQLTR-----PGSFLLTSTSGRQHTTFSAES 1324
hCLASP2 STPNINSVRNADSRGSLISTDSGNSLPERNSEKSNSLDKHQSSSTLGNSVVRCDKLDQSE 1227
hCLASP7 -----GEGDIAGTINPSVAMAIAGGPLAPGSR-----ASISQGPPTASRAGCALS AES 1249
hCLASP1 -----TVNHADSRASLASLDSNPSTNEKSSEKTDNCEKIPRPLALIGSTLRFDRLDQAE 1327
. : : : : . : : : :

hCLASP4 TRSLLMCYLYIVKMI SEDTLLTYWNKVSPQELINILILEVCLFHFYMGKRNIARVHDA 1273
hCLASP5 TRNLMICFLWIMKNADQSLRKWIADLPSTQLNRILDLLFICVLCFEYKKGQSSDKVSTQ 1268
hCLASP3 SRSLICLLWVLKNADETVLQKWFDTLSVLQNLRLDLLYLCVSCFEYKKGKVFERMNSL 1384
hCLASP2 IKSLLMCFYILKMSDDALFTYWNKASTSELMDFFTISEVCLHQFQYMGKRYIARNQEG 1287
hCLASP7 SRTLLACVLWVLKNTPEALLQRWATDLTLPQLGRLLDLLYLCLAAFEYKKGKAFERINSL 1309
hCLASP1 TRSLLMCFHIMKTISYETLIAYWQRAPEVSDFSI LDVCLQNFYLGKRNIIRKIAA 1387
: : * * : : : : : : : : : * * * : :

hCLASP4 WLSKHFGIDR-----KSQTMPALRNRSVGMQARLQHLSSLESS----- 1311
hCLASP5 VLQKSRDVKAR-----LEEALLRGEARGEMMRRAPGNDRFPGLNEN-- 1311
hCLASP3 TFKKSKDMRAK-----LEEAILGSIGARQEMVRRSRGQLERSPSGSAFGSQ 1430
hCLASP2 LGPIVHDKRS-----QTLFVSRNRTGMHARLQQLGSLDNS----- 1323
hCLASP7 TFKKSLDMKAR-----LEEAILGTIGARQEMVRRSRERSPFGNPEN----- 1350
hCLASP1 AFKFVQSTQNGTLKGSNPSCQTSGLLAQWMHSTSRHEGHKQHRSTLPIIRGKN----- 1442
. : : : : . : : : :

hCLASP4 -----FTLNHSSTTTEADIFHQALLEGNTATEVSLTVLDTISFFTCQFKTQLL 1359
hCLASP5 --LRWKKEQTHWRQANEKLDKTKELDQREALISGNLATEAHLIILDMQENIIQASS-ALD 1368
hCLASP3 ENLRWRKDMTHWRQNTTEKLDKSRAEIEHEALIDGNLATEANLIILDTLEIVQTVS-VTE 1489
hCLASP2 -----LTFNHSYGHSDADVLHQSLLEANIATEVCLTALDTLSLFTLAFKNQLL 1371
hCLASP7 --VRWRKSVTHWKQTS DRVDKTKDEMEHEALVEGNLATEASLVVLDLEIIIVQTVM-LSE 1407
hCLASP1 --ALSNPKLLQMLDNTMTSNSNEIDIVHHVDTEANIATEGCLTILDVLSLFTQTHQRQLQ 1500
. : : : : : : : : : : * * * * *

FIG. 13
4 f6

hCLASP4 MFDGHNPLMVKVFDIHLAFLKNGQSEVSLKHFVAFSLAFRAKISKFPFAAFFKGRVNMCAFCY 1419
hCLASP5 CKDS---LLGGVLRVLVNSLNCDSQSTTYLTHCFATLRALIAKFGDLLFEEVEQCFLDCH 1425
hCLASP3 SKES---ILGGVLKVLHLSMACNQSAVYLQHCFAQRALVSKFPPELLFEEETECCADLCL 1546
hCLASP2 ADHGHFPLMKKVFEDVYLCFLQKHQSETALKNVFTALRSLIYKFPSTFYEGRADMCALCY 1431
hCLASP7 ARE\$---VLGAVLKVVLYSLGSAQSALFLOHGLATQRALVSKFPPELLFEEDTELCDLCL 1464
hCLASP1 QCDCQNSLKRGFDTYMLFFQVNVQSATALKHVFASLRFLVCKFPFSAOFGPADLCGSFCY 1560
: : : : * : : : * : : : * : : : *

hCLASP4 EVLKCCSTKSISSSTRNEASALLYLLMRNNFEYTKRKTFLRTHLQIIIVASQLIADVALSGG 1479
hCLASP5 QVLHHCSSSMVDVTRSQCACATLYLLMR--FSFGATSNFARVKKQVMTSLASLVGRAPDFNE 1483
hCLASP3 RLLRHCCSSIGTIRSHPSALLYLLMR--QNFEIGNNFARVKKQVMTSLASLVGTQNFNE 1604
hCLASP2 EILKCCNSKLSSIRSTEASQLLYFLMRNNFDYTGKKSFVRTHLQVIVSVQLIADVVGIGE 1491
hCLASP7 RLLRHCCGRISTIRTHASALLYLLMR--QNFEIGNNFARVKKQVMTSLASLVGTQNFSE 1522
hCLASP1 EVLKCCNHRASSTQTEASALLYLFMRKNFEFNKQKSIVRSHLQLIKAVSQLIADAG-IGG 1619
.: : : : * : : : * : : : * : : : *

hCLASP4 SRFQESLF:INNfansDRPMKATAFFPAEVKDLTKRIRTVLMATAQMKEHEKDPEMLIDLQ 1539
hCLASP5 EHLRRSLRTILAYSEEDTAMQMPFPTQVEELLCNLSILYDVKMREFQEDPEMLMDLM 1543
hCLASP3 EFLRRSLKTLTYAEEDLELRETTFPDQVQDLVFNLMHLSDTVMMKEHQEDPEMLIDLH 1664
hCLASP2 TRFQCSLSIINNcansDRLIKHTSFSSDVKDLTKRIRTVLMATAQMKEHENDPEMLVDLQ 1551
hCLASP7 FHLRRSLKTLTYAEDMGLRDSTFAEQVQDLMLFNLMHLSDTVMMKEHQEDPEMLIDLH 1582
hCLASP1 SRFQHSLA:TNFANGDKQMKNSNFFAEVKDLTKRIRTVLMATAQMKEHEKDPEMLVDLQ 1679
: : * : : : * : : : * : : : * : : : *

transmembrane

hCLASP4 YSLAKSYASTFELRKTWLD\$MAKIHVKNGLFSEAMCYVHVAALVAEYILRRKK----- 1592
hCLASP5 YFLAKSYQASFDLRLTLWLQNMAGKH\$TKKKQYTEAMCLVHAAALVAEYILSMLEDH----- 1598
hCLASP3 YFLAKGYQTSFE-RLTWLQNMAGKH\$SERSNHAEEAQCLVHSAALVAEYILSMLEDR----- 1718
hCLASP2 YSLAKSYASTFELRKTWLD\$MARIHVKNGLLSEAMCYVHVTLVAEYILTRKG----- 1604
hCLASP7 YFLARGYQGSFDLRLTLWLQNMAGKH\$AELGNHAEAAQCMVHAAALVAEYILALLEDO----- 1637
hCLASP1 YSLANSYASTFELRRTWLESMAKIHARNGLLSEAMCYIHIAALIAEYILRRKGYWKVEKI 1739
: : : : * : : : * : : : * : : : *

hCLASP4 -----LFPNGCSAFKKITPNIDEEGAMKEDAGMMD----- 1622
hCLASP5 -----SYLPVGSVSFQNISSNVLEESVVS\$EDTLSPDEEDGV 1633
hCLASP3 -----KYLEVGCVTFQNISSNVLEESAVS\$DDVVS\$PDEEGI 1753
hCLASP2 -----VFRQGCTAFRVTITPNIDE\$EASMMEDVGMQD----- 1634
hCLASP7 -----RHLVPGCVSFQNISSNVLEESAIS\$DDILSPDEEGF 1672
hCLASP1 CTASLLSEDTHPCDSNSLLTTPSGGSMF\$MGWPAFLSITPNIKEGA\$AKEDSGMHD----- 1795
: : * : : : * : : : * : : : *

ITAM

hCLASP4 ---VHYSEVLLLELLEQCVDGLWKAERYEIISEISKLVPIYEKRRFEKLTQVYRTIHG 1679
hCLASP5 CAGQYFTESGLVGLLEQAAELFSTGGLYETVNEVYKLVIPILEAHREFRKLTLTHSKLQR 1693
hCLASP3 CSGKYFTESGLVGLLEQAAASFSMAGMYEAVNEVYKLVIPILEANRDAKKLSTIHGKLQE 1813
hCLASP2 ---VHFNEVDVLMELLEQCADGLWKAERYELIADYKLIIPYIEKRR----- 1677
hCLASP7 CSGKHFTELGLVGLLEQAGYFTMGGLYEAVNEVYKLVIPILEAHRDYKLLAAVHGKLQE 1732
hCLASP1 ---TPYNENILVEQLYMCGEFLWK\$ERYELIADV\$NKP IIAVFEKQRDFKLLSDLYYDIHR 1852
: : * : : : * : : : * : : : *

ITAM DOCK motif DOCK motif ITAM

hCLASP4 AYT\$KILEVMHTKRL\$LGIFFRVAEYQG\$FFEEELGKEYIYKEEKLTLGLSEISLR\$VKIY\$ 1739
hCLASP5 AFDSIVN\$KDH--KRMFGTYFRVGEFG-SKFGDLLEQE\$VYKEE\$AITKLPEISR\$LEAFY\$ 1750
hCLASP3 AFSKIVH\$QSTGWERMFGTYFRVGEFG-TKFGDLLEQE\$VYKEE\$AITKLAEISR\$LEGEY\$ 1872
hCLASP2 -----DFFEDELGKEYIYKEEKLTLGLSEISOR\$LLKIY\$ 1710
hCLASP7 AFTKIMH\$QSSGWERVFGTYFRVGEFG-AHFGDLLEQE\$VYKEE\$ITKLAEISR\$LEEFY\$ 1791
hCLASP1 SYLKVAE\$VNSEKRLFGRYRVAEYQG\$FFEEELGKEYIYKEEKLTLGLSEISOR\$LLKIYA 1912
: : : : * : : : * : : : *


FIG. 13
5 of 6

	ITAM	ITAM	
hCLASP4	ERFGTENVKIIQDSQKVNAKELDPK	YAHIOVTYVKE	YFDDKELTERKTEFERHNHISRFV 1799
hCLASP5	QCFGAEFVEVIKIDSTPVDKTKLDPK	KAYIQITFVE	YFDEYEMKDRVTYFEKNFNLRRE 1810
hCLASP3	ERFGEDVVEVIKIDSNPVDKCKLDPK	KAYIQITFVE	YFDTYEMKDRITYFDKNYNLRRE 1932
hCLASP2	DKFGSENVKMIQDSGKVNPKDLDS	YAYIOVTHVIE	FFDEKELQERKTEFERSHNLRRE 1770
hCLASP7	ERFGDDVVEIKDSYPVDKSKLDS	KAYIQITFVE	YFDTYELKDRVTYFDRNYGLRTFL 1851
hCLASP1	DKFGADNVKIIQDSNKNVPKDLDPK	YAYIOVTYVTF	FEKEIEDRKTD FEMHNNINRFV 1972
	: ** : *::** : * : .*** : **** : *	: ** : * : * : *	: * : * : *
		ITAM	DOCK motif
hCLASP4	FEAPYTLSGKKQGCIEEQKRRITILTSNSFF	YVVKRIPINCEQQIN	LKPIDGATDEIKD 1859
hCLASP5	YTPFTLEGRPERGELHEQYRRNTVLTTHAFFY	IKTRISVIQKEEFV	LTPIEVAIEDMKK 1870
hCLASP3	YCTPFTLDGRAHGEHQFQRKRTILTTHAFFY	IKTRNVNTHKEE	LTPIEVAIEDMQK 1992
hCLASP2	FEMPFTQTGKRQGGVEEQKRRITILTAHCF	FVVKRIPVMYQHHT	LNPIEVAIDEMSK 1830
hCLASP7	FCTPFTPDGRAHGEHQHQRKRTLLSTDHAF	FYIKTRIRVCHREET	VLTTPVEVAIEDMQK 1911
hCLASP1	FETPFTLSGKKHGGVAEQKRRITILTSHLF	FYVVKRIQVISQS	STLNPIEVAIDEMSR 2032
	: ** : * : * : * : * : * : *	: * : * : *	: * : * : *
		Coiled-coil	
hCLASP4	KTAELQKLCSSSTDVDMIQLQLKLQ	QWVSVQVNAGPLAYARAF	LNDSQASKYPKKVSELK 1919
hCLASP5	KTLQLAVAINQEPDPAKMLQMVLC	QSVGATVNQGP	LEVAQVFLAEIPADPKLYRHNKLR 1930
hCLASP3	KTQELAFATHQDPADPKMLQMVLC	QSVGTTVNQGP	LEVAQVFLSEIPSDPKLFRHNKLR 2052
hCLASP2	KVAELRQLCSSAEVDMIKLQLKLQ	QSVSVQVNAGPLAYARAF	LDNTNKRYPDNKVKLLK 1890
hCLASP7	KTRELAFATQEPDPAKMLQMVLC	QSVGPTVNQGP	LEVAQVFLAEIPEDPKLFRHNKLR 1971
hCLASP1	KVSELNQLCTMEEVDMISLQLKLQ	QSVSVKVNAGPMAYARAF	LEETNAKKYPDNQVKLLK 2092
	*. : *	* ** : * : *	: * : *
		Coiled-coil	
hCLASP4	DMFRKFIQACS	IALELNERLIKEDQVEYHEGLKSNFRDMVKELSD	IIHEQILQEDTMHSP 1979
hCLASP5	LCFKFIMRCGE	AVEKNKRLITADQREYQELKKNYNKLKLNLRPM	IERKIPELYKPIFR 1990
hCLASP3	LCFKDFTKRCE	DALRNKSLIGPVQREYQRELKGLSSP	----- 2090
hCLASP2	EVFRQFVEACG	QALAVNERLIKEDQLEYQEEMKANYREMAKELSE	IMHEQICPLEEKTS- 1949
hCLASP7	LCFKDFCKKCE	DALRNKALIGPDQKEYHRELENYCRLREALQPI	LTLQRLPQLMAPTP- 2030
hCLASP1	EIFRQFADACG	QALDVNERLIKEDQLEYQEELRSHYKMDLSELSTVMNE	QITGRDDLSKR 2152
	* : *	* : * : *	* : *
		PDZ ligand	
hCLASP4	WMSNTLHVFC	AISGTSSDRGYGSP	RYAEV-- 2008
hCLASP5	VESQKRDS	FHRSSFRCETQLSQS	----- 2015
hCLASP3	-----	-----	-----
hCLASP2	VLPNSLHI	FNAISGTPSTMTVHG	MTSSSSVV 1980
hCLASP7	---PGLRNSLNR	ASFRKADL	----- 2047
hCLASP1	GVDQCTRVISK	ATPALPTVIS	ISSAEV-- 2180


Human CLASP-2 expression in T cells upon activation

A)

hours post
activation



B)



7.5 kb - -

CLASP-2

2.5 kb —

28s rRNA Ethbr. staining

FIG. 14

THIS PAGE BLANK (USPTO)